

Multi-emission Kernels for Parton Branching Algorithms^a

Maximilian Löschner




Institute for Theoretical Physics (Project B1d, PI: S. Gieseke)
Erwin Schrödinger Institut (research stays in Vienna)



20 Sep 2022, Newcastle

^ain collaboration with Simon Plätzer and Emma Simpson Dore. [arXiv:2112.14454](https://arxiv.org/abs/2112.14454)



[Simon Plätzer]

$$d\sigma \simeq d\sigma_{\text{hard}}(Q) \times \text{PS}(Q \rightarrow \mu) \times \text{Had}(\mu \rightarrow \Lambda) \times \dots$$

Parton shower status

$$\int \left| \text{diagram 1} + \text{diagram 2} \right|^2 d\Phi_1 \bigg/ \int \left| \text{diagram 3} \right|^2 d\Phi_2$$

- Despite pushes for higher orders in parton showers (*e.g.* [Prestel, Hoeche—Phys.Rev.D 96 (2017) 7, 074017], [Skands, Li—PLB 771 (2017) 59-66])

Road to accuracy requires paradigm shift

- **Recoil, ordering, colour, correlations**


[Bewick, Seymour, Richardson—JHEP 04 (2020) 019], [Forshaw, Holguin, Plätzer—JHEP 09 (2020) 014], [Ruffa, Plätzer—JHEP 06 (2021) 007], [ML, Plätzer, Simpson—2112.14454], [also see PanScales]

- **Amplitude level** sets the complexity for resolving these

[Nagy, Soper], [DeAngelis, Forshaw, Plätzer—PRL 126 (2021) 11, 112001 & JHEP 05 (2018) 044]


Non-global observables

- ▶ **Coherent branching via angular ordering** essential for including large-angle soft contributions
- ▶ No global measure of deviation from jet configuration: **Coherent branching fails**
- ▶ **Dipole shower**: correct LL@LC for non-global, but issues in NLL@LC and LL@NLC for global observables



Non-global observables

- ▶ **Coherent branching via angular ordering** essential for including large-angle soft contributions
- ▶ No global measure of deviation from jet configuration: **Coherent branching fails**
- ▶ **Dipole shower**: correct LL@LC for non-global, but issues in NLL@LC and LL@NLC for global observables
- ▶ Require **dipole-type soft gluon evolution** (to account for change in colour structure)
- ▶ Even with a dipole approach, **$1/N_C$ effects possibly become comparable to subleading logs**, and intrinsically $\sim 10\%$ effects



Bucket list

⇒ Study approximations in emission iterations rather than iterations of one emission approximation.

Or: amplitude vs. cross-section level

Goal: NLL@NLC accuracy for global and non-global observables

▶ Going beyond iterated $1 \rightarrow 2$
splittings in parton showers

▶ Combine with global recoil scheme

▶ Include color and spin correlations

▶ Refine ad hoc models of MC-programs,
e.g. azimuthal correlations

▶ Define language for connecting fixed order to parton showers


} Systematic expansion
to handle uncertainties
⇔
higher logarithmic accuracy

Comparison to CS dipoles

- Catani-Seymour dipole operators reproduce the **partitioned soft and collinear behaviour** for one emission:

$$\mathcal{D}_{ij,k}(p_1, \dots, p_{m+1}) = -\frac{1}{2p_i \cdot p_j} \quad (5.2)$$

$$\cdot_m < 1, \dots, \tilde{i}, \dots, \tilde{k}, \dots, m+1 \mid \frac{\mathbf{T}_k \cdot \mathbf{T}_{ij}}{\mathbf{T}_{ij}^2} \mathbf{V}_{ij,k} \mid 1, \dots, \tilde{i}, \dots, \tilde{k}, \dots, m+1 >_m .$$



[Catani, Seymour '97]

$$\langle s | \mathbf{V}_{q_i g_j, k}(\tilde{z}_i; y_{ij, k}) | s' \rangle = 8\pi\mu^{2\epsilon} \alpha_S C_F \left[\frac{2}{1 - \tilde{z}_i(1 - y_{ij, k})} - (1 + \tilde{z}_i) - \epsilon(1 - \tilde{z}_i) \right] \delta_{ss'}$$


- Our idea: **algorithmic generation of such splitting kernels for > 1 emission**
- Generate partitioned soft behaviour via **power counting** instead of construction ‘by hand’
- Potential for constructing **subtraction terms**

Splitting kernels

Splitting kernels from amplitudes

From the cross-section level to decomposed amplitudes:


$$\sigma = \sum_n \int \text{Tr} [|\mathcal{M}(\mu)\rangle \langle \mathcal{M}(\mu)|] u(p_1, \dots, p_n) d\phi_n$$



Splitting kernels from amplitudes


From the cross-section level to decomposed amplitudes:

$$\sigma = \sum_n \int \text{Tr} [|\mathcal{M}(\mu)\rangle \langle \mathcal{M}(\mu)|] u(p_1, \dots, p_n) d\phi_n$$



Splitting kernel iterations

Density operator language is useful for discussing emissions in iterative manner:



[Forshaw, Holguin, Plätzer–JHEP 09 (2020) 014]

Partitioning


Disentangling different collinear sectors

- Use partition of one in terms of all possible collinear pairings

$$1 = \mathbb{P}_1^{(\mathcal{A})} + \mathbb{P}_2^{(\mathcal{A})} + \mathbb{P}_3^{(\mathcal{A})} + \dots$$

where $\mathbb{P}_i^{(\mathcal{A})}$ projects onto collinearity w.r.t. p_i for some amplitude \mathcal{A}

- **Disentangle overlapping collinear singularities**
- Keep smooth interpolation over whole phase space




Angular ordering and subtractions

- Radiation of a soft gluon leads to

$$d\sigma_{n+1} = d\sigma_n \times \frac{d\omega}{\omega} \frac{d\Omega}{2\pi} \frac{\alpha_S}{2\pi} \sum_{i,l} C_{il} W_{il}$$


where $W_{il} = \frac{\omega^2 p_i \cdot p_l}{p_i \cdot p_j p_l \cdot p_j} :$ 'Radiation function'



Angular ordering and subtractions

- Radiation of a soft gluon leads to

$$d\sigma_{n+1} = d\sigma_n \times \frac{d\omega}{\omega} \frac{d\Omega}{2\pi} \frac{\alpha_S}{2\pi} \sum_{i,l} C_{il} W_{il}$$



where $W_{il} = \frac{\omega^2 p_i \cdot p_l}{p_i \cdot p_j p_l \cdot p_j}$: 'Radiation function'

- Can decompose $W_{il} = W_{il}^{[i]} + W_{il}^{[l]}$

$$W_{il}^{[i]} = \frac{1}{2} \left(W_{il} - \frac{1}{1 - \cos \theta_{jl}} + \frac{1}{1 - \cos \theta_{ij}} \right)$$

- Then azimuthal averaging confines emissions to cone

$$\int_0^{2\pi} \frac{d\phi_{ij}}{2\pi} W_{il}^{[i]} = \begin{cases} \frac{1}{1 - \cos \theta_{ij}} & \text{if } \theta_{ij} < \theta_{il}, \\ 0 & \text{otherwise.} \end{cases}$$

- Textbook knowledge: subtraction partitioning implies angular ordering [Ellis, Stirling, Webber]

Subtraction partitioning

- ▶ As an alternative to fractional partitioning, define **subtraction scheme**:

$$\mathbb{P}_{(i||j)} \left[\frac{1}{S_{ij} S_{jl}} \right] = \frac{1}{2} \left(\frac{1}{S_{ij} S_{jl}} - \Delta_{(j||l)} + \Delta_{(i||j)} \right),$$

$$\mathbb{P}_{(j||l)} \left[\frac{1}{S_{ij} S_{jl}} \right] = \frac{1}{2} \left(\frac{1}{S_{ij} S_{jl}} - \Delta_{(i||j)} + \Delta_{(j||l)} \right),$$

$$\Delta_{(i||j)} = \frac{E_i}{E_j} \frac{1}{S_{il} S_{ij}}, \quad \Delta_{(j||l)} = \frac{E_l}{E_j} \frac{1}{S_{il} S_{jl}}.$$


by exploiting $S_{ij} \xrightarrow{(j||l)} E_i E_j n_i \cdot n_l = \frac{E_j}{E_l} S_{il}$

- ▶ $\mathbb{P}_{(i||j)} [\dots]$ non-singular in $(j || l)$ -limit while original singular behaviour is reproduced in $(i || j)$ -limit
- ▶ **Algorithmic generalisation to multi emissions under control**

Subtraction partitioning behaviour

Recent work: subtraction partitioning \Rightarrow angular ordering for 2E?

$$\mathcal{A} \propto \frac{1}{S_{ij}S_{jl}} :$$





Subtraction partitioning behaviour


Recent work: subtraction partitioning \Rightarrow angular ordering for 2E?

$$\mathcal{A} \propto \frac{1}{S_{ij} S_{jl}} :$$

$$\mathcal{A} \propto \frac{1}{S_{ij} S_{jk} S_{jl} S_{kl}} :$$



partitioning
(i||j)(k||l)



Fractional partitioning for two emissions

Alternatively: cancel out ‘unwanted’ collinear singularities by partitioning factors

- Read
 $(i \parallel j \parallel k) : S_{ijk} = (q_i + q_j + q_k)^2 \rightarrow 0$
- **Collect non-singular factors** in triple
collinear and coll-coll pairings

configuration	$\mathcal{A} \propto \frac{1}{S_{ij} S_{kl} S_{ijk} S_{jkl}}$
$i \parallel j \parallel k$	$S_{kl} S_{jkl}$
$i \parallel j \parallel l$	$S_{kl} S_{ijk} S_{jkl}$
$i \parallel k \parallel l$	$S_{ij} S_{ijk} S_{jkl}$
$j \parallel k \parallel l$	$S_{ij} S_{ijk}$
$(i \parallel j), (k \parallel l)$	$S_{ijk} S_{jkl}$
$(i \parallel k), (j \parallel l)$	\times
$(i \parallel l), (j \parallel k)$	\times

Fractional partitioning for two emissions

Alternatively: cancel out ‘unwanted’ collinear singularities by partitioning factors

- Read
 $(i \parallel j \parallel k) : S_{ijk} = (q_i + q_j + q_k)^2 \rightarrow 0$
- **Collect non-singular factors** in triple
collinear and coll-coll pairings

configuration	$\mathcal{A} \propto \frac{1}{S_{ij} S_{kl} S_{ijk} S_{jkl}}$
$i \parallel j \parallel k$	$S_{kl} S_{jkl}$
$i \parallel j \parallel l$	$S_{kl} S_{ijk} S_{jkl}$
$i \parallel k \parallel l$	$S_{ij} S_{ijk} S_{jkl}$
$j \parallel k \parallel l$	$S_{ij} S_{ijk}$
$(i \parallel j), (k \parallel l)$	$S_{ijk} S_{jkl}$
$(i \parallel k), (j \parallel l)$	\times
$(i \parallel l), (j \parallel k)$	\times

⇒ Construct partitioning factors of the form

$$\mathbb{P}_{(ijk)}^{(\mathcal{A})} = \frac{S_{kl} S_{jkl}}{S_{kl} S_{jkl} + S_{ij} S_{ijk} + S_{ijk} S_{jkl} + (S_{kl} + S_{ij}) S_{ijk} S_{jkl}}$$

- $\mathbb{P}_{(ijk)}^{(\mathcal{A})} \times \mathcal{A}$ **extracts the $(i \parallel j \parallel k)$ - singular behaviour**
- $\mathbb{P}_{(ijk)}^{(\mathcal{A})}$ **is non-singular in any collinear configuration**

Power Counting

Power counting

- Discuss soft and collinear **scaling of internal lines in general way**
- Sudakov-like decomposition of momenta:

$$q_I^\mu = \sum_{k \in I} r_{ik} = z_I \textcolor{red}{p}_i^\mu + \frac{S_I + p_{\perp,I}^2}{2z_I p_i \cdot n} n^\mu + \textcolor{red}{k}_{\perp,I}^\mu ,$$

- Decompose fermion and gluon lines (factors of $\sqrt{z_I}$ absorbed in vertices for fermions):

$$\circ \rightarrow \square \rightarrow \circ = \not{p}_i ,$$

$$\text{wavy} \square \text{wavy} = d^{\mu\nu}(p_i) ,$$

$$\circ \rightarrow \blacksquare \rightarrow \circ = \frac{S_I + p_{\perp,I}^2}{2z_I^2 p_i \cdot n} \not{n} ,$$

$$\text{wavy} \blacksquare \text{wavy} = \frac{S_I + p_{\perp,I}^2}{(z_I p_i \cdot n)^2} n^\mu n^\nu ,$$

$$\circ \rightarrow \square_{\perp} \rightarrow \circ = \frac{\not{k}_{\perp,I}}{z_I} ,$$



$$\text{wavy} \square_{\perp} \text{wavy} = \frac{k_{\perp,I}^\mu n^\nu + n^\mu k_{\perp,I}^\nu}{z_I p_i \cdot n} .$$

- Leads to **power counting rules** with potential connection to SCET



Soft and collinear scaling

- Algorithmically determine soft or collinear scaling of an emission amplitude via scaling of internal lines (and propagators)

Scaling of hard lines:

	h	h+c	h+s	h+c+s
	λ	λ	λ	λ (bal.)
	0	λ	λ	λ (unbal.)
	λ^2	λ^2	λ	λ (bal.)
	0	λ^2	λ	λ (unbal.)

Scaling of emissions:

	s	c	s+c
	1	λ	λ
	1	λ^2	λ

- Note differences between mappings, e.g. with and without balanced k_{\perp} -components

One emission amplitudes

- Determine list of all **relevant sub-amplitudes** via power counting rules
- Combine these in density operator (\simeq squared amp) to find **full splitting kernel**

	C	S		C	S
	λ	λ		λ	λ
	λ	1		λ^2	λ^2
	λ	λ		λ^2	λ^2
	λ^2	λ		λ^2	λ
	λ^2	λ^2		1	1
	λ^2	λ		λ	1

One emission example

Full one emission (ij) -splitting kernel (balanced mapping) consists of

$$\mathbb{U}_{(ij)} = \mathbb{P}_{(ij)} \left(\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \\ \text{Diagram 4} \\ \text{Diagram 5} \\ \text{Diagram 6} \end{array} \right)$$

The diagrams represent different ways a gluon emission can be attached to a quark line in a splitting process. Each diagram shows two incoming quark lines (left) and two outgoing quark lines (right), with a central vertical bar representing the splitting vertex. The diagrams are arranged in a 3x2 grid, with the first two diagrams in the first row enclosed in large parentheses. The diagrams show various configurations of gluon emission (represented by curly lines) and quark lines (represented by straight lines) connected by vertices (squares).


- ▶ Exhibits factorisation to hard amplitude
- ▶ Smooth interpolation between soft and collinear limits
- ▶ Algorithmically generalizable for more emissions

Balanced vs. unbalanced mapping

- Can test different implementations of momentum mappings, e.g. the balancing of transverse components

$$\mathbf{k}_{\perp, I}^{\mu} = \sum_{i \in I} \mathbf{k}_{\perp, I}^{(i), \mu},$$


- Yields different sets of diagrammatic contributions
- Nevertheless, the same collinear and soft behaviour is reproduced for one emission



- Still: can compare mappings and check for inconsistencies for > 1 emission


Check: One emission splitting function

- Reproduce **Splitting function** P_{qg} as a crosscheck



Check: One emission splitting function


- Reproduce **Splitting function** P_{qg} as a crosscheck



$$\rightarrow \frac{4\pi\alpha_s \mathbf{T}_i^2}{S_{ij}} \left[(d-2)\alpha_i + 4\frac{(1-\alpha_i)^2}{\alpha_i} + 4(1-\alpha_i) \right] \not{p}_i + \mathcal{O}(\lambda^{-1}).$$


Soft-Collinear Interplay

- Soft singular part of splitting function cancelled by:



$$\propto \frac{4\pi\alpha_s \mathbf{T}_i \cdot \mathbf{T}_k}{S_{jk}} \frac{4(1 - \alpha_i)^2}{\alpha_i} \frac{p_k \cdot n}{p_i \cdot n} [\not{p}_i][\not{p}_k]$$

- Eikonal part remains:







$$\propto \frac{4\pi\alpha_s \mathbf{T}_i \cdot \mathbf{T}_k}{S_{ij}S_{jk}} 4(1 - \alpha_i) \sqrt{\frac{\tilde{\beta}_i}{\alpha_i}} p_k \cdot n_{\perp} [\not{p}_i][\not{p}_k]$$

- Smooth interpolation between soft and collinear limits in $\mathbb{U}_{(ij)}$
- Current work: investigate this interplay for two emissions

Two emissions: splitting amplitudes


- ▶ Same procedure applies to two emissions
- ▶ Some amplitudes can **not be achieved by single emission iteration**
- ▶ Signals for **violation of exact factorisation** (drop out for two emissions though)

	$C_1 C_2$	$C_1 S_2$	$S_1 C_2$	$S_1 S_2$
	λ^2	λ	λ^2	λ
	λ^2	λ	λ	1
	λ^2	λ^2	λ^2	λ^2
	λ^3	λ^2	λ	λ
⋮				

Two emissions: combined contributions

- Determine amplitude scaling **algorithmically**:

`combinedAmpsB2[[c, s, c, s], 1]`



- Combine with partitioned propagator scaling to find all leading contributions for full kernel

	CC	CS	SC	SS
$A^{(1)}$	$1/\lambda^2$	$1/\lambda^3$	$1/\lambda$	$1/\lambda^4$
$A^{(2)}$	$1/\lambda^2$	$1/\lambda^2$	$1/\lambda^2$	$1/\lambda^4$
$A^{(3)}$	$1/\lambda^2$	$1/\lambda^2$	$1/\lambda$	$1/\lambda^3$
$A^{(4)}$	$1/\lambda^3$	$1/\lambda^2$	$1/\lambda$	$1/\lambda^4$
$A^{(5)}$	$1/\lambda^3$	$1/\lambda$	$1/\lambda$	$1/\lambda^4$
$B^{(1)}$	$1/\lambda^3$	$1/\lambda^4$	$1/\lambda^2$	$1/\lambda^4$
$B^{(2)}$	$1/\lambda^3$	$1/\lambda^3$	$1/\lambda^3$	$1/\lambda^4$
$B^{(3)}$	$1/\lambda^3$	$1/\lambda^3$	$1/\lambda^2$	$1/\lambda^4$
$B^{(4)}$	$1/\lambda^3$	$1/\lambda^4$	$1/\lambda^2$	$1/\lambda^4$
$B^{(5)}$	$1/\lambda^3$	$1/\lambda^3$	$1/\lambda^3$	$1/\lambda^4$
$B^{(6)}$	$1/\lambda^3$	$1/\lambda^3$	$1/\lambda^2$	$1/\lambda^4$
$X^{(1)}$	$1/\lambda^2$	$1/\lambda^2$	$1/\lambda^4$	$1/\lambda^4$
$X^{(2)}$	$1/\lambda^2$	$1/\lambda^2$	$1/\lambda^2$	$1/\lambda^2$
$E^{(1)}$	$1/\lambda^4$	$1/\lambda^4$	$1/\lambda^2$	$1/\lambda^4$
$E^{(2)}$	$1/\lambda^4$	$1/\lambda^3$	$1/\lambda^3$	$1/\lambda^4$
$E^{(3)}$	$1/\lambda^4$	$1/\lambda^3$	$1/\lambda^2$	$1/\lambda^4$
$E^{(4)}$	$1/\lambda^4$	$1/\lambda^3$	$1/\lambda^2$	$1/\lambda^4$
$E^{(5)}$	$1/\lambda^4$	$1/\lambda^2$	$1/\lambda^2$	$1/\lambda^4$

Conclusions

Goal: **universal algorithm for handling accuracy in multiple emissions** (for applications in parton showers and beyond)

- ▶ Density-operator formalism to study iterative behaviour of emissions
- ▶ Partitioning algorithms to separate overlapping singularities
- ▶ Momentum mapping for exposing collinear and soft factorization
- ▶ Global recoil via Lorentz transformation
- ▶ Set of power counting rules to single out leading amplitudes
- ▶ Can handle and compare different momentum mappings
- ▶ Two-emission kernels/power counting under control

Conclusions

Goal: **universal algorithm for handling accuracy in multiple emissions** (for applications in parton showers and beyond)


- ▶ Density-operator formalism to study iterative behaviour of emissions
- ▶ Partitioning algorithms to separate overlapping singularities
- ▶ Momentum mapping for exposing collinear and soft factorization
- ▶ Global recoil via Lorentz transformation
- ▶ Set of power counting rules to single out leading amplitudes
- ▶ Can handle and compare different momentum mappings
- ▶ Two-emission kernels/power counting under control

Thank you!

Backup slides

Coherent branching

- ▶ **Coherent emission of soft large angle gluons** from systems of collinear partons
- ▶ **Angular ordering** essential for including large-angle soft contributions




[Simon Plätzer]

- ▶ Resummation of global jet observables such as **thrust τ**
- ▶ **NLL accurate @Next-to-Leading-Colour (NLC)** if inclusive over secondary soft gluon emission

Applications

- Use projectors and helicity sums to represent emission amplitudes as **(complex) weights for numerical evaluation**


$$\mathbf{P}(q) \equiv \begin{cases} P^{\rho\sigma}(p) = d^{\rho\sigma}(p), & (\text{gluon}), \\ \not{p}(p) = \frac{\not{p}}{2n \cdot p}, & (\text{quark}), \end{cases} \quad \begin{aligned} d^{\mu\nu}(p) &= \epsilon_+^\mu(p, n) \epsilon_-^\nu(p, n) + (\mu \leftrightarrow \nu), \\ \not{p} &= \sum_\lambda u_\lambda(n) \bar{u}_\lambda(n), \end{aligned}$$



Applications

- Use projectors and helicity sums to represent emission amplitudes as **(complex) weights for numerical evaluation**

$$\mathbf{P}(q) \equiv \begin{cases} P^{\rho\sigma}(p) = d^{\rho\sigma}(p), & (\text{gluon}), \\ \not{p}(p) = \frac{\not{p}}{2n \cdot p}, & (\text{quark}), \end{cases} \quad \begin{aligned} d^{\mu\nu}(p) &= \epsilon_+^\mu(p, n) \epsilon_-^\nu(p, n) + (\mu \leftrightarrow \nu), \\ \not{p} &= \sum_\lambda u_\lambda(n) \bar{u}_\lambda(n), \end{aligned}$$




$$\begin{aligned} &\rightarrow \sum_{\lambda_i, \bar{\lambda}_i} \frac{u_{\lambda_1}}{\sqrt{2n \cdot p_i}} \left[\frac{\bar{u}_{\lambda_1}}{\sqrt{2n \cdot p_i}} \not{k}_\perp \not{\epsilon}_{\lambda_3} \not{p}_i \frac{u_{\lambda_2}}{\sqrt{2n \cdot p_i}} \right] \frac{\bar{u}_{\lambda_2}}{\sqrt{2n \cdot p_i}} \epsilon_{\lambda_3}^\sigma \\ &\times \frac{u_{\bar{\lambda}_1}}{\sqrt{2n \cdot p_i}} \left[\frac{\bar{u}_{\bar{\lambda}_1}}{\sqrt{2n \cdot p_k}} \not{p}_k \frac{u_{\bar{\lambda}_2}}{\sqrt{2n \cdot p_k}} p_k \cdot \epsilon_{\bar{\lambda}_3} \right] \frac{\bar{u}_{\bar{\lambda}_2}}{\sqrt{2n \cdot p_k}} \epsilon_{\bar{\sigma}, \bar{\lambda}_3}. \end{aligned}$$

Team

Karlsruhe/Manchester/Vienna network with support from **SFB** drives significant parts of the development, also relating to aspects such as **color reconnection** [e.g. Gieseke, Kirchgaesser, Plätzer-JHEP 11 (2018)

149]



Algorithm for subtraction partitioning

- General form of **partitioned propagator** P for config σ


$$\mathbb{P}_{\sigma}[P] = \frac{1}{m} \left(P + (m-1) \Delta_{\sigma; \tau_1, \dots, \tau_{m-1}}[P] - \sum_{i=1}^{m-1} \Delta_{\tau_i; \tau_1, \dots, \tau_{i-1}, \sigma, \tau_{i+1}, \dots, \tau_{m-1}}[P] \right),$$

- with **Subtraction terms**

$$\Delta_{\tau_1; \tau_2, \dots, \tau_m}[P] = \underbrace{\mathbb{F}_{\tau_1}[P]}_{\text{non-singular bits}} \left(\underbrace{\mathbb{S}_{\tau_1}[P]}_{\text{singular bits}} - \sum_{S/\tau_1} \Delta_{\tau_{i_1}; \tau_{i_2}, \dots, \tau_{i_{m-1}}}[\mathbb{S}_{\tau_1}[P]] \right),$$

- When partitioning e.g. to $\sigma = (i \parallel j \parallel k)$, subtract off all (sub-)divergences of other singular configs τ_i for propagator factor P .
- Combinatorial factor m : number of singular configs for P

Two emission example



► Partitioned version of $A^{(1)} \propto 1/S_{ij}S_{ijk}S_{kl}S_{jkl}$

$$\begin{aligned} \mathcal{P}(A^{(1)}) = & \frac{1}{3} \left(\frac{1}{S_{ij}S_{ijk}S_{kl}S_{jkl}} + 2\Delta_{(ijk)}[\mathcal{P}(A^{(1)})] - \Delta_{(jkl)}[\mathcal{P}(A^{(1)})] - \Delta_{(ij)(kl)}[\mathcal{P}(A^{(1)})] \right), \\ & + \frac{1}{3} \left(\frac{1}{S_{ij}S_{ijk}S_{kl}S_{jkl}} - \Delta_{(ijk)}[\mathcal{P}(A^{(1)})] + 2\Delta_{(jkl)}[\mathcal{P}(A^{(1)})] - \Delta_{(ij)(kl)}[\mathcal{P}(A^{(1)})] \right), \\ & + \frac{1}{3} \left(\frac{1}{S_{ij}S_{ijk}S_{kl}S_{jkl}} - \Delta_{(ijk)}[\mathcal{P}(A^{(1)})] - \Delta_{(jkl)}[\mathcal{P}(A^{(1)})] + 2\Delta_{(ij)(kl)}[\mathcal{P}(A^{(1)})] \right), \end{aligned}$$

where e.g.

$$\Delta_{(jkl)}[\mathcal{P}(A^{(1)})] = \frac{E_l^2}{E_j(E_j + E_k)} \frac{1}{S_{il}^2} \left(\frac{1}{S_{kl}S_{jkl}} - \frac{E_i E_l}{E_j(E_l + E_k)} \frac{1}{S_{il}S_{kl}} \right),$$

Check: Two Emissions



- **Reproduced from general two-emission kernel** which includes soft-limit too (here: in lightcone-gauge)

$$\begin{aligned}
 & \frac{\mu^{2\varepsilon}}{\hat{\alpha}^2 S_{i12}^2} \left\{ \begin{array}{l} \text{Diagram 1} + \text{Diagram 2} + \text{Diagram 3} + \text{Diagram 4} + (1 \leftrightarrow 2) \end{array} \right\} C_A C_F \\
 &= \left(\frac{8\pi\alpha_S}{\hat{\alpha} S_{i12}} \mu^\varepsilon \right)^2 C_A C_F \langle \hat{P}_{ggq}^{(\text{non-Ab})} \rangle \hat{p}_i + \mathcal{O}(\beta_{il}^{-3/2}).
 \end{aligned}$$

The diagrams represent two-emission kernels in lightcone-gauge. Each diagram shows a horizontal line with two external legs (shaded ellipses) and two internal legs (labeled \hat{p}_i). The internal legs are connected by a loop of gluons (curly lines). The diagrams are arranged in a 2x2 grid, with the top row containing two diagrams and the bottom row containing two diagrams. The diagrams are labeled with indices 1, 2, and i . The first diagram in the top row has a loop with a dashed line labeled i and a solid line labeled 1 . The second diagram in the top row has a loop with a dashed line labeled 2 and a solid line labeled 1 . The third diagram in the bottom row has a loop with a dashed line labeled 2 and a solid line labeled 1 . The fourth diagram in the bottom row has a loop with a dashed line labeled 2 and a solid line labeled 1 . The diagrams are summed and multiplied by $C_A C_F$ to give the final result.


Vertex rules

- Can find vertex rules such as:



Insights from Power Counting Rules

- Powerful vertex rule for lines belonging to same collinear sector:




A Feynman diagram representing a three-point vertex. It consists of a central vertex from which three lines extend outwards. The top-left line extends upwards and to the left, the top-right line extends upwards and to the right, and the bottom line extends straight downwards. Each of these three lines passes through a square box labeled p_i . The diagram is followed by an equals sign and a zero, indicating that this diagram evaluates to zero.

$$= 0,$$


Insights from Power Counting Rules

- Powerful vertex rule for lines belonging to same collinear sector:




$$= 0,$$

- Shows (known fact) that interference diagrams do not contribute in splitting function in a physical gauge
- **Reason:** denominator goes as $1/\lambda^{2k} S_{(\text{col})}^k$ for k coll. emissions
- Can only contribute in splitting function ($\propto 1/\lambda^{2k} S_{(\text{col})}^k$) if numerator goes as $\mathcal{O}(1)$, but the **only possible contribution** $\equiv 0$



$$= 0.$$

Global and non-global observables



[Dasgupta, Salam (2001)]

- Example: heavy and light jet mass (global) vs. hemisphere jet mass (non-global)
- Cancellations between large angle-soft and virtual contributions (from k_2) not guaranteed
⇒ **NLL enhancement from leftover $\alpha_S^2 L^2$ terms**

Partitioning

Amplitudes carry different singular S -invariants

$$\mathcal{A}(S_1, S_2) = \frac{\mathcal{N}(S_1, S_2)}{S_1 S_2},$$


Decomposition using partitioning factors:

$$\mathbb{P}_{(1)}^{(\mathcal{A})} = \frac{S_2}{S_1 + S_2}, \quad \mathbb{P}_{(2)}^{(\mathcal{A})} = \frac{S_1}{S_1 + S_2},$$

we can decompose \mathcal{A} into

$$\mathcal{A} = \left[\mathbb{P}_{(1)}^{(\mathcal{A})} + \mathbb{P}_{(2)}^{(\mathcal{A})} \right] \mathcal{A} = \frac{\mathcal{N}(S_1, S_2)}{S_1(S_1 + S_2)} + \frac{\mathcal{N}(S_1, S_2)}{S_2(S_1 + S_2)}.$$

Parton Shower



- **Soft** and **collinear** regions are of special interest:

$$S_{ij} \equiv (q_i + q_j)^2 = 2 q_i \cdot q_j = 2 q_i^0 q_j^0 [1 - \cos \theta_{ij}], \quad \text{for } q_{i/j}^2 = 0$$

- Amplitude goes as $\propto 1/S_{ij}$
 \Rightarrow becomes singular/enhanced when $S_{ij} \rightarrow 0$
- **Large logarithms** due to phase space integrations of the kind

$$\frac{dq_j^0}{q_j^0}, \quad \frac{d\theta_{ij}}{\theta_{ij}} \rightarrow \alpha_S \log^2 \frac{Q}{Q_0} \sim 1$$

for some scale $Q \in \{\theta, p_\perp, \dots\}$ and cut-off Q_0

Parton shower: collinear limit

- Single emission approach is then usually iterated in a probabilistic manner

$$W_{2+2} = \left(\int \left| \text{diagram}_1 \right|^2 + \left| \text{diagram}_2 \right|^2 + \left| \text{diagram}_3 \right|^2 + \left| \text{diagram}_4 \right|^2 d\Phi_2 \right) / \left| \text{diagram}_5 \right|^2$$

$$= 2^2 \int_{t_0}^t dt' \int_{t_0}^{t'} dt'' W(t') W(t'') = \frac{2^2}{2!} \left(\int_{t_0}^t dt W(t) \right)^2.$$

[Stefan Gieseke]

- Sum over any number of emissions: result exponentiates

$$\sigma_{>2}(t_0) = \sigma_2(t_0) \sum_{k=1}^{\infty} \frac{2^k}{k!} \left(\int_{t_0}^t dt W(t) \right)^k$$


- Sudakov Form Factor** (\simeq no emission probability in range $t \rightarrow t_0$)

$$\Delta(t_0) = \exp \left[- \int_{t_0}^t dt W(t) \right], \quad W(t) = \int_{z_-}^{z_+} \frac{\alpha_S(z, t)}{2\pi} \frac{\hat{P}(z, t)}{t} dz.$$

Momentum mapping

Momentum mapping


Adding emissions



- Start with **on-shell** (OS) momenta p_i (to be **emitters**) and p_r (to be **recoilers**) with overall momentum transfer $Q \equiv \sum_i p_i + \sum_r p_r$

Momentum mapping

Adding emissions



- Start with **on-shell** (OS) momenta p_i (to be **emitters**) and p_r (to be **recoilers**) with overall momentum transfer $Q \equiv \sum_i p_i + \sum_r p_r$
- Add emissions to the process with:
 1. Momentum conservation: $\sum_i q_i + \sum_{i,l} k_{il} + \sum_r q_r = Q$
 2. On-shellness of all partons
 3. Parametrization of soft & collinear behaviour for any # of emissions

Momentum mapping

$$q_r = \frac{\Lambda}{\alpha_L} p_r$$

$$k_{il} = \frac{\Lambda}{\alpha_L} \left[\alpha_{il} p_i + \tilde{\beta}_{il} n_i + \sqrt{\alpha_{il} \tilde{\beta}_{il}} n_{il}^\perp \right], \quad A_i \equiv \sum_l \alpha_{il}, \quad \tilde{\beta}_{il} = (1 - A_i) \beta_{il}$$

$$q_i = \frac{\Lambda}{\alpha_L} \left[(1 - A_i) p_i + (y_i - \sum_l \tilde{\beta}_{il}) n_i - \sum_l \sqrt{\alpha_{il} \tilde{\beta}_{il}} n_{il}^\perp \right]$$

- Decomposition w/ light-like momentum n_i and $n_{il}^\perp \cdot p_i = n_{il}^\perp \cdot n_i = 0$
- Need $\alpha_L^2 = (Q + N)^2 / Q^2$ for momentum conservation

$$Q = \sum_r q_r + \sum_i q_i + \sum_{i,l} k_{il} = \frac{\Lambda}{\alpha_L} \left[\underbrace{\sum_r p_r}_Q + \underbrace{\sum_i (p_i + y_i n_i)}_N \right]$$

- Lorentz transformation $\Lambda, \alpha_L \Rightarrow$ non-trivial **global recoil**

Momentum mapping II

- Using Λ and α_L , recoil effects are removed from considerations about factorization, due to Lorentz invariance and known mass dimension of the amplitudes:

$$|\mathcal{M}(q_1, \dots, q_n)\rangle = \frac{1}{\alpha_L^{2n-4}} |\mathcal{M}(\hat{q}_1, \dots, \hat{q}_n)\rangle .$$

- Soft and collinear power counting possible via scaling of α_{il} and β_{il} , i.e. $(p_i, n_i, n_{i\perp}^\perp)$ -components

	$(\alpha_{il}, y_i, \beta_{il})$
(forward) collinear	$(1, \lambda^2, \lambda^2)$
soft	$(\lambda, \lambda, \lambda)$.

- Facilitates study of an amplitude's singular behaviour for implementation in splitting kernels
- This mapping is just one possible instance. Can e.g. use different balancing of transverse components.

Two emissions: topologies

- Decompose squared amplitude in terms of set of topologies

$$|\mathcal{M}_{n+2}|^2 = \sum_i \sum_{\alpha} \left(E_{ijk}^{(\alpha)} + (j \leftrightarrow k) \right) \\ + \sum_i \sum_{l \neq i} \sum_{\alpha} \left(A_{ijkl}^{(\alpha)} + B_{ijkl}^{(\alpha)} + X_{ijkl}^{(\alpha)} + (j \leftrightarrow k) \right) + \dots$$

- Examples:

