

Two-loop amplitude for heavy-quark pair production via light-quark pair annihilation (and related processes)

Jonathan Ronca

In collaboration with:


*Bonciani, Broggio, Di Vita, Ferroglio, Laporta, Mastrolia, Mandal,
Mattiuzzi, Passera, Primo, Schubert, Torres Bobadilla, and Tramontano*

8th International Workshop on High Precision for Hard Processes

Discovery Museum, Newcastle-upon-Tyne
20 - 22 September 2022

20 Sept. 2022

Motivation :: Muon ($g - 2$)



[Muon g-2 Collaboration (2021)]

$$a_\mu^{\text{EXP}} = (116592089 \pm 63) \times 10^{-11} [0.54\text{ppm}] \quad \text{BNL E821}$$

$$a_\mu^{\text{EXP}} = (116592040 \pm 54) \times 10^{-11} [0.46\text{ppm}] \quad \text{FNAL E989 Run 1}$$

$$a_\mu^{\text{EXP}} = (116592061 \pm 41) \times 10^{-11} [0.35\text{ppm}] \quad \text{WA}$$

Standard Model prediction

$$a_\mu^{\text{SM}} = 116591810 (43) \times 10^{-11}$$

Significant discrepancy
between the **experiment** and the
SM prediction

$$\Delta a_\mu = a_\mu^{\text{EXP}} - a_\mu^{\text{SM}} = 251 (59) \times 10^{-11}$$

Motivation :: Muon ($g - 2$)

$$a_\mu^{\text{QED}} = 116584718.931 (19)(100)(23) \times 10^{-11}$$


$$a_\mu^{\text{EW}} = 153.6 (1.0) \times 10^{-11}$$

$$a_\mu^{\text{HLO}} = 6931 (40) \times 10^{-11}$$

Affected by the **largest** theoretical error


To extract $\Delta\alpha_{\text{had}}(t)$ from MUonE's measurement, the ratio of the SM cross sections in the signal and normalisation regions must be known at $\lesssim 10\text{ppm}$!

Large QED background



High-precision calculation of
electron-muon elastic scattering in QED


- Leading hadronic contribution computed via the usual dispersive (timelike) formula:



$$a_\mu^{\text{HLO}} = \frac{1}{4\pi^3} \int_{m_\pi^2}^\infty ds K(s) \sigma_{\text{had}}^{(0)}(s)$$

$$K(s) = \int_0^1 dx \frac{x^2 (1-x)}{x^2 + (1-x) (s/m_\mu^2)}$$

- Alternatively, simply exchanging the x and s integrations:



$$a_\mu^{\text{HLO}} = \frac{\alpha}{\pi} \int_0^1 dx (1-x) \Delta\alpha_{\text{had}}[t(x)]$$

$$t(x) = \frac{x^2 m_\mu^2}{x-1} < 0$$

Lautrup, Peterman, de Rafael, 1972

$\Delta\alpha_{\text{had}}(t)$ is the hadronic contribution to the running of α in the spacelike region: measure a_μ^{HLO} via scattering data!

Motivation :: Muon ($g - 2$)

$$a_\mu^{\text{QED}} = 116584718.931 (19)(100)(23) \times 10^{-11}$$

$$a_\mu^{\text{EW}} = 153.6 (1.0) \times 10^{-11}$$

$$a_\mu^{\text{HLO}} = 6931 (40) \times 10^{-11}$$

Affected
theory


To extract $\Delta\alpha_{\text{had}}(t)$ from MU cross sections in the signal known at $\lesssim 10\text{ppm}$!

Large QED background



High-precision calculation of
electron-muon elastic scattering in QED

- Leading hadronic contribution computed via the usual dispersive (timelike) formula:



$$a_\mu^{\text{HLO}} = \frac{1}{4\pi^3} \int_{m_\pi^2}^\infty ds K(s) \sigma_{\text{had}}^{(0)}(s)$$

$$= \int_0^1 dx \frac{x^2 (1-x)}{x^2 + (1-x) (s/m_\mu^2)}$$

The x and s integrations:


$$= \frac{\alpha}{\pi} \int_0^1 dx (1-x) \Delta\alpha_{\text{had}}[t(x)]$$

$$\frac{x^2 m_\mu^2}{x-1} < 0$$

Lautrup, Peterman, de Rafael, 1972

to the running of α in the
scattering data!

Calame, MP, Trentadue, Venanzoni, PLB 2015



[Abbiendi, Carloni Calame, Marconi, Matteuzzi,
Montagna, Nicrosini, Passera, Piccinini,
Tenchini, Trentadue, Venanzoni (2017)]

Cross Section and Scattering Amplitudes in pQFT

Target observable: **cross section**

$$\sigma(1 \rightarrow 2) = \alpha^2 \left[\sum_{j=0}^n \left(\frac{\alpha}{\pi} \right)^j \sigma_{\text{N}^j \text{LO}} + O(\alpha^{n+1}) \right]$$

related to the **scattering amplitude**

$$\mathcal{A} = 4\pi\mu^{-2\epsilon} \sum_j \left(\frac{\alpha}{\pi} \right)^j \mathcal{A}_j$$

σ_{LO}

$$\int \sum_{ij} \left[\begin{array}{c} \text{tree} \\ \text{tree} \end{array} \right] d\text{PS}_2$$


σ_{NLO}

$$\int \sum_{ij} \left[\begin{array}{c} \text{Virtual} \\ 1\text{L} \end{array} \right] d\text{PS}_2$$

σ_{NNLO}

$$\int \sum_{ij} \left[\begin{array}{c} \text{VV} \\ 2\text{L} \end{array} \right] d\text{PS}_2$$

$$\int \sum_{ij} \left[\begin{array}{c} \text{RV} \\ 1\text{L} \end{array} \right] d\text{PS}_2$$



Feynman
diagrams

$$\int \sum_{ij} \left[\begin{array}{c} \text{Real} \\ \text{tree} \end{array} \right] d\text{PS}_3$$

$$\int \sum_{ij} \left[\begin{array}{c} \text{RV} \\ 1\text{L} \end{array} \right] d\text{PS}_3$$

$$\int \sum_{ij} \left[\begin{array}{c} \text{RR} \\ \text{tree} \end{array} \right] d\text{PS}_4$$


Cross Section and Scattering Amplitudes in pQFT

Target observable: **cross section**

$$\sigma(1 \rightarrow 2) = \alpha^2 \left[\sum_{j=0}^n \left(\frac{\alpha}{\pi} \right)^j \sigma_{\text{N}^j \text{LO}} + O(\alpha^{n+1}) \right]$$

related to the **scattering amplitude**

$$\mathcal{A} = 4\pi\mu^{-2\epsilon} \sum_j \left(\frac{\alpha}{\pi} \right)^j \mathcal{A}_j$$



Feynman
diagrams

σ_{LO}

$$\int \sum_{ij} \left[\begin{array}{c} \text{tree} \\ \text{tree} \end{array} \right] d\text{PS}_2$$

σ_{NLO}

$$\int \sum_{ij} \left[\begin{array}{c} \text{1L} \\ \text{tree} \end{array} \right] d\text{PS}_2$$

σ_{NNLO}

$$\int \sum_{ij} \left[\begin{array}{c} \text{2L} \\ \text{1L} \end{array} \right] d\text{PS}_2$$


$$\int \sum_{ij} \left[\begin{array}{c} \text{tree} \\ \text{tree} \end{array} \right] d\text{PS}_3$$

$$\int \sum_{ij} \left[\begin{array}{c} \text{1L} \\ \text{1L} \end{array} \right] d\text{PS}_3$$

$$\int \sum_{ij} \left[\begin{array}{c} \text{tree} \\ \text{tree} \end{array} \right] d\text{PS}_4$$

Anatomy of $e^- \mu^+ \rightarrow e^- \mu^+$ two-loop NNLO QED contributions

$$\begin{aligned}\mathcal{M}^{(2)} &= \overline{\sum} 2\text{Re}[\mathcal{A}^{(0)*} \mathcal{A}^{(n)}] \\ &= (S_\epsilon)^2 \int d^d k_1 d^d k_2 \sum_G \frac{\mathcal{N}_G(\mathbf{k}, \mathbf{l})}{\prod_{\sigma \in G} D_\sigma(\mathbf{k}, \mathbf{l})}\end{aligned}$$



Two-loop Feynman integrals

- 4-point kinematics
- 4 mass-scales variables
 - 2 Mandelstam s, t
 - 2 masses m_e, m_μ

Observation: $\frac{m_e}{m_\mu} \simeq 10^{-5}$

$$\begin{aligned}m_e &= 0 \\ m_\mu &= M\end{aligned}$$

One less mass-scale


Opens the possibility of an
analytical calculation

*Simplification of the Dirac
trace algebra*

Feynman integrals are (in general) UV and IR divergent

Using **Dimensional Regularization**: space-time is treated as a free parameter $d = 4 - 2\epsilon$

Crossing: $e^- \mu^+ \rightarrow e^- \mu^+$ vs. $e^- e^+ \rightarrow \mu^- \mu^+$



$$e^-(p_1) \mu^+(p_2) \rightarrow e^-(p_3) \mu^+(p_4)$$

$$\begin{aligned} p_1^2 &= p_3^2 = 0 \\ p_2^2 &= p_4^2 = M^2 \end{aligned}$$

$$s = (p_1 + p_2)^2$$

$$t = (p_1 - p_3)^2$$


$$u = (p_2 - p_3)^2$$

$$s + t + u = M^2$$

$$\mathcal{M}^{(0)} = \frac{4(s - M^2)^2 + 4st + (d - 2)t^2}{t^2}$$

Crossing

$$\begin{aligned} s &\rightarrow t \\ t &\rightarrow s \\ u &\rightarrow u \end{aligned}$$



$$e^-(p_1) e^+(p_2) \rightarrow \mu^-(p_3) \mu^+(p_4)$$

$$\begin{aligned} p_1^2 &= p_2^2 = 0 \\ p_3^2 &= p_4^2 = M^2 \end{aligned}$$

$$s = (p_1 + p_2)^2$$

$$t = (p_1 - p_3)^2$$

$$u = (p_2 - p_3)^2$$

$$s + t + u = M^2$$

$$\mathcal{M}^{(0)} = \frac{4(t - M^2)^2 + 4st + (d - 2)s^2}{s^2}$$


From now on, we consider the cross-related **di-muon production**



Integrand decomposition of Feynman Integrals

$$\mathcal{I}(l) = \int_k \frac{\mathcal{N}(k, l)}{D_1(k, l) \cdots D_n(k, l)}$$

@1L



Polynomial division
modulo Gröbner basis

$$\begin{aligned} &= \frac{\mathcal{N}(k, l)}{D_1 \cdots D_j \cdots D_n} \\ &= \sum_{j=1}^n \frac{\Delta_{\hat{j}}(k, l)}{D_1 \cdots D_j \cdots D_n} + \frac{\Delta_{\hat{j}}(k, l)}{D_1 \cdots D_j \cdots D_n} \\ &= \sum_{j=1}^n \sum_{i_1 \cdots i_j} \frac{\Delta_{i_1 \cdots i_j}(k, l)}{D_{i_1} \cdots D_{i_j}} \end{aligned}$$

Iterating...

Contain Irreducible
Scalar Products

[Ossola, Papadopoulos, PiSau (2006)]
[Ellis, Giele, Kunszt, Melnikov (2007)]
[Mastrolia, Ossola, Papadopoulos, PiSau (2008)]

- Generalizable at n-loop
- Works with helicity amplitudes
- Possible automation

[Mastrolia, Ossola (2011)]
[Zhang (2012-2016)]
[Badger, Frellesvig, Zhang (2012-2013)]
[Mastrolia, Mirabella, Ossola, Peraro (2012)]

Adaptive Integrand Decomposition (AID)

$$\mathcal{I}(l) = \int_k \frac{\mathcal{N}(k, l)}{D_1(k, l) \cdots D_n(k, l)}$$

[Collins (1984)]
 [van Neerven and Vermaseren (1984)]
 [Kreimer (1992)]

Idea

$$d = d_{||} + d_{\perp}$$

$$k = k_{||} + k_{\perp}$$



$$k_j^\mu = k_{||j}^\mu + \lambda_j^\mu$$

$$k_i \cdot k_j = k_{||i} \cdot k_{||j} + \lambda_{ij}$$

$$\begin{aligned} \mathcal{I}(l) &= \int d^{d_{||}} k_{||} d^{d_{\perp}} k_{\perp} \frac{\mathcal{N}(k, l)}{D_1(k, l) \cdots D_n(k, l)} \\ &= \int d^{d_{||}} k_{||} \int \prod_{ij} G(\lambda_{ij}) d\lambda_{ij} k_{\perp} d\Theta_{\perp} \frac{\mathcal{N}(k_{||}, \lambda_{ij}, \Theta_{\perp})}{D_1(k_{||}, \lambda_{ij}) \cdots D_n(k_{||}, \lambda_{ij})} \end{aligned}$$

**Transverse direction
can be integrated out**

$$\int d\Theta_{\perp} = \int_{-1}^1 \prod_{i=1}^{4-d_{||}} \prod_{j=1}^L d \cos \theta_{i+j-1, j} (\sin \theta_{i+j-1, j})^{d_{\perp}-i-j-1}$$



$$\int_{-1}^1 dx (1-x^2)^{\alpha-\frac{1}{2}} C_n^{(\alpha)}(x) C_m^{(\alpha)}(x) = \delta_{nm} \frac{2^{1-2\alpha} \pi \Gamma(n+2\alpha)}{n! (n+\alpha) \Gamma^2(\alpha)}$$

[Mastrolia, Peraro, Primo (2016)]
 [Mastrolia, Peraro, Primo, Torres Bobadilla (2016)]

**Gegenbauer
polynomials**

Divide, Integrate, Divide again

$$\int dk \frac{\mathcal{N}(k, l)}{D_1(k, l) \cdots D_n(k, l)}$$

Divide

$$\sum_{j=1}^n \sum_{i_1 \cdots i_j} \int dk_{||} d\lambda d\Theta_{\perp} \frac{\Delta_{i_1 \cdots i_j}(k_{||}, \lambda, \Theta_{\perp})}{D_{i_1}(k_{||}, \lambda) \cdots D_{i_j}(k_{||}, \lambda)}$$

Integrate


$$\sum_{j=1}^n \sum_{i_1 \cdots i_j} \int dk_{||} d\lambda \frac{\Delta_{i_1 \cdots i_j}^{\text{int}}(k_{||}, \lambda)}{D_{i_1}(k_{||}, \lambda) \cdots D_{i_j}(k_{||}, \lambda)}$$

Divide

- Numerator reduced in terms of ISPs
- No need for integral identities at 1L
- At 2L, amplitude ready for IBPs

$$\sum_{j=1}^n \sum_{i_1 \cdots i_j} \int dk_{||} d\lambda \frac{\Delta'_{i_1 \cdots i_j}(k_{||})}{D_{i_1}(k_{||}, \lambda) \cdots D_{i_j}(k_{||}, \lambda)}$$

The AIDA framework



Double Virtuals for:

$e^+e^- \rightarrow \mu^+\mu^-$ @NNLO QED

[Bonciani, Broggio, Di Vita, Ferroglio, Mandal, Mastrolia, Mattiazzi, Primo, Schubert, Torres Bobadilla, Tramontano, JR (2021)]




$q\bar{q} \rightarrow t\bar{t}$ @NNLO QCD

[Mandal, Mastrolia, Torres Bobadilla, JR (2022)]

Di-muon production in QED: Feynman Diagrams


$$\mathcal{M}^{(1)} = A^{(1)} + n_l B_l^{(1)} + n_h C_h^{(1)}$$

$$\begin{aligned}\mathcal{M}^{(2)} = & A^{(2)} + n_l B_l^{(2)} + n_h C_h^{(2)} + n_l^2 D_l^{(2)} \\ & + n_l n_h D_{lh}^{(2)} + n_h^2 F_h^{(2)}\end{aligned}$$



[Bonciani, Broggio, Di Vita, Ferroglio, Mandal, Mastrolia, Mattiazzi, Primo, Schubert, Torres Bobadilla, Tramontano, JR (2021)]

Di-muon production in QED: Feynman Diagrams



$$\begin{aligned}
 m_e &= 0 \\
 m_\mu &= M \\
 s &= (p_1 + p_2)^2 \\
 t &= (p_1 - p_3)^2 \\
 u &= (p_2 - p_3)^2 \\
 s + t + u &= M^2
 \end{aligned}$$

$$\mathcal{M}^{(2)} = (S_\epsilon)^2 \int d^d k_1 d^d k_2 \sum_G \frac{\mathcal{N}_G(\mathbf{k}, \mathbf{l})}{\prod_{\sigma \in G} D_\sigma(\mathbf{k}, \mathbf{l})}$$

$O(10^4)$ terms
max rank = 4

AID

$$\mathcal{M}^{(2)} = \sum_{j=1}^n \sum_{i_1 \dots i_j} (S_\epsilon)^2 \int d^d k_1 d^d k_2 \frac{\Delta'_{i_1 \dots i_j}(\mathbf{k}, \mathbf{l})}{D_{i_1} \dots D_{i_j}}$$

$O(10^4)$ terms
max rank = 2

IBPs

$$\mathcal{M}^{(2)} = \sum_{j=1}^{N_{\text{MI}}} c_j(s, t, M; d) I_j^{(2)}(s, t, M; d)$$

Master Integrals

Master Integrals for $\mu e \rightarrow \mu e$
are known in literature.

$$I_{\mu e \rightarrow \mu e}^{(2)} = I_{ee \rightarrow \mu\mu}^{(2)}|_{s \leftrightarrow t}$$

Representation through Generalized PolyLogarithms

$$I_j^{(2)}(s, t, M; d) = \sum_i C_i(\{w\}_{ji}, d) G(\{w\}_{ji}, 1)$$

where

$$G(w_n, \dots, w_1; \tau) = \int_0^\tau \frac{dt}{t - w_n} G(w_{n-1}, \dots, w_1; t)$$

- O(4000) GPLs
- GPLs up to weight 4
- 18 letters


Letters
 $w_j = w_j(s, t, M)$

[Kotikov (1990)] [Gehrmann, Remiddi (1999)] [Henn (2013)]

[Argeri, Di Vita, Mastrolia, Mirabella, Schlenk, Schubert, Tancredi (2014)]

[Mastrolia, Passera, Primo, Schubert (2017)]

[Di Vita, Laporta, Mastrolia, Primo, Schubert (2018)]



Checks

$$\mathcal{M}^{(1)} = A^{(1)} + n_l B_l^{(1)} + n_h C_h^{(1)}$$

$$\begin{aligned}\mathcal{M}^{(2)} = & A^{(2)} + n_l B_l^{(2)} + n_h C_h^{(2)} + n_l^2 D_l^{(2)} \\ & + n_l n_h D_{lh}^{(2)} + n_h^2 F_h^{(2)}\end{aligned}$$

Evaluating the interferences with
HandyG and **GiNaC**

$$\frac{s}{M^2} = 5, \quad \frac{t}{M^2} = -\frac{5}{4}, \quad \mu = M$$


	ϵ^{-4}	ϵ^{-3}	ϵ^{-2}	ϵ^{-1}	ϵ^0	ϵ
$\mathcal{M}^{(0)}$	-	-	-	-	$\frac{181}{100}$	-2
$A^{(1)}$	-	-	$-\frac{181}{100}$	1.99877525	22.0079572	-11.7311017
$B_l^{(1)}$	-	-	-	-	-0.069056030	4.94328573
$C_h^{(1)}$	-	-	-	-	-2.24934027	2.54943566
$A^{(2)}$	$\frac{181}{400}$	-0.499387626	-35.4922919	19.4997261	49.0559119	-
$B_l^{(2)}$	-	$-\frac{181}{400}$	0.785712779	-16.1576674	-3.75247701	-
$C_h^{(2)}$	-	-	1.12467013	-9.50785825	-25.8771503	-
$D_l^{(2)}$	-	-	-	-	-3.96845688	-
$E_{hl}^{(2)}$	-	-	-	-	-4.88512563	-
$F_h^{(2)}$	-	-	-	-	-0.158490810	-

[Bonciani, Broggio, Di Vita, Ferroglio, Mandal, Mastrolia, Mattiazzi, Primo, Schubert, Torres Bobadilla, Tramontano, JR (2021)]

How do we check these terms?

Checks :: Literature

$q\bar{q} \rightarrow t\bar{t}$ in QCD




$e^-e^+ \rightarrow \mu^-\mu^+$ in QED

- Top-pair production admit a color decomposition
- 1-loop and 2-loop corrections already known in literature
- Abelian part get contributions from *QED-like diagrams* only

[Czakon(2008)]


[Bonciani, Ferroglio, Gehrmann, Maitre, Studerus (2008)]

[Bärnreuther, Czakon, Fiedler (2014)]



Abelian, Color-stripped

Remainder of
color factors



Full agreement with the abelian part of top-pair production

Checks :: IR structure

Two-loop IR poles from one-loop and tree (renormalised) contributions

$$\mathcal{M}^{(1)} \Big|_{\text{poles}} = \frac{1}{2} Z_1^{\text{IR}} \mathcal{M}^{(0)} \Big|_{\text{poles}}$$

$$\mathcal{M}^{(2)} \Big|_{\text{poles}} = \frac{1}{8} \left[\left(Z_2^{\text{IR}} - (Z_1^{\text{IR}})^2 \right) \mathcal{M}^{(0)} + 2 Z_1^{\text{IR}} \mathcal{M}^{(1)} \right] \Big|_{\text{poles}}$$

[Becher, Neubert (2009)]

[Hill (2017)]

IR Renormalisation Factor

$$\ln Z_{\text{IR}} = \frac{\alpha}{4\pi} \left(\frac{\Gamma'_0}{4\epsilon^2} + \frac{\Gamma_0}{2\epsilon} \right) + \left(\frac{\alpha}{4\pi} \right)^2 \left(-\frac{3\beta_0\Gamma'_0}{16\epsilon^3} + \frac{\Gamma'_1 - 4\beta_0\Gamma_0}{16\epsilon^2} + \frac{\Gamma_1}{4\epsilon} \right) + \mathcal{O}(\alpha^3)$$

Beta function

$$\text{Anomalous dimension} \quad \Gamma = \gamma_{\text{cusp}}(\alpha) \ln \left(-\frac{s}{\mu^2} \right) + 2\gamma_{\text{cusp}}(\alpha) \ln \left(\frac{t - M^2}{u - M^2} \right) + \gamma_{\text{cusp,M}}(\alpha, s) + 2\gamma_h(\alpha) + 2\gamma_\psi(\alpha)$$

$$\gamma_i = \sum_{j=0}^n \left(\frac{\alpha}{\pi} \right)^{j+1} \gamma_i^{(j)} + \mathcal{O}(\alpha^{n+1})$$




Full agreement with the direct calculation of the two-loop contribution


Extension to two-loop top-pair production @NNLO QCD

Full two-loop $q\bar{q} \rightarrow t\bar{t}$ @NNLO QCD
calculation available only **numerical**


More particles and interactions




More diagrams (~220)



New topologies occur



BUT



$$\propto T^a_{ij} T^a_{lk} f^{abc} (T^d T^c)_{kl} (T^b T^d T^a)_{ji} = 0 !$$

No additional Master Integrals are required

Extension to two-loop top-pair production @NNLO QCD

$$\mathcal{M}^{(1)} = 2(N_c^2 - 1) \left(N_c A^{(1)} + B^{(1)} + n_l C_l^{(1)} + n_h C_h^{(1)} \right)$$

$$\begin{aligned} \mathcal{M}^{(2)} = 2(N_c^2 - 1) & \left(N_c^2 A^{(2)} + B^{(2)} + \frac{C^{(2)}}{N_c^2} + n_l N_c D_l^{(2)} + n_h N_c D_h^{(2)} \right. \\ & \left. + n_l \frac{E_l^{(2)}}{N_c} + n_h \frac{E_h^{(2)}}{N_c} + n_l^2 F_l^{(2)} + n_l n_h F_{lh}^{(2)} + n_h^2 F_h^{(2)} \right) \end{aligned}$$

Evaluating the interferences with
HandyG and **GiNaC**

$$\frac{s}{M^2} = 5, \quad \frac{t}{M^2} = -\frac{5}{4}, \quad \mu = M$$

**First fully-analytical
calculation**

**Full agreement with
the literature**


- [Czakon(2008)]
- [Bonciani, Ferroglio, Gehrmann, Maitre, Studerus (2008)]
- [Bärnreuther, Czakon, Fiedler (2014)]
- [Fael, Passera (2019)]
- [Fael (2018)]

	ϵ^{-4}	ϵ^{-3}	ϵ^{-2}	ϵ^{-1}	ϵ^0	ϵ^1
$A^{(0)}$	-	-	-	-	$\frac{181}{100}$	-2
$A^{(1)}$	-	-	$-\frac{181}{400}$	0.1026418456757775	1.356145770566065	2.230403451742140
$B^{(1)}$	-	-	$\frac{181}{400}$	-0.3180868339485723	-5.763132746701004	2.913169881363488
$C_l^{(1)}$	-	-	0	0	-0.01726400752682416	1.235821434465827
$C_h^{(1)}$	-	-	0	0	-0.5623350683773134	0.6373589172648111
$A^{(2)}$	$\frac{181}{800}$	1.391733154324222	-2.298174307221209	-4.145752448999165	17.37136598564062	-
$B^{(2)}$	$-\frac{181}{400}$	-1.323646320375650	8.507455541210568	6.035611156200398	-35.12861106350758	-
$C^{(2)}$	$\frac{181}{800}$	-0.06808683394857230	-18.00716652035224	6.302454931016090	3.524044912826756	-
$D_l^{(2)}$	0		$-\frac{181}{800}$	0.2605057338631945	-0.7250180282219092	-1.935417246635768
$D_h^{(2)}$	0		0	0.5623350683773134	0.1045606449242690	-1.704747997587188
$E_l^{(2)}$	0		$\frac{181}{800}$	-0.3323207299541260	7.904121951420471	2.848697836597635
$E_h^{(2)}$	0		0	-0.5623350683773134	4.528240788258799	12.73232424278180
$F_l^{(2)}$	0		0	0	0	-1.984228442234312
$F_{lh}^{(2)}$	0		0	0	0	-2.442562819239786
$F_h^{(2)}$	0		0	0	0	-0.07924540546146283

[Mandal, Mastrolia, Torres Bobadilla, JR (2022)]

Results: One-loop di-muon production @NLO QED


$$\mathcal{M}_0^{(1)} = A^{(1)} + n_l B_l^{(1)} + n_h C_h^{(1)} \Big|_{\text{finite}}$$



$$\eta = \frac{s}{4M^2} - 1, \quad \phi = -\frac{(t - M^2)}{s}$$

Production region


$$\eta > 0, \quad \frac{1}{2} \left(1 - \frac{\eta}{1 + \eta} \right) < \phi < \frac{1}{2} \left(1 + \frac{\eta}{1 + \eta} \right)$$



[Bonciani, Broggio, Di Vita, Ferroglio, Mandal, Mastrolia, Mattiazzi, Primo, Schubert, Torres Bobadilla, Tramontano, JR (2021)]

Results: Two-loop di-muon production @NNLO QED

$$\mathcal{M}_0^{(2)} = A^{(2)} + n_l B_l^{(2)} + n_h C_h^{(2)} + n_l^2 D_l^{(2)} + n_l n_h E_{lh}^{(2)} + n_h^2 F_h^{(2)} \Big|_{\text{finite}}$$




[Bonciani, Broggio, Di Vita, Ferroglio, Mandal, Mastrolia, Mattiazzi, Primo, Schubert, Torres Bobadilla, Tramontano, JR (2021)]

$$\eta = \frac{s}{4M^2} - 1, \quad \phi = -\frac{(t - M^2)}{s}$$


Production region

$$\eta > 0, \quad \frac{1}{2} \left(1 - \frac{\eta}{1 + \eta} \right) < \phi < \frac{1}{2} \left(1 + \frac{\eta}{1 + \eta} \right)$$



Results: One-loop top-pair production @NLO QCD

$$\mathcal{M}_0^{(1)} = 2(N_c^2 - 1) \left(N_c A^{(1)} + \frac{B^{(1)}}{N_c} + n_l C_l^{(1)} + n_h C_h^{(1)} \right) \Big|_{\text{finite}}$$




[Mandal, Mastrolia, Torres Bobadilla, JR (2022)]

$$\eta = \frac{s}{4M^2} - 1, \quad \phi = -\frac{(t - M^2)}{s}$$


Production region

$$\eta > 0, \quad \frac{1}{2} \left(1 - \frac{\eta}{1 + \eta} \right) < \phi < \frac{1}{2} \left(1 + \frac{\eta}{1 + \eta} \right)$$



Results: Two-loop top-pair production @NNLO QCD

$$\mathcal{M}_0^{(2)} = 2(N_c^2 - 1) \left(N_c^2 A^{(2)} + B^{(2)} + \frac{C^{(2)}}{N_c^2} + n_l N_c D_l^{(2)} + n_h N_c D_h^{(2)} + n_l \frac{E_l^{(2)}}{N_c} + n_h \frac{E_h^{(2)}}{N_c} + n_l^2 F_l^{(2)} + n_l n_h F_{lh}^{(2)} + n_h^2 F_h^{(2)} \right) \Big|_{\text{finite}}$$




[Mandal, Mastrolia, Torres Bobadilla, JR (2022)]

$$\eta = \frac{s}{4M^2} - 1, \quad \phi = -\frac{(t - M^2)}{s}$$

Production region

$$\eta > 0, \quad \frac{1}{2} \left(1 - \frac{\eta}{1 + \eta} \right) < \phi < \frac{1}{2} \left(1 + \frac{\eta}{1 + \eta} \right)$$



Conclusions & Outlook


Electron-muon elastic scattering @NNLO QED is a crucial input for the **MuonE experiment**

- ✓ **Crossing:** the two-loop contributions to **di-muon production** @NNLO QED via electron-positron annihilation
 - ✓ First QED **analytical two-loop** calculation for di-muon production process
 - ✓ Complete **automation** through the **AIDA framework**
 - ✓ **Cross-checked** against
 - Independent calculations
 - IR structure cross-checked against the SCET prediction
 - ✓ Grid of 10500 phase-space points has been generated
- ✓ Extension to **two-loop** contributions to **top-pair production via quark-antiquark annihilation** @NNLO QCD
- ✓ Recent development: **analytical one-loop squared** contributions to di-muon production @NNLO QED
- ✓ Inclusion of **non-zero** electron mass to electron-muon elastic scattering calculation: **massification**
 - [Mitov, Moch (2006)]
 - [Becher, Melnikov (2007)]
 - [Engel, Gnendiger, Signer, Ulrich (2019)]
 - [Heller (2021)]
 - [Broggio, Engel, Ferroglio, Mandal, Mastrolia, Passera, Rocco, Signer, Torres Bobadilla, Ulrich, Zoller (to be published)]
- **Threshold expansion** for both di-muon and top-pair production @NNLO
- Inclusion of our contribution on MC generators

Thank you

UV Renormalization

$\mathcal{M}_b^{(2)}$ is UV divergent Renormalisation $\rightarrow \mathcal{M}^{(2)}$



$$\mathcal{M} = Z_{2,e} Z_{2,\mu} \mathcal{M}_b(\alpha_b = \alpha_b(\alpha), M_b = M_b(M))$$

where

$$M_b(M) = Z_M M$$

$$\alpha_s S_\epsilon = \alpha(\mu^2) \mu^{2\epsilon} Z_\alpha$$

Renormalisation constants:

$$Z_j = 1 + \left(\frac{\alpha}{\pi}\right) \delta Z_j^{(1)} + \left(\frac{\alpha}{\pi}\right)^2 \delta Z_j^{(2)} + O(\alpha^3)$$

Renormalisation schemes

- **On-shell** renormalisation $Z_{2,e}, Z_{2,\mu}, Z_M$
- **$\overline{\text{MS}}$** renormalisation Z_α

Renormalised interferences:

$$\mathcal{M}^{(0)} = \mathcal{M}_b^{(0)}$$

$$\mathcal{M}^{(1)} = \mathcal{M}_b^{(1)} + (\delta Z_{2,\mu}^{(1)} + Z_\alpha^{(1)}) \mathcal{M}_b^{(0)}$$

$$\mathcal{M}^{(2)} = \mathcal{M}_b^{(2)} + (\delta Z_{2,\mu}^{(1)} + Z_\alpha^{(1)}) \mathcal{M}_b^{(1)}$$

$$+ (\delta Z_{2,\mu}^{(2)} + \delta Z_{2,e}^{(2)} + Z_\alpha^{(2)} + \delta Z_{2,\mu}^{(1)} Z_\alpha^{(1)}) \mathcal{M}_b^{(0)}$$

$$+ \delta Z_M^{(1)} \mathcal{M}_{\text{massCT}}^{(1)}$$

