

The pT distribution of Higgs production at next-to-leading order in α_s

Vittorio Del Duca


ETH Zürich & U. Zürich & INFN

in collaboration with

R. Bonciani, H. Frellesvig, M. Hidding, V. Hirschi,
F. Moriello, G. Salvatori, G. Somogyi, F. Tramontano
J. Henn, L. Maestri, V. Smirnov

HP2 20 September 2022

Higgs p_T distribution at LHC



- ➊ high- p_T tail of the Higgs p_T distribution is sensitive to the structure of the loop-mediated Higgs-gluon coupling
New Physics particles circulating in the loop would modify it
- ➋ QCD NLO corrections to the top- and b -quark loop contributions to the Higgs p_T distribution, in the on-shell and MSbar mass renormalisation schemes

Higgs production at LHC




In proton collisions, the Higgs boson is produced mostly via gluon fusion

The gluons do not couple directly to the Higgs boson

For matter, the coupling is mediated by a heavy quark loop




The largest contribution comes from the top-quark loop

The production mode is (roughly) proportional to the top Yukawa coupling y_t^2



QCD NLO corrections (for any heavy quark mass)

Djouadi Graudenz Spira Zerwas 1991-1995



QCD NLO corrections are about 100% larger than leading order




QCD NNLO corrections are known for the top-quark loop only

Czakon Harlander Klappert Niggetiedt 2021

QCD NLO corrections




real radiation



K. Ellis Hinchliffe Soldate van der Bij 1988



virtual corrections



Djouadi Graudenz Spira Zerwas 1993

Anastasiou Beerli Bucherer Daleo Kunszt 2006


Aglietti Bonciani Degrassi Vicini 2006

} in terms of Harmonic Polylogarithms (HPL)

QCD NLO corrections



$m_H \ll 2m_t$



all amplitudes are reduced by one loop

σ_{EFT}^{LO}	15.05 pb	σ_{EFT}^{NLO}	34.66 pb
$R_{LO} \sigma_{EFT}^{LO}$	16.00 pb	$R_{LO} \sigma_{EFT}^{NLO}$	36.84 pb
$\sigma_{ex;t}^{LO}$	16.00 pb	$\sigma_{ex;t}^{NLO}$	36.60 pb
$\sigma_{ex;t+b}^{LO}$	14.94 pb	$\sigma_{ex;t+b}^{NLO}$	34.96 pb
$\sigma_{ex;t+b+c}^{LO}$	14.83 pb	$\sigma_{ex;t+b+c}^{NLO}$	34.77 pb

$$\frac{\sigma_{t+b}}{\sigma_t} - 1$$

$\text{LO } \mathcal{O}(\alpha_s^2)$	- 6.6 %
$\text{NLO } \mathcal{O}(\alpha_s^2) + \mathcal{O}(\alpha_s^3)$	- 4.5 %
$\text{NLO } \mathcal{O}(\alpha_s^3)$	- 2.8 %



$$R_{LO} = \frac{\sigma_{ex:t}^{LO}}{\sigma_{EFT}^{LO}} = 1.063$$

rescaled HEFT (rHEFT) does a good job (< 1%) in approximating the exact (only top) NLO σ but misses the $t\bar{b}$ interference

Anastasiou Duhr Dulat Furlan Gehrmann Herzog Lazopoulos Mistlberger 2016

QCD NNLO corrections



Top-quark mass corrections are known at NNLO

Czakon Harlander Klappert Niggetiedt 2021

channel	$\sigma_{\text{HEFT}}^{\text{NNLO}} [\text{pb}]$ $\mathcal{O}(\alpha_s^2) + \mathcal{O}(\alpha_s^3) + \mathcal{O}(\alpha_s^4)$	$(\sigma_{\text{exact}}^{\text{NNLO}} - \sigma_{\text{HEFT}}^{\text{NNLO}}) [\text{pb}]$ $\mathcal{O}(\alpha_s^3)$	$(\sigma_{\text{exact}}^{\text{NNLO}} / \sigma_{\text{HEFT}}^{\text{NNLO}} - 1) [\%]$
$\sqrt{s} = 8 \text{ TeV}$			
gg	$7.39 + 8.58 + 3.88$	+0.0353	+0.62
qg	$0.55 + 0.26$	-0.1397	-18
qq	$0.01 + 0.04$	+0.0171	-4
total	$7.39 + 9.15 + 4.18$	-0.0873	-0.10
$\sqrt{s} = 13 \text{ TeV}$			
gg	$16.30 + 19.64 + 8.76$	+0.0345	+0.62
qg	$1.49 + 0.84$	-0.3696	-16
qq	$0.02 + 0.10$	+0.0322	-15
total	$16.30 + 21.15 + 9.79$	-0.3029	-0.26



HEFT not so good for qg and qq channels






for top-quark mass, used $m_t^2/m_H^2 = 23/12$ (on-shell scheme)

The main obstacle when calculating the total cross section with full top-mass dependence
are the two-loop single-emission amplitudes.

Czakon Harlander Klappert Niggetiedt 2021

QCD NNLO corrections



Higgs + 4-parton amplitudes at one loop



OpenLoops

VDD Kilgore Oleari Schmidt Zeppenfeld 2001
Budge Campbell De Laurentis K. Ellis Seth 2020


Higgs + 3-parton amplitudes at two loops



top loop: Jones Kerner Luisoni 2018
Czakon Harlander Klappert Niggetiedt 2021

Bonciani VDD Frellesvig Moriello Hidding
Hirschi Salvatori Somogyi Tramontano 2022

gg \rightarrow Higgs amplitudes at three loops



one scale: one & two top loops
one top loop + light-quark loop


two scales: one top loop + b -quark loop

Czakon Niggetiedt 2020
Harlander Prausa Usovitsch 2019

Higgs p_T distribution at LHC



leading order



K. Ellis Hinchliffe Soldate van der Bij 1988

- high- p_T tail of the Higgs p_T distribution is sensitive to the structure of the loop-mediated Higgs-gluon coupling
New Physics particles circulating in the loop would modify it
- in high- p_T regime, clean signature of decay products ($H \rightarrow b\bar{b}$)



QCD NLO corrections

- for the top-quark, with on-shell scheme

Jones Kerner Luisoni 2018
Chen Huss Jones Kerner Lang Lindert Zhang 2021

- for the top-quark, with on-shell and MSbar schemes
for top- and b -quarks (for any heavy quark mass), with MSbar scheme

Bonciani VDD Frellesvig Moriello Hidding Hirschi Salvatori Somogyi Tramontano 2022



HEFT $m_H \ll 2m_t$ and $p_T \ll m_t$ Baur Glover 1990


QCD corrections are known at **NNLO** in HEFT, and yield a 15% increase wrt **NLO**

Boughezal Caola Melnikov Petriello Schulze 2015
Boughezal Focke Giele Liu Petriello 2015
Chen Cruz-Martinez Gehrmann Glover Jaquier 2016

Higgs p_T distribution at NLO



virtual corrections



top-quark loop

Jones Kerner Luisoni 2018


Czakon Harlander Klappert Niggetiedt 2021

any heavy quark in the loop

Bonciani VDD Frellesvig Henn Moriello V. Smirnov 2016

all above + Hidding Maestri Salvatori 2019


Bonciani VDD Frellesvig Moriello Hidding Hirschi Salvatori Somogyi Tramontano 2022




multi-scale problem with complicated analytic structure
elliptic iterated integrals appear



real corrections




VDD Kilgore Oleari Schmidt Zeppenfeld 2001
Budge Campbell De Laurentis K. Ellis Seth 2020





one-loop amplitudes for Higgs + 3-partons



leading order: up to $\mathcal{O}(\varepsilon^2)$

analytic: up to $\mathcal{O}(\varepsilon^0)$


K. Ellis Hinchliffe Soldate van der Bij 1988

numeric: up to $\mathcal{O}(\varepsilon^2)$

(numeric) derivative for mass renormalisation



one-loop amplitudes for Higgs + 4-partons



NLO real corrections: up to $\mathcal{O}(\varepsilon^0)$

analytic: unitarity-cut methods (taken from MCFM-9.I)

Budge Campbell De Laurentis K. Ellis Seth 2020


numeric: GoSam & MG5_aMC

run time

analytic: few ms/pt

numeric: $\mathcal{O}(100)$ times slower than analytic

two-loop amplitudes for Higgs + 3-partons



NLO virtual corrections

amplitude \rightarrow form factors \rightarrow scalar integrals \rightarrow Master Integrals
IBP

run time: 5 — 60 min/pt


FIRE-KIRA

4 scales, $s, t, m_H, m_t \rightarrow$ 3 external parameters

7 seven-propagator integral families

Bonciani VDD Frellesvig Henn Moriello Smirnov 2016 (A, B, C, D)
Bonciani VDD Frellesvig Henn Hidding Maestri Moriello Salvatori Smirnov 2019 (F)
Frellesvig Hidding Maestri Moriello Salvatori 2019 (G)
Bonciani VDD Frellesvig Moriello Hidding Hirschi Salvatori Somogyi Tramontano 2022 (H)

elliptic



A

B

C

D

MIs

A: 72

B: 5

C: 45

D: 17



F: 73

G: 84

H: 12

= 0

colour conservation



E

F


G

H

elliptic




two masses



Elliptic iterated integrals



2-loop sunrise graph




Sabry 1962: ...; Broadhurst 1989; ...; Bloch Vanhove 2013; ...
Brödel Duhr Dulat Penante Tancredi 2017-2019




2-loop 3-pt functions

electroweak form factor



Aglietti Bonciani Grassi Remiddi 2007


$t\text{-}t\bar{b}$



von Manteuffel Tancredi 2017




2-loop 4-pt function for Higgs + 1 jet



Bonciani VDD Frellesvig Henn Moriello Smirnov 2016

first instance of elliptic iterated integrals
in a genuine 4-pt topology

Family F: 73 MIs (65 in the polylogarithmic sector, 8 in the elliptic sector)
alphabet: 69 independent letters, with 12 independent square roots



Differential Equations



Differential Equation method to solve the MIs

$$\partial_i f(x_n; \varepsilon) = A_i(x_n; \varepsilon) f(x_n; \varepsilon)$$

f : N-vector of MIs, A_i : NxN matrix, $i=1,\dots,n$ external parameters

but in some cases ε -independent form

$$\partial_i f(x_n; \varepsilon) = \varepsilon A_i(x_n) f(x_n; \varepsilon)$$

Henn 2013

solution in terms of iterated integrals



mass values are floating →

DEs solved with 3 (top) or 4 (top and b) external parameters

DEs: Series Expansion Method

- Take two points (a_1, \dots, a_n) and (b_1, \dots, b_n) in the n -dim parameter space, and parametrise the contour $\gamma(t)$ that connects the two points

$$\gamma(t) : t \rightarrow \{x_1(t), \dots, x_n(t)\} \quad \vec{x}(0) = \vec{a}, \quad \vec{x}(1) = \vec{b}$$

and write the differential equation with respect to t .

Then find a solution about a point τ by series expanding the coefficient matrix A and then iteratively integrating it.

The procedure works for both polylogarithmic and elliptic sectors

Moriello 2019

- numerical solution of DEs through [DiffExp](#):

Mathematica implementation of Moriello's series expansion method

Hidding 2021

- checked with AMFlow Liu Ma Wang 2018

two-loop amplitudes for Higgs + 3-partons: Renormalisation

Bonciani VDD Frellesvig Moriello Hidding Hirschi Salvatori Somogyi Tramontano 2022




coupling constant: 5-flavour running in MSbar



renormalisation:

- top Yukawa coupling and top mass in OS scheme (massless b)
- top Yukawa coupling and top mass in MSbar scheme (massless b)
- top Yukawa coupling and top and b masses in MSbar scheme



massive b in Higgs- b loop
massless b in b loop

alternative:

massive b everywhere,

but requires 4-flavour running and including $gg \rightarrow Hbb$

two-loop amplitudes for Higgs + 3-partons: validation checks

IR poles

$$\mathcal{M}_{ij,IR}^{(2)} \propto I_{ij}^{(1)}(\{p\}, \epsilon) \mathcal{M}_{ij}^{(1)}$$



with insertion operators

$$I_{gg}^{(1)}(\{p\}, \epsilon) = -\frac{\alpha_S}{\pi} \frac{(4\pi)^\epsilon}{\Gamma(1-\epsilon)} \left(\frac{N_c}{\epsilon^2} + \frac{\beta_0}{\epsilon} \right) \left[\left(\frac{\mu^2}{-s} \right)^\epsilon + \left(\frac{\mu^2}{-t} \right)^\epsilon + \left(\frac{\mu^2}{-u} \right)^\epsilon \right]$$

$$I_{q\bar{q}}^{(1)}(\{p\}, \epsilon) = -\frac{\alpha_S}{2\pi} \frac{(4\pi)^\epsilon}{\Gamma(1-\epsilon)} \left\{ - \left(\frac{N_c}{\epsilon^2} + \frac{3N_c}{4\epsilon} + \frac{\beta_0}{2\epsilon} \right) \left[\left(\frac{\mu^2}{-t} \right)^\epsilon + \left(\frac{\mu^2}{-u} \right)^\epsilon \right] + \frac{1}{N_c} \left(\frac{1}{\epsilon^2} + \frac{3}{2\epsilon} \right) \left(\frac{\mu^2}{-s} \right)^\epsilon \right\}$$

agreement with HEFT limit

$$\mathcal{M} = \mathcal{M}_{HEFT} + \mathcal{O}\left(\frac{1}{M_t}\right)$$




two-loop amplitudes for Higgs + 3-partons: validation checks



soft and collinear limits

(these are checks on real-virtual parts of NNLO cross section,
however they are feasible on our two-loop amplitudes)



Aglietti Bonciani Degrassi Vicini 2006

one-loop 2-parton splitting functions

Bern Dixon Dunbar Kosower 1994
Bern Kilgore Schmidt VDD 1998-99
Kosower Uwer 1999

one-loop 1-soft-gluon factor

Bern Kilgore Schmidt VDD 1998-99
Catani Grazzini 2000



checked also “two-loop photon correction”

Higgs p_T distribution at NLO: checks with previous results


- inclusive p_T distribution ($p_{T,j} > 30$ GeV)
with OS mass renormalisation

our result

$$\sigma_{NLO} = 14.37 \pm 0.05 \text{ pb}$$

Chen Huss Jones Kerner Lang Lindert Zhang 2021
(Jones Kerner Luisoni 2018-2021)

$$\sigma_{NLO} = 14.15 \pm 0.07 \text{ pb}$$




- high p_T tail of distribution

checked with approximate high- p_T distribution

Lindert Melnikov Kudashkin Wever 2018

based on approximate high- p_T two-loop amplitudes


Melnikov Kudashkin Wever 2018



Higgs p_T distribution at LHC



QCD NLO corrections for the top-quark (on-shell mass renormalisation)



Jones Kerner Luisoni 2018

Chen Huss Jones Kerner Lang Lindert Zhang 2021

$$\frac{d\sigma}{dp_T^2} \propto \frac{1}{p_T^2} \quad \text{in HEFT NLO corrections}$$


$$\frac{d\sigma}{dp_T^2} \propto \frac{1}{(p_T^2)^2} \quad \text{in top NLO corrections}$$

NLO/LO in HEFT and top loop agree to $O(10\%)$



QCD NLO corrections to top- b interference, using top-quark loop in HEFT and b -quark loop in small m_b limit

Lindert Melnikov Tancredi Wever 2017



Higgs p_T distribution at NLO



p_T distribution computed with

CoLorFULNLO

Somogyi 2009

dual subtraction

Prisco Tramontano 2020



evaluated on:

3×10^4 pt for OS top (1.4×10^4 pt on basic grid, 1.6×10^4 pt on biased grid)

9×10^4 pt for MSbar top

1.8×10^5 pt for MSbar top and b



set-up

$\sqrt{s} = 13$ TeV

$p_{T,j_1} > 20$ GeV

$m_H = 125.25$ GeV

anti-kt algorithm with $R = 0.4$

$m_t^{\text{OS}} = 172.5$ GeV

7-pt scale variation about:

$m_t^{\overline{\text{MS}}} (m_t^{\overline{\text{MS}}}) = 163.4$ GeV

$$\mu_R^0 = \mu_F^0 = \frac{H_T}{2} = \frac{1}{2} \left(\sqrt{m_H^2 + p_T^2} + \sum_i |p_{T,i}| \right)$$

$m_b^{\overline{\text{MS}}} (m_b^{\overline{\text{MS}}}) = 4.18$ GeV

$G_F = 1.16639 \cdot 10^{-5}$ GeV $^{-2}$

NNPDF40_nlo_as_01180

inclusive Higgs p_T distribution



QCD NLO corrections

Bonciani VDD Frellesvig Moriello Hidding Hirschi Salvatori Somogyi Tramontano 2022

for the top-quark, with on-shell and MSbar schemes
for top- and b -quarks with MSbar scheme

renormalisation of internal masses	σ_{LO} [pb]	σ_{NLO} [pb]
top+bottom- $(\overline{\text{MS}})$	$12.318^{+4.711}_{-3.117}$	$19.89(8)^{+2.84}_{-3.19}$
top- $(\overline{\text{MS}})$	$12.538^{+4.822}_{-3.183}$	$19.90(8)^{+2.66}_{-2.85}$
top-(OS)	$12.551^{+4.933}_{-3.244}$	$20.22(8)^{+3.06}_{-3.09}$



from LO to NLO large k factor and reduction of scale uncertainty





top- b interference is a negative correction at $\mathcal{O}(\alpha_s^3)$ but positive at $\mathcal{O}(\alpha_s^4)$



effect of top mass renormalisation utterly negligible at LO
but 15 times bigger at NLO

$$\frac{\sigma_{t(\text{OS})}}{\sigma_{t(\overline{\text{MS}})}} - 1 = \begin{cases} 0.1\% \text{ at LO} \\ 1.6\% \text{ at NLO} \end{cases}$$



Higgs p_T distribution at low-intermediate p_T



20-40 GeV bin
 260^{+16}_{-83} fb/GeV
 249^{+21}_{-65} fb/GeV
 238^{+27}_{-98} fb/GeV

- at LO no events below 20 GeV since $p_{T,j} > 20$ GeV
- at LO no appreciable difference between $t(\text{OS})$ and $t(\text{MSbar})$
- at NLO sizeable shape distortion in the lowest bins
- at NLO agreement (not shown) between exact and rHEFT in the low-middle p_T range
 $\text{HEFT} \quad m_H \ll 2m_t \quad \text{and} \quad m_b \ll p_T \ll m_t$
- scale uncertainty bands (not shown) are much larger than differences

Higgs p_T distribution at LHC




scale uncertainty bands = ratio of bands at NLO over central value at LO



k factor almost always larger than 2 for MSbar, and about 2 for OS

Ratios of Higgs p_T distributions



- from LO to NLO, reduction of scale uncertainty and of mass renormalisation scheme dependence
- except in the lowest bins, no appreciable difference between $t+b(\text{MSbar})$ and $t(\text{MSbar})$
The b quark, and thus top- b interference, is negligible, except at low end of p_T range
- p_T distribution for $t(\text{MSbar})$ falls off faster than same for $t(\text{OS})$ as p_T increases because μ_R increases with p_T and so $m_t^{\text{MS}}(\mu_R)$ decreases
- mass renormalisation scheme difference between $t(\text{MSbar})$ and $t(\text{OS})$ is same size as scale uncertainty at high end of p_T range, both at LO and NLO

Conclusions


- we computed the Higgs p_T distribution at NLO in QCD including for the first time top and b quarks and the MSbar mass scheme
- computation has excellent numerical stability
- b quark, and thus top- b interference, is negligible, except at low end of p_T range, where it affects the shape of the distribution
- in the intermediate to high p_T range, use of top quark only is warranted, but sizeable dependence on mass renormalisation scheme
- p_T distribution can be improved:
mixed QCD-EW corrections (we already have $gg \rightarrow Hg$),
resummation,
top-charm interference, ...

Back-up slides

Series Expansion Method: patching the contour

F. Moriello at Amplitudes 2020

- Local series solution converges up to the closest singular point: need multiple series to patch the contour
- Truncated series: to ensure fast convergence, radius set to half distance between the expansion point and closest singularity




- The series depend on boundary constants fixed by using boundary point and continuity at the contact points.


By setting each series to zero outside its radius:

$$\mathbf{I}^{(i)}(t) = \mathbf{I}_0^{(i)}(t) + \mathbf{I}_1^{(i)}(t) + \mathbf{I}_2^{(i)}(t) + \mathbf{I}_3^{(i)}(t), \quad t \in [0, 1]$$

Higgs p_T distribution due to QCD-EW interference



Becchetti Bonciani VDD Hirschi Moriello Schweitzer 2020



gg-initiated QCD-EW p_T spectrum harder than HEFT

QCD-EW Higgs+3-parton master integrals at two loops


4 scales, $s, t, m_H, m_V \rightarrow 3$ external parameters

7 seven-propagator integral families


48 MIs (planar), 61 MIs (non-planar)

alphabet: square roots are present, but an MPL representation is possible

Becchetti Bonciani Casconi VDD Moriello 2018 (planar MIs)
Becchetti Moriello Schweitzer 2021 (non-planar MIs)



planar



non-planar

solved through generalised power series expansion Moriello 2019

Higgs production



QCD corrections have been computed at **N³LO** in HEFT

Anastasiou Duhr Dulat Herzog Mistlberger 2015
Mistlberger 2018
(in terms of MPLs and elliptic integrals)



including quark-mass effects and **QCD-EW** interference the cross section is

$$\sigma = 48.58 \text{ pb}^{+2.22 \text{ pb} (+4.56\%)}_{-3.27 \text{ pb} (-6.72\%)} \text{ (theory) } \pm 1.56 \text{ pb (3.20\%)} \text{ (PDF} + \alpha_s \text{)}$$



The breakdown of the cross section

$$\begin{aligned} 48.58 \text{ pb} &= 16.00 \text{ pb } (+32.9\%) && (\text{LO, rEFT}) \\ &+ 20.84 \text{ pb } (+42.9\%) && (\text{NLO, rEFT}) \\ &- 2.05 \text{ pb } (-4.2\%) && ((t, b, c), \text{exact NLO}) \\ &+ 9.56 \text{ pb } (+19.7\%) && (\text{NNLO, rEFT}) \\ &+ 0.34 \text{ pb } (+0.2\%) && (\text{NNLO, } 1/m_t) \\ &+ 2.40 \text{ pb } (+4.9\%) && (\text{EW, QCD-EW}) \\ &+ 1.49 \text{ pb } (+3.1\%) && (\text{N}^3\text{LO, rEFT}) \end{aligned}$$

Anastasiou Duhr Dulat Furlan Gehrmann Herzog Lazopoulos Mistlberger 2016
Handbook 4 of LHC Higgs Cross Sections 2016

Higgs production

Handbook 4 of LHC Higgs Cross Sections 2016



- 6 sources of uncertainties due to:
 - higher orders
 - truncation of the threshold expansion
 - PDFs
 - NLO corrections to QCD-EW interference
 - quark mass effects (2: top mass and top-b interference) at NNLO

$\delta(\text{scale})$	$\delta(\text{trunc})$	$\delta(\text{PDF-TH})$	$\delta(\text{EW})$	$\delta(t, b, c)$	$\delta(1/m_t)$
+0.10 pb -1.15 pb	$\pm 0.18 \text{ pb}$	$\pm 0.56 \text{ pb}$	$\pm 0.49 \text{ pb}$	$\pm 0.40 \text{ pb}$	$\pm 0.49 \text{ pb}$
+0.21% -2.37%	$\pm 0.37\%$	$\pm 1.16\%$	$\pm 1\%$	$\pm 0.83\%$	$\pm 1\%$

$$\delta(\text{trunc}) = 0.11 \text{ pb} \quad \text{Mistlberger 2018}$$

$$\delta(1/m_t) = -0.26\% \quad \text{Czakon Harlander Klappert Niggetiedt 2021}$$

Polylogarithms



classical polylogarithms

$$\text{Li}_m(z) = \int_0^z dt \frac{\text{Li}_{m-1}(t)}{t} = \sum_{n=1}^{\infty} \frac{z^n}{n^m}$$
$$\text{Li}_1(z) = \sum_{n=1}^{\infty} \frac{z^n}{n} = -\ln(1-z)$$



harmonic polylogarithms (HPLs)

Euler 1768
Spence 1809

$$H(a, \vec{w}; z) = \int_0^z dt f(a; t) H(\vec{w}; t)$$
$$f(-1; t) = \frac{1}{1+t}, \quad f(0; t) = \frac{1}{t}, \quad f(1; t) = \frac{1}{1-t}$$

with $\{a, \vec{w}\} \in \{-1, 0, 1\}$

Remiddi Vermaseren 1999



classical polylogarithms are multiple polylogarithms with specific roots (0 and constant a)

$$G(\vec{0}_n; x) = \frac{1}{n!} \ln^n x \quad G(\vec{a}_n; x) = \frac{1}{n!} \ln^n \left(1 - \frac{x}{a}\right) \quad G(\vec{0}_{n-1}, a; x) = -\text{Li}_n \left(\frac{x}{a}\right)$$



when the root equals +1,-1,0 multiple polylogarithms become HPLs

Multiple polylogarithms

$$G(a, \vec{w}; z) = \int_0^z \frac{dt}{t-a} G(\vec{w}; t), \quad G(a; z) = \ln \left(1 - \frac{z}{a}\right)$$

$a, \vec{w} \in \mathbb{C}$

Goncharov 1998-2001



multiple polylogarithms (MPL) form a shuffle algebra

$$G_{\omega_1}(z) G_{\omega_2}(z) = \sum_{\omega} G_{\omega}(z) \quad \text{with } \omega \text{ the shuffle of } \omega_1 \text{ and } \omega_2$$

For a constant
Poincaré Kummer
Lappo-Danilevsky 1935

example

$$\begin{aligned} G(a; z) G(b; z) &= \int_0^z \frac{dt_1}{t_1 - a} \int_0^z \frac{dt_2}{t_2 - b} \\ &= \int_0^z \frac{dt_1}{t_1 - a} \int_0^{t_1} \frac{dt_2}{t_2 - b} + \int_0^z \frac{dt_2}{t_2 - a} \int_0^{t_2} \frac{dt_1}{t_1 - b} \\ &= G(a, b; z) + G(b, a; z) \end{aligned}$$



$$\lim_{z \rightarrow 0} G(a_1, \dots, a_n; z) = 0 \quad \text{unless} \quad \vec{a} = \vec{0}$$



$$\frac{\partial}{\partial z} G(a_1, \dots, a_k; z) = \frac{1}{z - a_1} G(a_2, \dots, a_k; z)$$




MPLs can be represented as nested harmonic sums

$$\sum_{n_1=1}^{\infty} \frac{u_1^{n_1}}{n_1^{m_1}} \sum_{n_2=1}^{n_1-1} \dots \sum_{n_k=1}^{n_{k-1}-1} \frac{u_k^{n_k}}{n_k^{m_k}} = (-1)^k G \left(\underbrace{0, \dots, 0}_{m_1-1}, \frac{1}{u_1}, \dots, \underbrace{0, \dots, 0}_{m_k-1}, \frac{1}{u_1 \dots u_k}; 1 \right)$$

QCD NNLO corrections

virtual corrections



Harlander Prausa Usovitsch 2019


(one top & one light quark,
in terms of HPLs)

Czakon Niggetiedt 2020

(one & two top)

Anastasiou Deutschmann Schweitzer 2020

real-virtual corrections




Jones Kerner Luisoni 2018

Czakon Harlander Klappert Niggetiedt 2021

(top)

double-real radiation



VDD Kilgore Oleari Schmidt Zeppenfeld 2001

Budge Campbell De Laurentis K. Ellis Seth 2020



iterated integrals on $\mathcal{M}_{0,p}$ are multiple polylogarithms

Brown 2006

$\mathcal{M}_{0,p}$ = space of configurations of p points on the Riemann sphere

$$G(a, \vec{w}; z) = \int_0^z \frac{dt}{t-a} G(\vec{w}; t), \quad G(a; z) = \ln\left(1 - \frac{z}{a}\right) \quad a, \vec{w} \in \mathbb{C}$$




iterated integrals on a torus ...

Brown Levin 2011

$$\tilde{\Gamma} \left(\begin{smallmatrix} n_1 \dots n_k \\ z_1 \dots z_k \end{smallmatrix} ; z, \tau \right) = \int_0^z dt g^{(n_1)}(t - z_1, \tau) \tilde{\Gamma} \left(\begin{smallmatrix} n_2 \dots n_k \\ z_2 \dots z_k \end{smallmatrix} ; t, \tau \right)$$

kernels $g^{(n)}$ have at most simple poles at $z = m + n\tau$



... are elliptic multiple polylogarithms (eMPL)

$$E_3 \left(\begin{smallmatrix} n_1 \dots n_k \\ z_1 \dots z_k \end{smallmatrix} ; z, \vec{a} \right) = \int_0^z dt \varphi_{n_1}(z_1, t, \vec{a}) E_3 \left(\begin{smallmatrix} n_2 \dots n_k \\ z_2 \dots z_k \end{smallmatrix} ; t, \vec{a} \right) \quad n_i \in \mathbb{Z}, \quad z_i \in \mathbb{C} \quad a_i \in \mathbb{R}$$

with $\vec{a} = (a_1, a_2, a_3)$ are the zeroes of the elliptic curve

$$y^2 = (x - a_1)(x - a_2)(x - a_3)$$

and $E_3(; z, \vec{a}) = 1$




2-loop sunrise can be written in terms of eMPLs

Brödel Duhr Dulat Penante Tancredi 2017

QCD-EW interference

- 💡 The Higgs boson may (indirectly) couple to gluons also via the gauge coupling i.e. through a double (electroweak boson + quark) loop



Aglietti Bonciani Degrassi Vicini 2004
(light fermion loop)
Degrassi Maltoni 2004
Actis Passarino Sturm Uccirati 2008
(heavy fermion loop)

(in terms of MPLs)

(numerically
... elliptic integrals appear)

$$O(\alpha_s^2 \alpha^2)$$

- 💡 the top loop yields a 2% correction to the 5 light fermion loops

- 💡 gg-initiated QCD NLO corrections (light fermion loop) computed in various approximations:

— $m_{W,Z} \rightarrow \infty$ limit

Anastasiou Boughezal Petriello 2009

— soft approximation

Bonetti Melnikov Tancredi 2018

— $m_{W,Z} \rightarrow 0$ limit

Anastasiou VDD Furlan Mistlberger Moriello Schweitzer Specchia 2018


and found to be about 5% wrt NLO (HEFT) cross section

QCD-EW interference




gg-initiated QCD NLO corrections (light fermion loop): $\mathcal{O}(\alpha_s^3 \alpha^2)$

Becchetti Bonciani VDD Hirschi Moriello Schweitzer 2020



Bonetti Melnikov Tancredi 2016



Becchetti Bonciani Casconi VDD Moriello 2018

Bonetti Panzer V. Smirnov Tancredi 2020

Becchetti Moriello Schweitzer 2021

IR local subtraction schemes

MadGraph MC@NLO

Frixione Kunszt Signer 1995
Frederix Frixione Maltoni Stelzer 2009

COLORFUL

VDD Somogyi Trocsanyi 2006
Somogyi 2009
VDD Deutschmann Lionetti 2019

$$\text{LO} \quad \sigma_{gg \rightarrow H+X}^{(\alpha_s^2 \alpha^2)} = 0.68739^{+23.4\%+2.0\%}_{-17.3\%-2.0\%} \text{ pb}$$

$$\text{NLO} \quad \sigma_{gg \rightarrow H+X}^{(\alpha_s^2 \alpha^2 + \alpha_s^3 \alpha^2)} = 1.467(2)^{+18.7\%+2.0\%}_{-14.6\%-2.0\%} \text{ pb}$$

i.e. NLO 110% wrt LO

gg-initiated NLO corrections in HEFT

$$\sigma_{gg \rightarrow H+X}^{(\text{HEFT}, \alpha_s^2 \alpha + \alpha_s^3 \alpha)} = 30.484^{+19.8\%+1.9\%}_{-15.3\%-1.9\%} \text{ pb}$$

thus our NLO result 4.8% wrt gg-initiated NLO HEFT