New Horizons in Primordial Black Hole physics (NEHOP)



Contribution ID: 43

Type: Talk

Small Primordial Black Holes as Window on Quantum Gravity

Tuesday, 20 June 2023 17:20 (20 minutes)

Hawking's calculation of particle production by a black hole is based on the semi-classical limit of a fixed metric. This approximation may break down after a finite time as a black hole evolves due to back-reaction. Therefore, I shall argue that two far-reaching questions remain to be answered:

(1) How long is the semi-classical description valid?

(2) What happens after a potential breakdown?

Primordial black holes (PBHs) offer unique opportunities to address these issues by observations. Since one possible option is a slowdown of evaporation, it is conceivable that PBHs below 10^{15} g can have survived until today. Such long-lived small PBHs could not only act as dark matter, but detecting even a single one of them would also greatly advance our understanding of quantum gravity. Finally, I will attempt to answer the above questions using explicitly solvable analogue models. An non-perturbative computation of real-time evolution reveals indications that (1) the semi-classical description can break down long before half of the mass is lost and that indeed (2) evaporation slows down drastically at this point.

Based on:

M. Michel, S. Zell, *The Timescales of Quantum Breaking*, arXiv:2203.XXXXX, to appear.
G. Dvali, L. Eisemann, M. Michel, S. Zell, *Black hole metamorphosis and stabilization by memory burden*, Phys. Rev. D 102 (2020) 103523, arXiv:2006.00011.

Primary authors: MICHEL, Marco (Ben-Gurion University of the Negev); ZELL, Sebastian (UCLouvain)

Presenter: ZELL, Sebastian (UCLouvain)

Session Classification: Session 8