



Contribution ID: 26

Type: Long talk (20 mins)

Non-Gaussianity from preheating of non-minimally coupled inflaton

Friday, 16 December 2022 10:10 (30 minutes)

Full non-linear simulations of massless preheating model have revealed a large non-Gaussianity can be generated. We study a more observationally viable model consisting of inflaton non-minimally coupled to gravity that decays into a massless scalar spectator during preheating.

Including the scale-dependence of Hubble rate places tight constraints on the ‘cosmic variance’, the values which the mean spectator field value can take. A negligible cosmic variance forces mean close to zero, and symmetry of many inflation potentials around zero dictates leading order non-perturbative treatment shall fail. Thus, we calculate sub-leading order term in the non-perturbative delta N formalism. As this calculation also needs to be performed keeping in mind the scale dependence, we show a scale-invariant way of solving momentum integrals is insufficient to yield the correct non-Gaussianity from preheating.

Type of presentation

20 minute talk

Would you be interested in receiving feedback on your presentation?

Yes

Are you happy for your talk to be recorded?

Yes

Other categories:

Please select the most relevant category

Cosmology

Primary author: GHODERAO, Pulkit (Imperial College London)

Presenter: GHODERAO, Pulkit (Imperial College London)

Session Classification: Full Length Talks