



Contribution ID: 39

Type: Long talk (20 mins)

Flavonstrahlung in the $B_3 - L_2 Z'$ Model at Current and Future Colliders

Friday, 16 December 2022 14:30 (30 minutes)

The $B_3 - L_2 Z'$ model may explain some gross features of the fermion mass spectrum as well as $b \rightarrow s\ell\ell$ anomalies. A TeV-scale physical scalar field associated with gauged $U(1)_{B_3-L_2}$ spontaneous symmetry breaking, the flavon field ϑ , affects Higgs phenomenology via mixing. In this talk, I will discuss the collider phenomenology of the flavon field. Higgs data are used to place bounds upon parameter space. I then examine “flavonstrahlung” ($Z'^* \rightarrow Z'\vartheta$ production) at colliders as a means to directly produce and discover flavon particles, providing direct empirical evidence tying it to $U(1)_{B_3-L_2}$ symmetry breaking. A 100 TeV FCC-hh or a 10 TeV muon collider would have high sensitivity to flavonstrahlung, whereas the HL-LHC can observe it only if the flavon charge is larger than unity.

Type of presentation

20 minute talk

Would you be interested in receiving feedback on your presentation?

Yes

Are you happy for your talk to be recorded?

Yes

Other categories:

Please select the most relevant category

Beyond the Standard Model

Primary author: LOISA, Eetu (University of Cambridge)

Presenter: LOISA, Eetu (University of Cambridge)

Session Classification: Full Length Talks