

Contribution ID: 147

Type: Talk

QCD Anderson transition with overlap valence quarks on a twisted-mass sea

Monday, 29 July 2024 14:55 (20 minutes)

In this work we probe the QCD Anderson transition by studying spectral distributions of the massless overlap operator on gauge configurations created by the *twisted mass at finite temperature collaboration* (tmfT) with 2+1+1 flavors of dynamical quarks and the Iwasaki gauge action. We assess finite-size and discretization effects by considering two different lattice spacings and several physical volumes, and mimic the approach to the continuum limit through stereographic projection. Fitting the inflection points of the participation ratios of the overlap Dirac eigenmodes, we obtain estimates of the temperature dependence of the mobility edge, below which quark modes are localized. We observe that it is well described by a quadratic polynomial and systematically vanishes at temperatures below the pseudocritical one of the chiral transition. In fact, our best estimates within errors overlap with that of the chiral phase transition temperature of QCD in the chiral limit. In addition to the published version of this work, more recent results on lower temperatures and gauge configurations smoothed with gradient flow will be presented.

Based on: 10.1103/PhysRevD.109.074512, 10.48550/arXiv.2304.13617

Primary authors: Dr SMITH, Dominik (Justus Liebig University Giessen, Facility for Antiproton and Ion Research in Europe GmbH); VON SMEKAL, Lorenz (Justus Liebig University Giessen, Helmholtz Forschungsakademie Hessen für FAIR); KEHR, Robin (Justus Liebig University Giessen)

Presenter: KEHR, Robin (Justus Liebig University Giessen)

Session Classification: QCD at non-zero temperature

Track Classification: QCD at Non-zero Temperature