

Contribution ID: 303

Type: Talk

## The Hubbard interaction at finite temperature on a Hexagonal lattice

Tuesday, 30 July 2024 16:35 (20 minutes)

The temporal finite volume induces significant effects in Monte Carlo simulations of systems in low dimensions. An example is graphene, a 2-D hexagonal system known for its unique electronic properties and numerous potential applications.

In this work, we explore the behavior of fermions on a graphene sheet with a Hubbard-type interaction characterized by coupling U. This system exhibits zero or near zero-energy excitations that are highly sensitive to finite temperature effects. Therefore, accounting for this dependence is essential to obtaining reliable zerotemperature extrapolations. We compute corrections to the self-energy and the effective mass of low-energy excitations, as well as the shift in ground state energy. These analyses are conducted for both zero and finite temperatures. Our findings reveal that the first-order  $\mathcal{O}(U)$  contributions are absent, leading to non-trivial corrections starting at  $\mathcal{O}(U^2)$ . We validate our calculations against numerical results from Hybrid Monte Carlo simulations on small lattices and extrapolate the behavior at low temperatures.

**Primary author:** RAZMADZE, Lado (Forschungszentrum Juelich)

**Co-authors:** BERKOWITZ, Evan (Forschungszentrum Jülich); OSTMEYER, Johann (Bonn University); LUU, Thomas (Forschungszentrum Jülich / University of Bonn)

**Presenter:** RAZMADZE, Lado (Forschungszentrum Juelich)

Session Classification: Applications outside particle physics

Track Classification: Applications Outside Particle Physics