

Contribution ID: 443

Type: Talk

NRQCD Bottomonium at non-zero temperature using time-derivative moments

Wednesday, 31 July 2024 11:35 (20 minutes)

We will discuss our recent results for the thermal mass and width of bottomonium. A well-known challenge for the lattice community is calculating the spectral function from the Euclidean correlator. We have approximated the spectral function and derived the mass and width of particles through the time derivatives of the lattice correlator.

We have focused on extracting the properties of bottomonium states, specifically Upsilon and Chi_b1, from lattice correlators. We will give an overview of the time-derivative moments approach and look at results for the temperature dependance of the mass and width of both bottomonium states. The zero temperature results are consistent with experimental values, while results at higher temperatures are similar to those obtained using other methods.

Primary author: HOROHAN D'ARCY, Rachel (Maynooth University)

Co-authors: SMECCA, Antonio (Swansea University); JAEGER, Benjamin (University of Southern Denmark); ALL-TON, Chris (Swansea University); AARTS, Gert (Swansea University); SKULLERUD, Jon-Ivar (Maynooth University); LOMBARDO, Maria Paola (INFN Firenze); BIGNELL, Ryan (Trinity College Dublin); KIM, Seyong (Sejong University); RYAN, Sinead (Trinity College Dublin); BURNS, Timothy (Swansea University)

Presenter: HOROHAN D'ARCY, Rachel (Maynooth University)

Session Classification: QCD at non-zero temperature

Track Classification: QCD at Non-zero Temperature