

Contribution ID: 28

Type: Talk

Phase shift in doubly Charmed H-like dibaryon $\Lambda_c \Lambda_c$ scattering at $M_{\pi} \approx 303 MeV$

Tuesday 30 July 2024 11:15 (20 minutes)

To explore the possibility of H-like dibaryon $\Lambda_c \Lambda_c (0^+)$, we proceed calculation on lattice. Two Wilson-Clover ensembles are used which were generated by CLQCD collaboration with similar settings that $m_{\pi} \approx 303$ MeV, $m_D \approx 1.966$ GeV, $m_{D^*} \approx 2.077$ GeV. In this work, finite-volume scattering with single channel Luscher's formula is adopted because the coupling between $\Lambda_c \Lambda_c$ and $\Xi_{cc} N$ is quite small according to our calculation. There is a sign of the CDD zero pole here. By parameterizing the Luscher's scattering matrix, S and D wave phase shift can be given then. However, we find no bound state within a single channel $\Sigma_c \Sigma_c$. Further, the coupling between $\Lambda_c \Lambda_c$ and $\Sigma_c \Sigma_c$ is a little strong.

Primary author: GENG(耿), Yiqi(乙淇)(南京师范大学,中国科学院近代物理所)

Co-authors: LIU, Liuming (IMP); SUN, peng (IMP); WU, Jia-jun (GUCAS); XING, Hanyang (IMP); ZHU, Ruilin (Nanjing Normal U.)

Presenter: GENG(耿), Yiqi(乙淇)(南京师范大学,中国科学院近代物理所)

Session Classification: Structure of hadrons and nuclei

Track Classification: Structure of Hadrons and Nuclei