

Contribution ID: 281

Type: Talk

## Scale setting of $\mathrm{SU}(N)$ Yang–Mills theories via Twisted Gradient Flow

Monday, 29 July 2024 14:35 (20 minutes)

We present preliminary results for the scale setting of SU(N) Yang–Mills theories using twisted boundary conditions and the gradient-flow scale  $\sqrt{t_0}$ . The end goal of this study is to determine the SU(N)  $\Lambda$ -parameter through the step-scaling method. The scale  $\sqrt{t_0}$ , being defined from the flowed action density of the gauge fields, is correlated with their topological charge and thus could be affected by topological freezing. We deal with this problem with the Parallel Tempering on Boundary Conditions algorithm, which we found to be effective for the same numerical setup in previous work.

**Primary authors:** GIORGIERI, Andrea (University of Pisa); BONANNO, Claudio (IFT UAM/CSIC Madrid); Dr DASILVA GOLÁN, Jorge Luis (Physics Department, Brookhaven National Laboratory); GARCIA-PEREZ, Margarita (Instituto de Fisica Teorica UAM-CSIC); D'ELIA, Massimo (University of Pisa and INFN)

Presenter: GIORGIERI, Andrea (University of Pisa)

Session Classification: Vacuum structure and confinement

Track Classification: Vacuum Structure and Confinement