

Contribution ID: 259

Type: Talk

Intrinsic width of the flux tube in 2+1 dimensional Yang-Mills theories

Wednesday 31 July 2024 11:55 (20 minutes)

In this talk we present a study of the flux tube in Yang-Mills theories with a particular focus on its intrinsic width.

In the Effective String Theory description of flux tubes, this quantity is typically neglected, since it has no measurable effects on the inter-quark static potential. However, it can be directly observed in the profile of the flux tube as a deviation from the expected gaussian shape.

In the recent past numerical lattice results on these deviations have been interpreted in terms of a dual superconductor model. We discuss the pros and cons of this proposal and compare it with the results of an effective description of the flux tube based on the Svetitsky-Yaffe mapping of Yang-Mills theories into spin models in the vicinity of the deconfinement transition.

We study in particular the SU(2) and SU(3) models in 2+1 dimensions, which are mapped into the Ising and Z_3 spin models in two dimensions respectively. The main feature of this description is that the intrinsic width is not a new independent quantity but can be related to the same physical scale appearing in the confining inter-quark static potential.

We test this prediction with precise numerical simulations of the SU(2) and SU(3) models, finding a good agreement with the numerical results.

Primary authors: NADA, Alessandro (Università di Torino); PANFALONE, Dario (Universita' di Torino); CELLINI, Elia (University of Turin / INFN Turin); VERZICHELLI, Lorenzo (Università di Torino, INFN sezione di Torino); CASELLE, Michele (Università di Torino)

Presenter: VERZICHELLI, Lorenzo (Università di Torino, INFN sezione di Torino)

Session Classification: Vacuum structure and confinement

Track Classification: Vacuum Structure and Confinement