

Contribution ID: 73

Type: Talk

A proposal for removing πN -state contamination from the nucleon induced pseudoscalar form factor in lattice QCD

Monday 29 July 2024 14:15 (20 minutes)

In the PACS10 project, the PACS collaboration have generated three sets of the PACS10 gauge configurations at the physical point with lattice volume larger than $(10 \text{ fm})^4$ and three different lattice spacings. The isovector nucleon form factors had been already calculated by using two sets of the PACS10 gauge configurations. In our strategy, the smearing parameters of the nucleon interpolation operator were highly optimized to eliminate as much as possible the contribution of excited states in the nucleon two-point function. This strategy was quite successful in calculations of the electric (G_E), magnetic (G_M) and axial-vector (F_A) form factors, while the induced pseudoscalar (F_P) and pseudoscalar (G_P) form factors remained strongly affected by residual contamination of πN -state contribution. In this talk, we propose a simple method to remove the πN -state contamination from the F_P form factor, and then evaluate the induced pseudoscalar charge g_P^* and the pionnucleon coupling $g_{\pi NN}$ from existing data in a new analysis. Applying this method to the G_P form factor will also be considered.

Primary author: SASAKI, Shoichi (Department of Physics, Tohoku University)

Co-authors: AOKI, Yasumichi (RIKEN); ISHIKAWA, Ken-Ichi (Hiroshima University); KURAMASHI, Yoshinobu (University of Tsukuba/RIKEN AICS); SATO, Kohei (University of Tsukuba); SHINTANI, Eigo (University of Tsukuba); TSUJI, Ryutaro (KEK); WATANABE, Hiromasa (YITP, Kyoto University); YAMAZAKI, Takeshi (Univ. of Tsukuba)

Presenter: SASAKI, Shoichi (Department of Physics, Tohoku University)

Session Classification: Structure of hadrons and nuclei

Track Classification: Structure of Hadrons and Nuclei