



Contribution ID: 274

Type: Talk

## Form factor curves consistent with unitarity for semileptonic decays

*Tuesday, 30 July 2024 11:35 (20 minutes)*

We discuss a method to generate form factor curves across the entire kinematic range for semileptonic (SL) pseudoscalar to pseudoscalar decays, for example  $B \rightarrow \pi \mu \nu$  and  $B_s \rightarrow K \mu \nu$ .

The work builds upon the Dispersive Matrix (DM) method. Using known form factor information at specific discrete  $q^2$  points as input, the DM method allows model-independent extrapolation to any desired  $q^2$  value in the SL physical region. Here  $q$  is the outgoing lepton-pair 4-momentum. The main obstacle in using DM results to determine phenomenological predictions, such as forward-backward asymmetry, is that it is not obvious how to exploit the bounds over continuous ranges of  $q^2$  when integrating, for example, the differential decay rate over the physical  $q^2$  range or over bins in  $q^2$ .

Using this method, we can generate a family of curves, each consistent with unitarity constraints, that can be used in the same way as a set generated from a parametrized fit (e.g. a  $z$ -fit). This allows integration over any desired bins. We further show some techniques to increase the computational efficiency of the method.

We demonstrate the application to determining  $V_{ub}$ .

**Primary author:** RADLEY-SCOTT, Callum (University of Southampton)

**Co-authors:** FLYNN, Jonathan (University of Southampton); HUSUNG, Nikolai (IFT UAM/CSIC)

**Presenter:** RADLEY-SCOTT, Callum (University of Southampton)

**Session Classification:** Quark and lepton flavour physics

**Track Classification:** Quark and Lepton Flavour Physics