



Contribution ID: 355

Type: Talk

Virtual radiative Leptonic decays of charged Kaons

Tuesday, 30 July 2024 11:15 (20 minutes)

We present preliminary results for $K \rightarrow l\nu l'^+ l'^-$ decays, which are mediated by an off-shell W boson and a virtual photon. These decays are suppressed in the Standard Model since their rates start at $\mathcal{O}(\alpha_{em}^2)$, making them very interesting in the search for new physics.

We compute the four form factors needed to describe the structure-dependent part of the decay amplitudes using the $N_f = 2 + 1 + 1$ gauge ensembles generated by the Extended Twisted Mass Collaboration (ETMC) at the pion physical point.

We choose several values of both the momentum and the virtuality of the photon k^2 to cover all the allowed kinematic range. In the region above the two-pion threshold ($k^2 > 4m_\pi^2$), where issues related to the analytic continuation are present, we reconstruct the form factors employing the so-called HLT method.

As a byproduct of our analysis, we reassess the form factors entering $K \rightarrow l\nu\gamma$.

Primary author: DI PALMA (ON BEHALF OF THE RM123 COLLABORATION), Roberto (Università Roma Tre)

Presenter: DI PALMA (ON BEHALF OF THE RM123 COLLABORATION), Roberto (Università Roma Tre)

Session Classification: Quark and lepton flavour physics

Track Classification: Quark and Lepton Flavour Physics