

Forward-limit GPDs of the η_c -meson

Jose Manuel Morgado Chávez

With: B. Blossier, C. Mezrag and T. San José

41st Lattice Conference

University of Liverpool, United Kingdom.

July 28th – August 3rd 2024.

Email: jose-manuel.morgadochavez@cea.fr




Motivation: η_c -mesons

η_c meson

Composition: $c\bar{c}$
 J^{PC} : 0^{-+}

Mass: 2983.9 ± 0.4 MeV

Width: 32.0 ± 0.7 MeV



Motivation: η_c -mesons

η_c meson

Composition:

$c\bar{c}$

J^{PC} :


0^{-+}

Mass:

2983.9 ± 0.4 MeV

Width:

32.0 ± 0.7 MeV



η_c -hadron structure

- How does it emerge from the bounding of a pair $c\bar{c}$?
- Comparison with lighter 0^- mesons: Assess quark-mass effect on hadron structure.

Introduction: t -dependent Parton Distribution Functions (tPDFs)

Off-forward parton distribution functions: Non-local, light-like separated, quark or gluon operators, evaluated between hadron states in non-forward kinematics and projected onto the light-front. [Fortsch.Phys.:42(1994)101, Phys.Lett.B:380(1996)417, Phys.Rev.D:55(1997)7114]

Example: Twist-two chiral-even quark GPD of a spinless hadron.

$$H_{q/h}(x, \xi, t, \mu^2) = \frac{1}{2} \int \frac{dy^-}{2\pi} e^{ix \frac{p+p'}{2} \cdot z} \left\langle h(p') \right| \bar{\psi}_q(-z/2) \gamma^+ \widehat{\mathcal{W}}[-z/2, z/2] \psi_q(z/2) \left| h(p) \right\rangle \Big|_{\substack{z^+ = 0 \\ z_\perp = \mathbf{0}_\perp}}$$

Introduction: t -dependent Parton Distribution Functions (tPDFs)

Off-forward parton distribution functions: Non-local, light-like separated, quark or gluon operators, evaluated between hadron states in non-forward kinematics and projected onto the light-front. [Fortsch.Phys.:42(1994)101, Phys.Lett.B:380(1996)417, Phys.Rev.D:55(1997)7114]

Example: Twist-two chiral-even quark GPD of a spinless hadron.

$$H_{q/h}(x, \xi, t, \mu^2) = \frac{1}{2} \int \frac{dy^-}{2\pi} e^{ix \frac{p+p'}{2} \cdot z} \left\langle h(p') \right| \bar{\psi}_q(-z/2) \gamma^+ \widehat{\mathcal{W}}[-z/2, z/2] \psi_q(z/2) \left| h(p) \right\rangle \Big|_{\substack{z^+ = 0 \\ z_\perp = \mathbf{0}_\perp}}$$

t -dependent Parton Distribution Functions

$$q_h(x, t, \mu^2) = H_{q/h}(x, \xi = 0, t, \mu^2) \quad [p \cdot z = p' \cdot z \equiv p^+]$$

Introduction: t -dependent Parton Distribution Functions (tPDFs)

Off-forward parton distribution functions: Non-local, light-like separated, quark or gluon operators, evaluated between hadron states in non-forward kinematics and projected onto the light-front. [Fortsch.Phys.:42(1994)101, Phys.Lett.B:380(1996)417, Phys.Rev.D:55(1997)7114]

Example: Twist-two chiral-even quark GPD of a spinless hadron.

$$H_{q/h}(x, \xi, t, \mu^2) = \frac{1}{2} \int \frac{dy^-}{2\pi} e^{ix \frac{p+p'}{2} \cdot z} \left\langle h(p') \right| \bar{\psi}_q(-z/2) \gamma^+ \widehat{\mathcal{W}}[-z/2, z/2] \psi_q(z/2) \left| h(p) \right\rangle \Big|_{\substack{z^+ = 0 \\ z_\perp = \mathbf{0}_\perp}}$$

t -dependent Parton Distribution Functions

$$q_h(x, t, \mu^2) = H_{q/h}(x, \xi = 0, t, \mu^2) \quad [p \cdot z = p' \cdot z \equiv p^+]$$

Properties:

- Universality *i.e.* hadron-specific objects.
- Contain parton distribution functions and electromagnetic form factors.
- Non-perturbative description of hadron structure: (3D) Tomography.

Project: To compute tPDFs of the η_c -meson

Pseudo-distributions in a nutshell (I)

Definition: Ioffe-time tPDF ($\nu \equiv -p \cdot z$)

[Nucl.Phys.B:311(1989)541, Phys.Rev.D:51(1995)6036, Phys.Rev.D:100(2019)116011]

$$q_h(\nu, t, \mu^2) \equiv \int_{-1}^1 dx e^{i\nu x} q_h(x, t, \mu^2) = \frac{1}{2p^+} \langle h(p') | \bar{\psi}_q(-z/2) \gamma^+ \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle \Big|_{\substack{z^+ = 0 \\ z_\perp = 0_\perp}}$$

Pseudo-distributions in a nutshell (I)

Definition: Ioffe-time tPDF ($\nu \equiv -p \cdot z$)

[Nucl.Phys.B:311(1989)541, Phys.Rev.D:51(1995)6036, Phys.Rev.D:100(2019)116011]

$$q_h(\nu, t, \mu^2) \equiv \int_{-1}^1 dx e^{i\nu x} q_h(x, t, \mu^2) = \frac{1}{2p^+} \langle h(p') | \bar{\psi}_q(-z/2) \gamma^+ \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle \Big|_{\substack{z^+ = 0 \\ z_\perp = 0_\perp}}$$

1. Consider a **generic matrix element** with $z^\mu \in \mathbb{R}^{3,1}$ (or even $z^\mu \in \mathbb{R}^4$)

[Phys.Rev.D:96(2017)034025, Phys.Rev.D:96(2017)094503, Phys.Rev.D:100(2019)116011]

$$\begin{aligned} M_q^\mu(p, p', z) &= \langle h(p') | \bar{\psi}_q(-z/2) \gamma^\mu \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle \\ &= (p + p')^\mu \mathcal{F}(\nu, t, z^2) - (p' - p)^\mu \mathcal{G}(\nu, t, z^2) + z^\mu \mathcal{Z}(\nu, t, z^2), \quad \nu \equiv -p \cdot z \end{aligned}$$

Pseudo-distributions in a nutshell (I)

Definition: Ioffe-time tPDF ($\nu \equiv -p \cdot z$)

[Nucl.Phys.B:311(1989)541, Phys.Rev.D:51(1995)6036, Phys.Rev.D:100(2019)116011]

$$q_h(\nu, t, \mu^2) \equiv \int_{-1}^1 dx e^{i\nu x} q_h(x, t, \mu^2) = \frac{1}{2p^+} \langle h(p') | \bar{\psi}_q(-z/2) \gamma^+ \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle \Big|_{\substack{z^+ = 0 \\ z_\perp = \mathbf{0}_\perp}}$$

1. Consider a **generic matrix element** with $z^\mu \in \mathbb{R}^{3,1}$ (or even $z^\mu \in \mathbb{R}^4$)

[Phys.Rev.D:96(2017)034025, Phys.Rev.D:96(2017)094503, Phys.Rev.D:100(2019)116011]

$$\begin{aligned} M_q^\mu(p, p', z) &= \langle h(p') | \bar{\psi}_q(-z/2) \gamma^\mu \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle \\ &= (p + p')^\mu \mathcal{F}(\nu, t, z^2) - (p' - p)^\mu \mathcal{G}(\nu, t, z^2) + z^\mu \mathcal{Z}(\nu, t, z^2), \quad \nu \equiv -p \cdot z \end{aligned}$$

2. Light-front projection, *i.e.* $z^\mu \propto (1, 0, 0, -1)$ and $\mu = +$

$$M_q^+(p, p', z) \Big|_{\substack{z^+ \rightarrow 0 \\ z_\perp \rightarrow \mathbf{0}_\perp}} = \langle h(p') | \bar{\psi}_q(-y/2) \gamma^+ \hat{\mathcal{W}}[-y/2; y/2] \psi_q(y/2) | h(p) \rangle = 2p^+ \mathcal{F}(\nu, t, z^2 \rightarrow 0)$$

Pseudo-distributions in a nutshell (I)

Definition: Ioffe-time tPDF ($\nu \equiv -p \cdot z$)

[Nucl.Phys.B:311(1989)541, Phys.Rev.D:51(1995)6036, Phys.Rev.D:100(2019)116011]

$$q_h(\nu, t, \mu^2) \equiv \int_{-1}^1 dx e^{i\nu x} q_h(x, t, \mu^2) = \frac{1}{2p^+} \langle h(p') | \bar{\psi}_q(-z/2) \gamma^+ \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle \Big|_{\substack{z^+ = 0 \\ z_\perp = \mathbf{0}_\perp}}$$

1. Consider a **generic matrix element** with $z^\mu \in \mathbb{R}^{3,1}$ (or even $z^\mu \in \mathbb{R}^4$)

[Phys.Rev.D:96(2017)034025, Phys.Rev.D:96(2017)094503, Phys.Rev.D:100(2019)116011]

$$\begin{aligned} M_q^\mu(p, p', z) &= \langle h(p') | \bar{\psi}_q(-z/2) \gamma^\mu \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle \\ &= (p + p')^\mu \mathcal{F}(\nu, t, z^2) - (p' - p)^\mu \mathcal{G}(\nu, t, z^2) + z^\mu \mathcal{Z}(\nu, t, z^2), \quad \nu \equiv -p \cdot z \end{aligned}$$

2. Light-front projection, *i.e.* $z^\mu \propto (1, 0, 0, -1)$ and $\mu = +$

$$M_q^+(p, p', z) \Big|_{\substack{z^+ \rightarrow 0 \\ z_\perp \rightarrow \mathbf{0}_\perp}} = \langle h(p') | \bar{\psi}_q(-y/2) \gamma^+ \hat{\mathcal{W}}[-y/2; y/2] \psi_q(y/2) | h(p) \rangle = 2p^+ \mathcal{F}(\nu, t, z^2 \rightarrow 0)$$

3. Connection with Ioffe-time distributions

$$q_h(\nu, t, \mu^2) = \mathcal{F}(\nu, t, z^2 \rightarrow 0)$$

Pseudo-distributions in a nutshell (II)

Pseudo-tPDF: $\tilde{q}_h(\nu, t, z^2) = \mathcal{F}(\nu, t, z^2) \Big|_R$, $\lim_{z^2 \rightarrow 0} \tilde{q}_h(\nu, t, z^2) \rightarrow q_h(\nu, t, \mu^2)$

Pseudo-distributions in a nutshell (II)

$$\text{Pseudo-tPDF: } \tilde{q}_h(\nu, t, z^2) = \mathcal{F}(\nu, t, z^2) \Big|_{\mathbf{R}}, \quad \lim_{z^2 \rightarrow 0} \tilde{q}_h(\nu, t, z^2) \rightarrow q_h(\nu, t, \mu^2)$$

Euclidean setup:

- $p^\mu = (E, \mathbf{p}_\perp, p_3)$ and $p'^\mu = (E, -\mathbf{p}_\perp, p_3)$
- $z^\mu = (0, 0, 0, z_3)$

Pseudo-distributions in a nutshell (II)

$$\text{Pseudo-tPDF: } \tilde{q}_h(\nu, t, z^2) = \mathcal{F}(\nu, t, z^2) \Big|_{\mathbf{R}}, \quad \lim_{z^2 \rightarrow 0} \tilde{q}_h(\nu, t, z^2) \rightarrow q_h(\nu, t, \mu^2)$$

Euclidean setup:

- $p^\mu = (E, \mathbf{p}_\perp, p_3)$ and $p'^\mu = (E, -\mathbf{p}_\perp, p_3)$
- $z^\mu = (0, 0, 0, z_3)$

1. Compute

$$M^0(p, p', z) = \langle h(p') | \bar{\psi}_q(-z/2) \gamma^0 \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle = 2E\mathcal{F}(\nu, t, z^2)$$

Pseudo-distributions in a nutshell (II)

$$\text{Pseudo-tPDF: } \tilde{q}_h(\nu, t, z^2) = \mathcal{F}(\nu, t, z^2) \Big|_{\text{R}}, \quad \lim_{z^2 \rightarrow 0} \tilde{q}_h(\nu, t, z^2) \rightarrow q_h(\nu, t, \mu^2)$$

Euclidean setup:

- $p^\mu = (E, \mathbf{p}_\perp, p_3)$ and $p'^\mu = (E, -\mathbf{p}_\perp, p_3)$
- $z^\mu = (0, 0, 0, z_3)$

1. Compute

$$M^0(p, p', z) = \langle h(p') | \bar{\psi}_q(-z/2) \gamma^0 \hat{\mathcal{W}}[-z/2; z/2] \psi_q(z/2) | h(p) \rangle = 2E\mathcal{F}(\nu, t, z^2)$$

2. Renormalize through a RGI ratio

[Phys. Lett. B: 767 (2017) 314]

$$\mathcal{F}(\nu, t, z^2) \Big|_{\text{R}} \equiv \frac{M^0(p, p', z)}{M^0(0, 0, z)} \frac{M^0(0, 0, 0)}{M^0(p, p', 0)} = \frac{\mathcal{F}(\nu, t, z^2)}{\mathcal{F}(0, 0, z^2)} \frac{\mathcal{F}(0, 0, 0)}{\mathcal{F}(0, t, 0)} = \tilde{q}_h(\nu, t, z^2) + \text{h.t.}$$

Lattice QCD calculation

Numerical setup

- $N_f = 2$ ensembles (CLS) [Nucl.Phys.B:865(2012)397, PoSLATTICE2013:(2014)475]
 - Wilson gauge action.
 - $\mathcal{O}(a)$ -improved Wilson fermions.
 - $\kappa_u = \kappa_d \equiv \kappa_k$.
 - No Symanzik improvement for M^μ .

id	β	a [fm]	L/a	am_π	m_π [MeV]	$m_\pi L$	κ_c	κ_l
A5	5.2	0.0755(9)(7)	32	0.1265(8)	331	4.0	0.12531	0.13594
E5	5.3	0.0658(7)(7)	32	0.1458(3)	437	4.7	0.12724	0.13625
F7			48	0.0885(3)	265	4.3	0.12713	0.13638
N6	5.5	0.0486(4)(5)	48	0.0838(2)	340	4.0	0.13026	0.13667

- One hadron-interpolator and four smearings (source and sink).


$$\eta_c^s(x) = \psi_c^s(x) \gamma_5 \psi_c^s(x) \quad , \quad J^{PC} = 0^{-+}$$
$$\psi_q^s(x) = (1 + 0.125 \Delta_{\text{APE}})^{N_s} \psi_q(x) \quad , \quad N_s \in \{0, 30, 50, 80\}$$

- Twisted boundary conditions and a symmetric frame.

Matrix elements from LQCD

Computation of hadron three-point functions



$$C_3^{(ss')}(\vec{p}, t_{src}) = \sum_{\vec{x}, \vec{z}} e^{-i\vec{p}\cdot\vec{x} - i\vec{q}\cdot\vec{z}} \langle \eta_c^s(\vec{x}, t_{src}) \bar{\psi}_c(\vec{z} - \Delta\vec{z}, \tau) \gamma^0 \hat{\mathcal{W}}[\vec{z} - \Delta\vec{z}, \tau; \vec{z}, \tau] \psi_c(\vec{z}, \tau) \bar{\eta}_c^s(\vec{0}, 0) \rangle$$



- Consider connected diagrams only: Sequential propagator technique.
- Project ground state ($\eta_c(1s)$) according to GEVP.
- Compute ratios to isolate matrix elements: [PoSLATTICE2005:(2006)360]

$$R(\tau) = \frac{C_3^{(P)}(\vec{p}, \vec{p}', t_{src}, \tau)}{\sqrt{C_2^{(P)}(\vec{p}', t_{src}) C_2^{(P)}(\vec{p}, t_{src})}} \sqrt{\frac{C_2^{(P)}(\vec{p}, t_{src} - \tau) C_2^{(P)}(\vec{p}', \tau)}{C_2^{(P)}(\vec{p}', t_{src} - \tau) C_2^{(P)}(\vec{p}, \tau)}} \rightarrow \frac{M^0(p, p', z)}{4\sqrt{E(\vec{p}) E(\vec{p}')}}$$

Pseudo-tPDF data



- Pseudo-distribution data approaching a universal line for all ensembles.
- Signal-to-noise ratio significantly good over the entire kinematic range.

Light-cone matching

From pseudo- to light-cone distributions (I)

Light-cone matching: $\lim_{z^2 \rightarrow 0} \tilde{q}_h(\nu, t, z^2) \rightarrow q_h(\nu, t, \mu^2)$

From pseudo- to light-cone distributions (I)

1. Light-cone operator product expansion: $z^2 \rightarrow 0$

[Nucl.Phys.B:27(1971)541]

$$\begin{aligned}\tilde{q}_h(\nu, t, z^2) &\sim \sum_i \sum_{n=0}^{\infty} \mathcal{C}_n^{(i)}(z^2) \langle h(p') | \tilde{\mathcal{O}}_{(i)}^{\{\mu\mu_1\cdots\mu_n\}} | h(p) \rangle z_\mu \prod_{k=1}^n z_{\mu_k} + \text{h.t.} \\ &= \sum_i \sum_{n=0}^{\infty} \mathcal{C}_n^{(i)}(z^2) (-2)^n \tilde{a}_n^{(i)}(t, z^2) \nu^n + \text{h.t.}\end{aligned}$$

With $\tilde{a}_n^{(i)}(t, z^2) = (1 + z^2 \tilde{f}_n^{(i)}(\nu, t, z^2, m^2)) \tilde{a}_n^{(i)}(t, \log z^2)$, $\tilde{f}_n^{(i)}$ include target-mass correct.

From pseudo- to light-cone distributions (I)

1. Light-cone operator product expansion: $z^2 \rightarrow 0$

[Nucl.Phys.B:27(1971)541]

$$\begin{aligned}\tilde{q}_h(\nu, t, z^2) &\sim \sum_i \sum_{n=0}^{\infty} \mathcal{C}_n^{(i)}(z^2) \langle h(p') | \tilde{\mathcal{O}}_{(i)}^{\{\mu\mu_1\cdots\mu_n\}} | h(p) \rangle z_\mu \prod_{k=1}^n z_{\mu_k} + \text{h.t.} \\ &= \sum_i \sum_{n=0}^{\infty} \mathcal{C}_n^{(i)}(z^2) (-2)^n \tilde{a}_n^{(i)}(t, z^2) \nu^n + \text{h.t.}\end{aligned}$$

With $\tilde{a}_n^{(i)}(t, z^2) = (1 + z^2 \tilde{f}_n^{(i)}(\nu, t, z^2, m^2)) \tilde{a}_n^{(i)}(t, \log z^2)$, $\tilde{f}_n^{(i)}$ include targe-mass correct.

2. Recast as convolution: Light-cone matching to $\overline{\text{MS}}$ distribution

[Phys.Rev.D:98(2018)014019, Phys.Rev.D:98(2018)050004, Phys.Rev.D:97(2018)074508]

$$\tilde{q}_h(\nu, t, z^2) = \int_0^1 dw \mathcal{C}(w, z^2 \mu^2, \alpha_s) q_h(w\nu, t, \mu^2)$$

From pseudo- to light-cone distributions (I)

1. Light-cone operator product expansion: $z^2 \rightarrow 0$

[Nucl.Phys.B:27(1971)541]

$$\begin{aligned}\tilde{q}_h(\nu, t, z^2) &\sim \sum_i \sum_{n=0}^{\infty} \mathcal{C}_n^{(i)}(z^2) \langle h(p') | \tilde{\mathcal{O}}_{(i)}^{\{\mu\mu_1\cdots\mu_n\}} | h(p) \rangle z_\mu \prod_{k=1}^n z_{\mu_k} + \text{h.t.} \\ &= \sum_i \sum_{n=0}^{\infty} \mathcal{C}_n^{(i)}(z^2) (-2)^n \tilde{a}_n^{(i)}(t, z^2) \nu^n + \text{h.t.}\end{aligned}$$

With $\tilde{a}_n^{(i)}(t, z^2) = (1 + z^2 \tilde{f}_n^{(i)}(\nu, t, z^2, m^2)) \tilde{a}_n^{(i)}(t, \log z^2)$, $\tilde{f}_n^{(i)}$ include targe-mass correct.

2. Recast as convolution: Light-cone matching to $\overline{\text{MS}}$ distribution

[Phys.Rev.D:98(2018)014019, Phys.Rev.D:98(2018)050004, Phys.Rev.D:97(2018)074508]

$$\tilde{q}_h(\nu, t, z^2) = \int_0^1 dw \mathcal{C}(w, z^2 \mu^2, \alpha_s) q_h(w\nu, t, \mu^2)$$

Challenges of the light-cone matching

- Integral relation
- Acts on light-cone distributions

From pseudo- to light-cone distributions (II)

Solution: Ioffe-time distributions, onto and outside the light-front, are analytic functions in the Ioffe-time variable:

$$q_h(\nu, t, \mu^2) = \sum_{m=0}^{\infty} a_m(t, \mu^2) \nu^m, \quad \tilde{q}_h(\nu, t, z^2) = \sum_{m=0}^{\infty} \tilde{a}_m(t, z^2) \nu^m.$$

- Matching becomes a standard product

$$\tilde{a}_m(t, z^2) = \mathcal{C}_m(z^2 \mu^2, \alpha_s) a_m(t, \mu^2), \quad \mathcal{C}_m(z^2 \mu^2) \equiv \int_0^1 dw w^m \mathcal{C}(w, z^2 \mu^2, \alpha_s).$$

- Expansion coefficients identified in LC-OPE: Inclusion of target-mass corrections

$$\tilde{a}_m^{(i)}(t, z^2) \rightarrow (1 + z^2 \tilde{f}_m^{(i)}) \tilde{a}_m^{(i)}(t, \log z^2)$$

- Connection with Mellin moments

$$\mathfrak{m}_m^{(i)}(t, \mu^2) = i^m m! a_m^{(i)}(t, \mu^2)$$

From pseudo- to light-cone distributions (III)

Strategy:

1. Fit pseudo-distribution data to

$$\tilde{q}_h(\nu, t, z^2) = \sum_{m=0}^N \left(\tilde{A}_m(t, z^2) + z^2 \tilde{B}_m(t, z^2) \right) \nu^m$$

From pseudo- to light-cone distributions (III)

Strategy:

1. Fit pseudo-distribution data to

$$\tilde{q}_h(\nu, t, z^2) = \sum_{m=0}^N \left(\tilde{A}_m(t, z^2) + z^2 \tilde{B}_m(t, z^2) \right) \nu^m \xrightarrow{\text{Cut-off}} \tilde{A}_m(t, z^2) = \tilde{A}_m(t, z^2) \Big|_{\text{Cont.}} + a \tilde{\mathcal{A}}_m(t, z^2)$$

and extract as many coefficients as possible.

From pseudo- to light-cone distributions (III)

Strategy:

1. Fit pseudo-distribution data to

$$\tilde{q}_h(\nu, t, z^2) = \sum_{m=0}^N \left(\tilde{A}_m(t, z^2) + z^2 \tilde{B}_m(t, z^2) \right) \nu^m \xrightarrow{\text{Cut-off}} \tilde{A}_m(t, z^2) = \tilde{A}_m(t, z^2) \Big|_{\text{Cont.}} + a \tilde{\mathcal{A}}_m(t, z^2)$$

and extract as many coefficients as possible.

2. Match coefficients to the light-cone

$$\tilde{A}_m(t, z^2) \Big|_{\text{Cont.}} \longrightarrow \tilde{a}_m(t, z^2); \quad a_m(t, \mu^2) = \frac{\tilde{a}_m(t, z^2)}{\mathcal{C}_m(z^2 \mu^2, \alpha_s)}$$

From pseudo- to light-cone distributions (III)

Strategy:

1. Fit pseudo-distribution data to

$$\tilde{q}_h(\nu, t, z^2) = \sum_{m=0}^N \left(\tilde{A}_m(t, z^2) + z^2 \tilde{B}_m(t, z^2) \right) \nu^m \xrightarrow{\text{Cut-off}} \tilde{A}_m(t, z^2) = \tilde{A}_m(t, z^2) \Big|_{\text{Cont.}} + a \tilde{\mathcal{A}}_m(t, z^2)$$

and extract as many coefficients as possible.


2. Match coefficients to the light-cone

$$\tilde{A}_m(t, z^2) \Big|_{\text{Cont.}} \longrightarrow \tilde{a}_m(t, z^2); \quad a_m(t, \mu^2) = \frac{\tilde{a}_m(t, z^2)}{\mathcal{C}_m(z^2 \mu^2, \alpha_s)}$$


3. Reconstruct light-cone Ioffe-time $\overline{\text{MS}}$ distribution accordingly

$$q_h(\nu, t, \mu^2) = \sum_{m=0}^N a_m(t, \mu^2) \nu^m$$

Ioffe-time tPDFs: Non-singlet distribution



Ioffe-time tPDFs (II)



Conclusions and future steps

Summary

- Ongoing study of η_c -meson's structure through GPDs within lattice QCD.
- Study light-cone matching in Ioffe-time space:
 - Model bias reduction to truncation error.
 - Systematic extraction of arbitrary order Mellin moments.
 - Transparent treatment of target mass corrections.
 - Possible improvement in matching.

Future steps

- Extend analysis to the non-singlet sector.
- Extend kinematics and statistics: t -values and new ensembles.
- Refine analysis of lattice artifacts: e.g. finite volume or quark masses.
- Reconstruction of x -space distributions.


Thank you!

Back-up slides

t-dependent Parton Distribution Functions (tPDFs)


Hadron structure

How do quarks and gluons combine to make hadrons up?



Generalized Bjorken limit
 $Q^2 \rightarrow \infty$ with $Q^2 >> t$
and ξ fixed.



Factorization
[Phys. Rev. D59(1999)074009]



$$\mathcal{H}(\xi, t, Q^2) = \sum_{p=q,g} \int_{-1}^1 \frac{dx}{\xi} \mathcal{K}^p \left(\frac{x}{\xi}, \frac{Q^2}{\mu_F^2}, \alpha_s(\mu_F^2) \right) \textcolor{red}{H^p(x, \xi, t, \mu_F^2)}$$

Generalized Parton distributions: **Off-forward parton distribution functions**

Two-point functions: Effective masses





Energy spectrum compatible with expectation within finite-volume and cut-off effects.

Systematics: - Fit range: Model averaging (AIC) [Phys. Rev. D:103(2021)114502]



- Excited states: GEVP [Nucl. Phys. B:259(1985)58, JHEP:04(2009)094]

Two-point functions: Dispersion relations



Consistency check: Expected energy-momentum dispersion relations fulfilled.

Three-point functions: Ratio fits



Three-point functions

