



Contribution ID: 126

Type: Talk

Heavy-light Meson Decay Constants and Hyperfine Splittings with the Heavy-HISQ Method

Tuesday, 30 July 2024 16:55 (20 minutes)

We present preliminary new lattice QCD results for high-precision vector-to-pseudoscalar and tensor-to-vector ratios of decay constants of the $B^{(*)}$, $D^{(*)}$, $B_s^{(*)}$ and $D_s^{(*)}$ mesons, in which many uncertainties cancel, using the heavy-HISQ method. We use the Highly Improved Staggered Quark (HISQ) action for all valence quarks and second generation MILC $n_f = 2 + 1 + 1$ HISQ gluon field configurations, with lattice spacings ranging from 0.15 fm down to 0.045 fm. Our pion masses range from ≈ 300 MeV down to the physical value, and our heavy quark masses range from the physical charm up to the physical bottom on the three finest ensembles. These quantities enable powerful tests of Standard Model flavour phenomenology and aid the search for new physics through their sensitivity to beyond-Standard-Model effects.

Primary authors: SMECCA, Antonio (Swansea University); DAVIES, Christine (University of Glasgow); Dr HARRISON, Judd (University of Glasgow); MILLER, Kerr (University of Glasgow)

Presenter: MILLER, Kerr (University of Glasgow)

Session Classification: Quark and lepton flavour physics

Track Classification: Quark and Lepton Flavour Physics