



Contribution ID: 388

Type: Talk

Valence leading isospin breaking contributions to $a_\mu^{\text{HVP-LO}}$

Thursday, 1 August 2024 10:20 (20 minutes)

In the framework of the ongoing computation by ETMC of the leading-order hadronic vacuum polarization (HVP) contribution to the muon anomalous magnetic moment a_μ^{HVP} in QCD+QED, we present preliminary results about the valence quark-connected isospin-breaking corrections to $a_\mu^{\text{HVP-LO}}$, at leading order in $\alpha_{\text{em}} = e^2/4\pi$ and $(m_d - m_u)/\Lambda_{\text{QCD}}$. In our computation, we employ the RM123 approach to QCD+QED and here we focus on two different volumes ($L \sim 3.8$ fm and $L \sim 5.1$ fm) and a fixed lattice spacing (corresponding to $a_{\text{isoQCD}} \sim 0.07951(4)$ fm).

Primary author: EVANGELISTA, Antonio (University and INFN of Rome Tor Vergata)

Co-author: COLLABORATION, ETM

Presenter: EVANGELISTA, Antonio (University and INFN of Rome Tor Vergata)

Session Classification: Quark and lepton flavour physics

Track Classification: Quark and Lepton Flavour Physics