



Contribution ID: 123

Type: Talk

## $\pi^0 \rightarrow \gamma^* \gamma^*$ transition form factor and the pion pole contribution to $a_\mu$ on CLS ensembles

*Monday, 29 July 2024 11:35 (20 minutes)*

We present the status of the Mainz group's lattice QCD calculation of the pion transition form factor  $\mathcal{F}_{\pi^0 \gamma^* \gamma^*}$ , which describes the interaction of an on-shell pion with two off-shell photons. This form factor is the main ingredient in the calculation of the pion-pole contribution to hadronic light-by-light scattering in the muon  $g - 2$ .

We use the  $N_f = 2 + 1$  CLS gauge ensembles, and we update our previous work by including a physical pion mass ensemble (E250). We compute the transition form factor in the pion rest frame as well as in a moving frame in order to have access to a wider range of photon virtualities. In addition to the quark-line connected correlator we also compute the quark-line disconnected diagrams that contribute to the form factor.

In this final stage of the analysis, we combine the result on E250 with the previous work published in 2019 to extrapolate the form factor to the continuum and to physical quark masses. Testing different Ansätze for the fit, we explore the systematic uncertainties of the extrapolation. The contribution from the disconnected diagrams is also scrutinized.

**Primary authors:** GERARDIN, Antoine (Centre de Physique Théorique de Marseille); VON HIPPEL, Georg (University of Mainz); MEYER, Harvey B. (University of Mainz); KOPONEN, Jonna (JGU Mainz); OTTNAD, Konstantin (University of Mainz)

**Presenter:** KOPONEN, Jonna (JGU Mainz)

**Session Classification:** Quark and lepton flavour physics

**Track Classification:** Quark and Lepton Flavour Physics