



Contribution ID: 469

Type: Talk

Quark-mass dependence of the $\Delta(1232)$ resonance parameters.

Wednesday, 31 July 2024 11:15 (20 minutes)

We present results for the $N\pi$ finite-volume spectra on three different lattice volumes with fixed lattice spacing $a = 0.116$ fm and m_π ranging from 249 MeV to 137 MeV. The calculations employ $N_f = 2 + 1$ doubly-HEX-smearred clover fermions. We perform the calculations with total momenta up to $\vec{P} = (1, 1, 1) \frac{2\pi}{L}$ and all relevant irreps. Using the Lüscher method we fit Breit-Wigner parametrizations of the $N\pi$ scattering amplitude to the spectra to obtain the mass and decay width of the lightest baryon resonance, the $\Delta(1232)$.

Primary author: PAUL, Srijit (University of Maryland)

Co-authors: Dr POCHINSKY, Andrew (MIT); PITTLER, Ferenc (The Cyprus Institute); Prof. NEGELE, John W (MIT); LESKOVEC, Luka (University of Ljubljana); PETSCHLIES, Marcus (U. Bonn); SYRITSYN, Sergey (Stony Brook University); KRIEG, Stefan (JSC, Forschungszentrum Juelich/HISKP, University of Bonn); MEINEL, Stefan (University of Arizona / RIKEN BNL Research Center)

Presenter: PAUL, Srijit (University of Maryland)

Session Classification: Hadronic and nuclear spectrum and interactions

Track Classification: Hadronic and Nuclear Spectrum and Interactions