



Contribution ID: 437

Type: Talk

## Towards glueball scattering in lattice Yang-Mills theory

*Thursday, 1 August 2024 09:40 (20 minutes)*

We present progress in the determination of glueball masses and scattering parameters computed using lattice  $SU(3)$  Yang-Mills theory. Focusing on the vacuum-quantum-number ( $J^{PC} = 0^{++}$ ) sector, the computation makes use of a number of techniques to improve the extraction of energies from Euclidean correlators, including anisotropic lattices, the multi-level algorithm, and a large set of operators applied in a variational method. In addition to various algorithmic considerations, our main result is a volume-dependent extraction of the ground state energy. This enables a prediction of the mass in the infinite volume limit and the trilinear coupling governing the coefficient of the leading volume-dependent exponential. This result is a first step towards using excited finite-volume states to extract the glueball scattering amplitude, with potential applications in QCD and strongly interacting dark matter models.

**Primary authors:** Prof. RAGO, Antonio (QTC & IMADA, University of Southern Denmark); BRUNO, Mattia (Universita' di Milano-Bicocca); HANSEN, Maxwell (University of Mainz)

**Presenter:** HANSEN, Maxwell (University of Mainz)

**Session Classification:** Hadronic and nuclear spectrum and interactions

**Track Classification:** Hadronic and Nuclear Spectrum and Interactions