

Contribution ID: 74

Type: Talk

Tetraquarks $\overline{b}\overline{b}ud$, $I(J^P) = 0(1^-)$ and $\overline{b}\overline{c}ud$ with $I(J^P) = 0(0^+), 0(1^+)$ from Lattice QCD Static Potentials

Tuesday, 30 July 2024 14:25 (20 minutes)

We use existing antistatic-antistatic potentials computed with two flavors of twisted mass fermions to study antiheavy-antiheavy-light-light four-quark systems in the Born-Oppenheimer approximation. We do not only compute masses of bound states, but also search for poles in the scattering amplitude above the lowest meson-meson threshold, to detect resonances and determine their parameters. We include effects due to the heavy quark spins, by considering coupled channel Schroedinger equations and using the experimentally measured mass splittings of the *B* and *B*^{*} meson as input. We present indications for a $\bar{b}\bar{b}ud$ tetraquark resonance with quantum numbers $I(J^P) = 0(1^-)$ close to the B^*B^* threshold. We also discuss first results for $\bar{b}\bar{c}ud$ systems with $I(J^P) = 0(0^+)$ and $I(J^P) = 0(1^+)$.

Primary author: HOFFMANN, Jakob (Goethe University, Frankfurt am Mian)

Co-authors: Mr MÜLLER, Lasse (Goethe University, Frankfurt am Mian); Prof. WAGNER, Marc (Goethe University, Frankfurt am Mian)

Presenter: HOFFMANN, Jakob (Goethe University, Frankfurt am Mian)

Session Classification: Hadronic and nuclear spectrum and interactions

Track Classification: Hadronic and Nuclear Spectrum and Interactions