



Contribution ID: 61

Type: Talk

Four-quark operators with $\Delta F = 2$ in the GIRS scheme

Friday, 2 August 2024 14:35 (20 minutes)

We calculate the mixing matrices of four-quark operators that change flavor numbers by two units. Our approach employs two schemes: the coordinate-space Gauge Invariant Renormalization Scheme (GIRS) and the Modified Minimal Subtraction scheme. From our perturbative computations, we extract the conversion factors between these two renormalization schemes at the next-to-leading order. A significant challenge in the study of four-quark operators is that they mix among themselves upon renormalization. Additionally, computations in GIRS at a given order in perturbation theory require Feynman diagrams with at least one additional loop. The extraction of the conversion factors involves calculating two-point Green's functions, which include products of two four-quark operators, and three-point Green's functions, which involve one four-quark operator and two bilinear operators, with all operators located at distinct spacetime points. Furthermore, we focus on both parity-conserving and parity-violating four-quark operators. This calculation is crucial for determining Cabibbo–Kobayashi–Maskawa (CKM) matrix elements from numerical simulations using the GIRS scheme.

Primary authors: Dr SPANOUEDES, Gregoris (University of Cyprus); PANAGOPOULOS, Haralambos (University of Cyprus); Mr HERODOTOU, Herodotos (University of Cyprus); KOSTA, Marios (University of Cyprus, Rinnoco Ltd); Dr CONSTANTINO, Martha (Temple University)

Presenter: KOSTA, Marios (University of Cyprus, Rinnoco Ltd)

Session Classification: Quark and lepton flavour physics

Track Classification: Quark and Lepton Flavour Physics