



Contribution ID: 161

Type: Talk

η invariant of massive Wilson Dirac operator and the index

Tuesday, 30 July 2024 11:35 (20 minutes)

We mathematically show an equality between the index of a Dirac operator on a flat continuum torus and the η invariant of the Wilson Dirac operator with a negative mass when the lattice spacing is sufficiently small. Unlike the standard approach, our formulation using the K -theory does not require the Ginsparg-Wilson relation or the modified chiral symmetry on the lattice. We prove that a one-parameter family of continuum massive Dirac operators and the corresponding Wilson Dirac operators belong to the same equivalence class of the K^1 group at a finite lattice spacing. Their indices, which are evaluated by the spectral flow or equivalently by the η invariant at finite masses, are proved to be equal.

Primary authors: AOKI, Shoto (The University of Tokyo); FUKAYA, Hidenori (Osaka Univ.); Prof. FURUTA, Mikio (U. of Tokyo); Prof. MATSUO, Shinichiroh (Nagoya Univ.); ONOGI, Tetsuya (Osaka University); Prof. YAMAGUCHI, Satoshi (Osaka Univ.)

Presenter: FUKAYA, Hidenori (Osaka Univ.)

Session Classification: Theoretical developments

Track Classification: Theoretical Developments