

Lattice Weyl Fermion on a single spherical domain-wall 1

Shoto Aoki The Univ. of Tokyo
with Hidenori Fukaya, Naoto Kan

August 2, 2024 @Liverpool
arXiv:2402.09774

————→ Kan's Talk is in the next session!



Contents

Introduction


S^2 domain-wall in \mathbb{R}^3

S^2 domain-wall on $3D$ square lattice

Summary

Motivation

Every curved manifold can be isometrically embedded into some higher-dimensional Euclidean spaces.



Edge modes are localized at the curved domain-wall.

= They feel "gravity" by the equivalence principle.

→ Are they chiral?

Embedding a Curved Space [Nash, 1956]

For any n -dim. Riemann space (Y, g) , there is an embedding $f : Y \rightarrow \mathbb{R}^m$ ($m \gg n$) such that Y is identified as

$$x^\mu = x^\mu(y^1, \dots, y^n) \quad (\mu = 1, \dots, m)$$

$$\begin{pmatrix} x^\mu & : & \text{Cartesian coordinates of } \mathbb{R}^m \\ y^i & : & \text{Coordinates of } Y \end{pmatrix}$$


and the metric is written as

$$g_{ij} = \sum_{\mu\nu} \delta_{\mu\nu} \frac{\partial x^\mu}{\partial y^i} \frac{\partial x^\nu}{\partial y^j}.$$

————→ Vielbein and spin connection are also induced!

Any Riemannian manifold can be identified as a submanifold of a flat Euclidean space!

“Gravity” in Condensed Matter Physics



[Onoe et al., 2012] observed a gravitational effect on 1D uneven peanut-shaped C_{60} polymer.

The Hamiltonian on a curved surface is

[Jensen and Koppe, 1971; da Costa, 1981]

$$H = -\frac{\hbar^2}{2m_*} \left[\frac{1}{\sqrt{g}} \partial_i (\sqrt{g} g^{ij} \partial_j) + h^2 - k \right], \quad \begin{cases} h: \text{Extrinsic curvature} \\ k: \text{Gaussian curvature} \end{cases}$$


→ Density of states depends on the curvatures.

Weyl Fermion on Single Spherical Domain-wall cf. [Sen's Talk]

We investigate a Free fermion system with S^2 domain-wall.

$$D = \sum_{i=1}^3 \sigma^i \frac{\partial}{\partial x^i} - m$$
$$\rightarrow \mathbb{D}^{S^2} \frac{1}{2} (1 + \sigma^3)$$

- Spectrum
- Edge modes
- Continuum Limit
- Restoration of Symmetry



Kan will assign $U(1)$ gauge connection and talk about an obstacle in formulating lattice chiral gauge theory.

Contents

Introduction

S^2 domain-wall in \mathbb{R}^3

S^2 domain-wall on 3D square lattice

Summary

Contents

Introduction


S^2 domain-wall in \mathbb{R}^3

S^2 domain-wall on 3D square lattice

Summary

Domain wall:

$$m(r) = \begin{cases} -m & (r < r_0) \\ +M \rightarrow +\infty & (r \geq r_0) \end{cases}$$



Dirac operator:

$$D = \sum_{i=1}^3 \sigma_i \frac{\partial}{\partial x^i} + m(r) = \sigma_r \left(\frac{\partial}{\partial r} + \frac{1}{r} \right) + m(r) - \sigma_r \frac{D^{S^2}}{r},$$

$$D^{S^2} = \sum_{i=1}^3 \sigma^i L_i + 1, \quad \sigma_r = \frac{x}{r} \sigma_1 + \frac{y}{r} \sigma_2 + \frac{z}{r} \sigma_3.$$

Boundary condition: $\sigma_r \psi(x) = \pm \psi(x)$ at $r = r_0$

Effective Dirac op and Dirac op. of S^2

A local Lorentz transformation

$$R = \begin{pmatrix} e^{-i\frac{\phi}{2}} \cos(\frac{\theta}{2}) & -e^{-i\frac{\phi}{2}} \sin(\frac{\theta}{2}) \\ e^{i\frac{\phi}{2}} \sin(\frac{\theta}{2}) & e^{i\frac{\phi}{2}} \cos(\frac{\theta}{2}) \end{pmatrix} e^{i\frac{\phi}{2}}$$

changes $\psi \rightarrow R^{-1}\psi$ and

$$D^{S^2} \rightarrow i \left(\sigma_1 \frac{\partial}{\partial \theta} + \frac{\sigma_2}{\sin \theta} \left(\frac{\partial}{\partial \phi} + \underbrace{\frac{i}{2} - \frac{\cos \theta}{2} \sigma_1 \sigma_2}_{\text{Spin}^c \text{ connection on } S^2} \right) \right),$$

$$\sigma_r \rightarrow \sigma_3$$

Edge states feel gravity through the induced connection!

[Takane and Imura, 2013].

Eigenstate of $D^\dagger D$ and DD^\dagger

Let χ_\pm satisfy

$$D^{S^2} \chi_\pm = \lambda \chi_{\mp}, \quad (\lambda = 1, 2, \dots)$$

$$\sigma_r \chi_\pm = \pm \chi_\pm.$$

In the large m limit, we assume $\psi_\pm = \frac{1}{r} e^{-m|r-r_0|} \chi_\pm$, then we get

$$D\psi_+ = \left(\sigma_r \left(\frac{\partial}{\partial r} + \frac{1}{r} \right) + m\epsilon - \sigma_r \frac{D^{S^2}}{r} \right) \psi_+ \simeq \frac{\lambda}{r_0} \psi_-.$$

$$D^\dagger \psi_- = \left(-\sigma_r \left(\frac{\partial}{\partial r} + \frac{1}{r} \right) + m\epsilon + \sigma_r \frac{D^{S^2}}{r} \right) \psi_- \simeq \frac{\lambda}{r_0} \psi_+.$$

$$\longrightarrow D^\dagger D\psi_+ = \left(\frac{\lambda}{r_0} \right)^2 \psi_+ \quad \text{and} \quad DD^\dagger \psi_- = \left(\frac{\lambda}{r_0} \right)^2 \psi_-$$

Chiral fermion appears at the wall!

Contents

Introduction

S^2 domain-wall in \mathbb{R}^3

S^2 domain-wall on 3D square lattice

Summary

S^2 Domain-wall Fermion on Lattice

We consider a lattice on B^3 with the radius r_0 .


The (Wilson) Dirac op is

$$D = \frac{1}{a} \left(\sum_{i=1}^3 \left[\sigma^i \frac{\nabla_i - \nabla_i^\dagger}{2} + \frac{1}{2} \nabla_i \nabla_i^\dagger \right] - m \right).$$

$$(\nabla_i \psi)_x = \psi_{x+\hat{i}} - \psi_x, \quad (\nabla_i^\dagger \psi)_x = \psi_{x-\hat{i}} - \psi_x$$


+OBC

We analyze $D^\dagger D$ and DD^\dagger .



Spectrum and Edge modes of $D^\dagger D$


We solve $D^\dagger D\psi = E^2\psi$ when $r_0 = 24a, ma = 0.35$.



- Localized at the boundary
- $\sigma_r\psi = +\psi$
- There is a gap from zero (as a gravitational effect)
- Agrees well with the continuum prediction

Spectrum and Edge modes DD^\dagger

We also compute $DD^\dagger\psi = E^2\psi$.




The result is almost the same as $D^\dagger D$, but edge modes are negative chiral modes:

$$\sigma_r\psi = -\psi$$

It seems that a chiral theory is possible...


Continuum Limit and Finite-volume Effect

Continuum limit $a \rightarrow 0$
($mr_0 = 8.4$ is fixed)




Agree well with
the conti. prediction!

Large volume limit $r_0 \rightarrow \infty$
($ma = 0.35$ is fixed)




Saturates in the large r_0 limit!


Restoration of Rotational Symmetry




$$a/r_0 = 1/8$$



$$a/r_0 = 1/16$$



$$a/r_0 = 1/32$$



The rotational symmetry automatically recovers in the continuum limit!

Contents

Introduction

S^2 domain-wall in \mathbb{R}^3

S^2 domain-wall on 3D square lattice

Summary

Conclusion

We investigate a **FREE** fermion system with a S^2 domain-wall.


- Weyl fermions appear at the wall.
- Continuum limit is good!
- Low-energy theory seems chiral.

But what if link variables exist?

————→ Kan will explain this situation in detail.

Outlook

- Symmetric mass generation
[Tong's talk]
- Embedding
 $4D$ Schwarzschild space into
 $6D$ flat space [Kasner, 1921].




$$\begin{aligned} ds^2 &= dx^2 + dy^2 + dz^2 + dX(r, \tau)^2 + dY(r, \tau)^2 + dZ(r, \tau)^2 \\ &= \left(1 - \frac{\beta}{4\pi r}\right) d\tau^2 + \frac{dr^2}{1 - \frac{\beta}{4\pi r}} + r^2 d\Omega^2 \end{aligned}$$

Reference i

- da Costa, R. C. T. (1981). Quantum mechanics of a constrained particle. Phys. Rev. A, 23:1982–1987.
- Jensen, H. and Koppe, H. (1971). Quantum mechanics with constraints. Annals of Physics, 63(2):586–591.
- Kaplan, D. B. (1992). A method for simulating chiral fermions on the lattice. Physics Letters B, 288(3):342–347.
- Kasner, E. (1921). Finite representation of the solar gravitational field in flat space of six dimensions. American Journal of Mathematics, 43(2):130–133.
- Nash, J. (1956). The imbedding problem for riemannian manifolds. Annals of Mathematics, 63(1):20–63.
- Onoe, J., Ito, T., Shima, H., Yoshioka, H., and ichi Kimura, S. (2012). Observation of riemannian geometric effects on electronic states. EPL (Europhysics Letters), 98(2):27001.
- Takane, Y. and Imura, K.-I. (2013). Unified description of dirac electrons on a curved surface of topological insulators. Journal of the Physical Society of Japan, 82(7):074712.


Curved Kaplan Domain-Wall [SA and H. Fukaya, 2022]



- Edge modes ($\gamma_{\text{normal}} = +1$) appear at the wall.
- They feel gravity through the induced spin connection.
- The continuum limit is unique (lattice spacing $\rightarrow 0$).
- Rotational symmetry is automatically recovered.

cf. flat case [Kaplan, 1992], spherical TI [Takane and Imura, 2013]

With nontrivial $U(1)$ link variables



When the monopole exists in the ball, another 0-mode appears around the monopole.

—→ An obstacle in formulating lattice chiral gauge theory.