



Contribution ID: 237

Type: Poster

B -meson semileptonic decay form factors from highly improved staggered quarks

Tuesday, 30 July 2024 17:15 (1 hour)

We present an update for our ongoing calculation of B -meson semileptonic form factors, using the highly improved staggered quark (HISQ) action on FNAL-MILC $N_f = 2+1+1$ HISQ gauge ensembles. We compute the scalar, vector, and tensor form factors for the $B \rightarrow \pi$, $B \rightarrow K$, $B_s \rightarrow K$, and $B_{(s)} \rightarrow D_{(s)}$ transitions. We have recently added data with $a \approx 0.03$ fm into our analysis, allowing simulation directly at the physical b -quark mass.

Primary authors: EL-KHADRA, Aida (University of Illinois Urbana-CHampaign); JAY, William (MIT)

Co-authors: Dr VAQUERO, Alejandro (University of Utah); KRONFELD, Andreas (Fermilab); LYTLE, Andrew (University of Illinois at Urbana-Champaign); DETAR, Carleton (University of Utah); GAMIZ, Elvira (University of Granada); LAIHO, Jack (Syracuse University); SIMONE, James (Fermilab); GOTTLIEB, Steven (Indiana University)

Presenters: EL-KHADRA, Aida (University of Illinois Urbana-CHampaign); JAY, William (MIT)

Session Classification: Poster session and reception

Track Classification: Quark and Lepton Flavour Physics