

## Spectroscopy of lattice gauge theories from spectral densities

*Wednesday, 3 July 2024 16:25 (25 minutes)*

We critically discuss the algorithmic process of estimating spectral densities using the Hansen-Lupo-Tantalo method. A novel approach at finite volume is deployed to extract the spectrum of lattice gauge theories. As a case study, our discussion takes as an example the study of beyond-Standard-Model (BSM) symplectic gauge theories with matter field consisting of a mixed fermion representation—fundamental and two-index antisymmetric one. We discuss potential sources of systematic effects. The results obtained with the spectral densities are critically compared with conventional data analysis techniques, vastly used in lattice QCD and BSM, such as the generalised eigenvalue problem. It will be also stressed how this algorithm and code can be applied to both QCD and BSM theories on a lattice.

**Primary author:** FORZANO, Niccolo (Swansea University)

**Co-authors:** LUPO, Alessandro (Aix-Marseille Université); Mr LUCINI, Biagio (Swansea University); Mr LIN, C.-J David (National Yang Ming Chiao Tung University); VADACCHINO, Davide (University of Plymouth); Mr HONG, Deog Ki (Pusan National University); BENNETT, Ed (Swansea University); ZIERLER, Fabian (Swansea University); HSIAO, Ho (National Yang Ming Chiao Tung University); Mr LEE, Jong-Wan (Institute for Basic Science (IBS)); DEL DEBBIO, Luigi (Edinburgh); Mr PIAI, Maurizio (Swansea University); Mr HILL, Ryan (Edinburgh University)

**Presenter:** FORZANO, Niccolo (Swansea University)

**Session Classification:** Talks