

Studies of Baryon Transition Form Factors with HADES

Tuesday, 2 July 2024 14:30 (30 minutes)

The High Acceptance Di-Electron Spectrometer (HADES) [1], installed at GSI/FAIR Helmholtzzentrum in Darmstadt, was designed for spectroscopy of positron-electron pairs in heavy-ion reactions in the SIS-18 energy range (1-2 GeV/nucleon). The main goal of this experiment is to study inclusive $e+e-$ production in pion, proton and ion induced reactions at various energies to provide information on a radiation from baryonic matter. Various models predict that this radiation proceed via intermediate rho meson. The properties of the meson are strongly modified in the cold or hot dense nuclear matter due to vector meson-baryon couplings. These couplings can be directly studied in the resonance Dalitz decays $R \rightarrow Ne+e-$. Such decays provide also information on the electromagnetic baryon-resonance transition form factors (eTFF) in the time-like region. The studies of the resonance Dalitz decays offer a great opportunity to study eTFF in a direct way. The HADES collaboration has measured the Delta(1232) Dalitz decay in p+p collisions [2] delivering, for the first time, the $\Delta \rightarrow pe+e-$ branching ratio. In the next step, using combined measurements of hadronic and dielectron final states in π -N collisions and Partial Wave Analysis (PWA) developed by the Bonn-Gatchina group [3], the contributions of $N(1440)$, $N(1520)$ and $N(1535)$ to two pion and dielectron final states have been studied. As a result cross sections for $\Delta\pi$, $N\rho$, $N\sigma$ isobar contributions have been extracted. In the dielectron channel the off-shell ρ meson contribution to the Dalitz decays of $N(1520)$ and $N(1535)$ have been obtained and allowed for extraction of the mass dependence of the effective time-like eTFF [4]. Studies of angular distributions of emitted electrons have provided an additional important information on hadronic spin density matrix elements [5]. The recent upgrade of the HADES detector [6] made possible to study also electromagnetic decays of hyperons. First measurements at HADES on both virtual and real photon decays, $Y \rightarrow Ye+e-$ and $Y^* \rightarrow Y\gamma$, have been performed in p+p collisions at 4.5 GeV [7].

The results of the HADES collaboration obtained with proton and pion beams will be presented. The eTFF will be compared to various versions of the Vector Dominance Model, to the quark-constituent model [8] and the effective Lagrangian model [5]. Prospects for HADES measurements at SIS-18 in the near future and in the further future at SIS-100 within the FAIR programme will also be discussed.

- [1] G. Agakichiev et al. (HADES), Eur. Phys. J. A 41, 243 (2009).
- [2] J. Adamczewski-Musch et al. (HADES), Phys. Rev. C 95, 06520 (2017).
- [3] J. Adamczewski-Musch et al. (HADES), Phys. Rev. C 102, 024001 (2020).
- [4] R. Abou Yassine et al. (HADES), arXiv:2205.15914 [nucl-ex]
- [5] M. Zétényi et al., Phys. Rev. C 104, 015201 (2021).
- [6] J. Adamczewski-Musch et al. (HADES), Eur. Phys. J. A 57, 138 (2021).
- [7] J. Rieger (HADES, PANDA@ HADES), EPJ Web of Conferences 291, 05005 (2024).
- [8] G. Ramalho, M. T. Pena, Phys. Rev. D 95, 014003 (2017).

Primary author: CIEPAŁ, Izabela (Institute of Nuclear Physics PAS, Kraków)

Presenter: CIEPAŁ, Izabela (Institute of Nuclear Physics PAS, Kraków)

Session Classification: Talks