



Direct dark matter searches: overview

Laura Baudis
University of Zurich
Wandering in the dark




 Search for interactions of a new particle from the Big Bang which we haven't yet detected in the lab







- Elastic (or inelastic) DM collisions
 with atomic nuclei (NRs)
- Inelastic DM collisions with electrons in the atomic shell (ERs)
- Absorption of light bosons via
 - axio-electric effect (ALPs)
 - kinetic mixing (dark photons)



Inputs from several fields required to predict expected event rates




Interaction cross section versus mass




Main experimental challenges


- To observe a signal which is:
 - very small → low recoil energies ~eV to keV (perhaps ~meV)
 - very rare \rightarrow <1 event/(kg y) at low masses and < 1 event/(t y) at high masses
 - hidden in backgrounds → deep underground & low-radioactivity materials




Backgrounds

- Cosmic rays, radioactivity
 - Deep underground, large shields
 - Ultimate: neutrinos














Backgrounds

- Cosmic rays, radioactivity
 - Deep underground, large shields
 - Ultimate: neutrinos










Neutrino signals/backgrounds

NRs (CEvNS) and ERs (all other reactions)



Experiments and signatures







See also talk by C. McCabe

Signatures

- DM particle mass
- Target material
- Time and direction





Towards the neutrino fog




LB and S. Profumo, PDG 2025


Cryogenic experiments







CRESST (Al₂O₃, CaWO₄, Si, LiAlO₂)




CRYOSEL (Ge)

Super-CDMS (Ge, Si)


- Sub-keV (< 100 eV) energy thresholds
- Phonons and/or ionisation/light ⇒background discrimination
- Optimised to probe sub-GeV DM
- All see low-energy excess (LEE)

CCDs: SENSEI and DAMIC

- Low ionisation energy, low noise, and particle tracks for background reduction \Rightarrow ~ eV energy thresholds and particle ID
- SENSEI at SNOLAB, DAMIC-M at Modane; DAMIC-M prototype: 1.3 kg-d exposure at Modane, probes freeze-in scenario
- OSCURA: next-gen detector 10 kg active, construction planned for SNOLAB



DAMIC-M, DM-electron PRL 135, 2025




Here for ultralight mediator

 $F_{DM} \sim 1/q^2$


Spherical proportional counters

- NEWS-G: light targets (H, He, Ne), pulse shape discrimination against surface events, low energy threshold (very low capacitance)
- S140 (140 cm Ø Cu sphere): 10 d physics data at Modane, then shipped to SNOLAB, taking data
- DarkSPHERE (3 m ø Cu sphere) in preparation for Boulby

Talks by G. Rogers, P. Walters



SD limits: PRL 134, 2025






S140 at Modane in its shield

Light DM: existing technologies + many new ideas

See talk by Ashlea Kemp



- Reach energy thresholds at the eV-scale (R&D towards meV-scale)
- Exposure less relevant, best constraints for kg-d or less
- Particle discrimination in general not feasible
- Low energy excess: seen by many experiments (not related to particle interactions)

Rouven Essig, Carin Cain

R. Essig/Stony Brook University; APS/Carin Cain

Noble liquids: LAr






DEAP-3600 at SNOLAB

DarkSide-20K at LNGS (Hall C)

- DEAP: detector upgrades finalised, new DM search run in 2025
- DarkSide-20k: 50 t underground LAr in TPC with 21 m² cryogenic SiPM arrays
 - PDUs: DCR < 0.01 Hz/mm2 at 77 K, PDE: ~45%, ASIC
 - Under construction: in Hall C at LNGS






DS collaboration, EPJ-C 85, 2025

Noble liquids: LAr

Low mass sensitivity

High mass sensitivity



DS collaboration, Communications Physics 7, 2024

- DarkSide-20k: first detector filling: 2027, first data taking in 2028
- Argo: 400 t (300 t fiducial) LAr in single or two-phase, proposed for SNOLAB CubeHall

Noble liquids: LXe

LZ: talks by I. Darlington, S. Dave



LUX-ZEPLIN at SURF



XENONnT at LNGS



PandaX-4T at JingPing


• Existing experiments:

- > TPCs with 2 arrays of 3-inch PMTs
- Kr and Rn removal techniques
- Ultra-pure water shields, n & μ vetos




LXe: Recent results

Distribution of events in S2 versus S1 space in the TPC




Recent results: DM-Nucleus

DM mass range: ~ 3 GeV - 10 TeV



LUX-ZEPLIN PRL 135, 2025




First Search for Light Dark Matter in the Neutrino Fog with XENONnT

Dark Matter Search Results from 4.2 Tonne – Years of Exposure of the LUX-ZEPLIN (LZ) Experiment


Recent results: DM-Electron

DM mass range: ~ 5 MeV - 1 GeV

$$F_{DM} = 1$$



$$F_{DM} \propto \frac{1}{q^2}$$



PandaX-4T, PRL 130, 2023, XENONnT, PRL 134, 2025

Heavy dark photon A' mediator

Ultra-light dark photon A' mediator


Recent results: DM Absorption

DM mass range: ~ 0.5 keV - 100 keV

Dark photons

10⁻²⁶ 10⁻²⁷ 10⁻²⁸ 10⁻²⁸ 10⁻³⁰ XENONIT (S2 only) 10⁻³¹ 10⁻³² 10⁻³³ XENONIT (S2 only) 10⁻³¹ 10⁻³² 10⁻³³ 10⁰ M_{HP} [keV/c²]


Axion like particles



XENONnT: PRL 129, 2022; LZ: PRD 108, 2023

Backgrounds

Dominated by electronic recoils from ²²²Rn decays in the LXe



Other searches with LXe detectors


- *pp and ⁷Be solar neutrinos
- *solar axions


- * $\beta\beta$ -decay ¹³⁶Xe ($Q_{\beta\beta}$ = 2.46 MeV)
- *DEC, EC β +-decay ¹²⁴Xe (Q_{DEC} = 2.86 MeV)

- *8B solar neutrinos
- *SN neutrinos

- *inelastic DM
- ***EFT DM models**

Solar neutrinos: v-Nucleus Scattering







$$\nu_x + A \rightarrow \nu_x + A$$

- *Complementary measurements to experiments using CC reactions
- * "Flavour democratic" (C. Lunardi, Neutrino-2024)

Solar neutrinos: v-Nucleus Scattering

- First detection of elastic NRs from astrophysical v's
- First measurement of CEvNS process with Xe targets
- First step into the "neutrino fog" by DM experiments








XENONnT: PRL 123, 2024

PandaX-4T: PRL 123, 2024

Very small solar ν detectors









Future LXe experiments

XLZD (XENON-LZ-DARWIN) PandaX-20T PandaX-40T





78 t LXe (60 t active target)



20 t and 40 t LXe in total; 20 t in 2027, mostly funded

PandaX-xT: SCPMN 68, 2025






XLZD, arXiv: 2410.17137

XLZD









~450 members

79 institutions

17 countries

xlzd.org


Talks by H. Araujo, R. Matheson, H. Zhang



Collaboration meeting at LNGS, July 2025

Design Book: 2410.17137 (accepted in EPJ-C)

> 13 σ stat significance, R. Bernabei et al., Nucl. Phys. At. Energy 22 (2021) 329-342




- Test DAMA/LIBRA annual modulation signal with Nal(TI)
- No evidence for annual modulation from ANAIS-112 (6 y of data, 626 kg y, ongoing) and COSINE-100 (6 y of data, Eth=0.7 keV, concluded in March 2023)
- Combined ANAIS and COSINE analysis: arXiv:2503.19559
- In construction: COSINUS, SABRE and COSINE-200


ANAIS: PRL 135, 2025, COSINE-100: 2501.13665

Combined analysis: PRL 135, 2025, 12

P. Schreiner, Santander June 2025




> 13 σ stat significance, R. Bernabei et al., Nucl. Phys. At. Energy 22 (2021) 329-342




- Test DAMA/LIBRA annual modulation signal with Nal(TI)
- No evidence for annual modulation from ANAIS-112 (6 y of data, 626 kg y, ongoing) and COSINE-100 (6 y of data, Eth=0.7 keV, concluded in March 2023)
- Combined ANAIS and COSINE analysis: arXiv:2503.19559
- In construction: COSINUS, SABRE and COSINE-200

ANAIS: PRL 135, 2025, COSINE-100: 2501.13665

Combined analysis: PRL 135, 2025, 12




> 13 σ stat significance, R. Bernabei et al., Nucl. Phys. At. Energy 22 (2021) 329-342




Combined analysis: PRL 135, 2025, 12

- SABRE at LNGS & SUPL: ultra-pure NaI powder; active shield; finish installation end of 2027 & 2025, reach 5σ in two years
- COSINE-200 at Yemilab: in-house crystal growing, improved light yields; 0.2 dru backgrounds; move to Yemilab in progress
- COSINUS at LNGS: detects also phonons in un-doped Nal (apart from scintillation) at few mK, for active background rejection; first data expected by end of 2025



SABRE South at SUPL





COSINUS at LNGS


- SABRE at LNGS & SUPL: ultra-pure NaI powder; active shield; finish installation end of 2027 & 2025, reach 5σ in two years
- COSINE-200 at Yemilab: in-house crystal growing, improved light yields; 0.2 dru backgrounds; move to Yemilab in progress
- COSINUS at LNGS: detects also phonons in un-doped Nal (apart from scintillation) at few mK, for active background rejection; first data expected by end of 2025

In addition:


Maria Martinez, TAUP2025

expression of intent to INFN-LNGS to merge know-how, experience, and equipment among ANAIS, COSINE, and SABRE collaborations for operating DAMA/LIBRA detectors:

- Direct search for modulation in same/different setup
- Independent measurement of DAMA/LIBRA NR quenching factors




SABRE South at SUPL





COSINUS at LNGS

Directional detectors




CYGNUS-UK
Boulby, UK

CYGNO
Gran Sasso, Italy

CYGNUS-OZ
Stawell, Aus.


- Gaseous TPCs to measure the recoil direction correlated to galactic motion towards Cygnus
- Challenge: good angular resolution & head
 vs. tails at low recoil energies
- Cygnus: coordinates efforts for global 1000 m^3 (He-CF₄) TPCs with ~ 1 keV threshold

F. Mayet et al., Physics Reports 627(2016)




SI cross section vs time

Constraints on SI cross section (50 GeV DM) over the past 40 years



New tools, new discoveries



see, e.g., LB., Neutrinos with dark matter detectors. Zenodo. https://doi.org/10.5281/zenodo.12745235

Summary & Outlook

- DD experiments: many different technologies, covering a large range in DM masses & cross sections
- Challenges: reduce energy thresholds further (with existing & new technologies), reduce & understand backgrounds, increase target masses
- Goals: discover a new, dark species; explore the parameter space until astrophysical neutrinos will dominate event rates
- Explore a variety of DM candidates and break new grounds in neutrino physics, solar axions, etc
- Must always be prepared for a discovery & for the unexpected ;-)


Extra material

Approaching the Neutrino Fog

- Here shown for nuclear recoils (v floor as boundary to "v fog")
- Region where experiments leave the Poissonian regime*


The "fog" for different targets

72 H 10⁻⁴¹



DM mass $[GeV/c^2]$

Effect of ν fluxes uncertainties




C. O'Hare, PRL 127, 2021

^{*} σ where the DM discovery limit scales as $\sim (Mt)^{-1/n}$


DM-electron scattering

Overview of recent results

Heavy mediator




Ultralight mediator




Dark Matter Spectroscopy

 Capability to reconstruct the DM mass and cross section for various masses - here 25, 50, 250 GeV/c² and cross sections

fixed galactic model




including galactic uncertainties



Xenon + Germanium + Argon

XLZD Dark Matter: Exclusion




XLZD collaboration
Design Book
arXiv: 2410.17137

Systematic limit imposed by CEvES from atmospheric neutrinos

Exposure of **1000 t y needed** to dive into the atmospheric neutrino fog (where > 1, 10, 100,... events are expected in the WIMP ROI)

At contour n: a 10 x lower XS sensitivity requires an increase in exposure of at least 10ⁿ

XLZD Dark Matter: Detection



Exposure of 1000 t y: 1-,2- and 3-sigma (yellow, orange, red) Cls

XLZD collaboration
Design Book
arXiv: 2410.17137


Evidence contours for 20 GeV and 80 GeV WIMPs

Theory:
EW multiplet, S. Bottaro
et al., EPJ-C 82, 2022,
Single complex EW *n*-plet
with non-zero
hypercharge added to SM


Solar v-Electron Scattering

Main challenge: reduce ²²²Rn (²¹⁴Pb β-decay) background to x 10 below the pp rate (0.1 μBq/kg)


XLZD preliminary



600 t y exposure, 1 and 4 energy bins



DARWIN collaboration, EPJ-C 80 12 (2020)




46

SN v-Nucleus Scattering v

- Sensitivity to all v flavours: few events/ton expected from SN at ~10 kpc
- Main challenge: low energies, understand few-e- backgrounds
- XLZD: sensitivity beyond SMC; part of SNEWS2.0


Rates for $27\,M_{\odot}$ SN at 10 kpc


XENONnT/LZ and XLZD




Atmospheric Neutrinos

• In general, exposures > few 1000 t y are needed for 5- σ detection





Some dark matter candidate particles



Nature Phys 13, 224-231 (2017)