

CTEQ-MCnet school on
QCD Analysis and Phenomenology
and the Physics and Techniques of Event Generators

LECTURE 3


Introduction to the Parton Model and Perturbative QCD

Fred Olness (SMU)

Lauterbad (Black Forest), Germany


26 July - 4 August 2010

Recap: Parton Model, Factorization, Evolution




DIS
AT
NLO


DIS at NLO



Sample NLO contributions to DIS



DIS NLO Kinematics




$$k_1 \equiv q^\mu = \left(\frac{s - Q^2}{2\sqrt{s}}, 0, 0, \frac{(s + Q^2)}{2\sqrt{s}} \right) \quad -q^2 = Q^2 > 0$$

$$k_2 \equiv p^\mu = \left(\frac{s + Q^2}{2\sqrt{s}}, 0, 0, \frac{-(s + Q^2)}{2\sqrt{s}} \right) \quad p^2 = 0$$

$$k_3^\mu = \frac{\sqrt{s}}{2} (1, +\sin\theta, 0, +\cos\theta) \quad k_3^2 = 0$$

$$k_4^\mu = \frac{\sqrt{s}}{2} (1, -\sin\theta, 0, -\cos\theta) \quad k_4^2 = 0$$

Mandelstam Variables {s,t,u}



$$s = (k_1 + k_2)^2 \equiv (k_3 + k_4)^2$$

$$t = (k_1 - k_3)^2 \equiv (k_2 - k_4)^2$$

$$u = (k_1 - k_4)^2 \equiv (k_2 - k_3)^2$$

$$s + t + u = m_1^2 + m_2^2 + m_3^2 + m_4^2$$

Exercise

{s,t,u} are partonic


$$s = +Q^2 \frac{(1-x)}{x} \quad t = -Q^2 \frac{(1-z)}{2x} \quad u = -Q^2 \frac{(1+z)}{2x}$$

$$x = \frac{Q^2}{2p \cdot q} \quad x \in [0, 1]$$

$$z \equiv \cos\theta \quad z \in [-1, 1]$$

Homework

1) Let's work out the general 2→2 kinematics for general masses.



a) Start with the incoming particles.


Show that these can be written in the general form:

$$p_1 = (E_1, 0, 0, +p) \quad p_1^2 = m_1^2$$

$$p_2 = (E_2, 0, 0, -p) \quad p_2^2 = m_2^2$$

... with the following definitions:

$$E_{1,2} = \frac{\hat{s} \pm m_1^2 \mp m_2^2}{2\sqrt{\hat{s}}} \quad p = \frac{\Delta(\hat{s}, m_1^2, m_2^2)}{2\sqrt{\hat{s}}}$$



$$\Delta(a, b, c) = \sqrt{a^2 + b^2 + c^2 - 2(ab + bc + ca)}$$

Note that $\Delta(a, b, c)$ is symmetric with respect to its arguments, and involves the only invariants of the initial state: s, m_1^2, m_2^2 .


b) Next, compute the general form for the final state particles, p_3 and p_4 . Do this by first aligning p_3 and p_4 along the z-axis (as p_1 and p_2 are), and then rotate about the y-axis by angle θ .

Homework Part 2

PROBLEM #2: Consider the reaction:

$pp \rightarrow pp$ ($12 \rightarrow 34$) with **CMS** scattering angle θ . The CMS energy is $\sqrt{s} = 2 \text{ TeV}$.

- Compute the boost from the CMS frame to the rest frame of #2 (lab frame)
- Compute the energy of #1 in the lab frame.
- Compute the scattering angle θ_{lab} as a function of the CMS θ and invariants.



Hint: by using invariants you can keep it simple. I.e., don't do it the way Goldstein does.

The power of invariants

$$|\mathcal{M}|^2 = \frac{s}{-t} + \frac{-t}{s} + \frac{2uQ^2}{st}$$


For the real 2→2 graphs

$$\simeq \frac{2(1-x)}{(1-z)} + \frac{2(1-z)}{(1-x)} + \frac{2x(1+z)}{(1-x)(1-z)}$$

Singular at $z=1$

$$z \rightarrow 1, \quad \cos \theta \rightarrow 1$$

$$\theta \rightarrow 0, \quad t \rightarrow 0$$




Collinear Singularity

Separate infinity, absorb in PDF

Singular at $x=1$

$$x \rightarrow 1, \quad s \rightarrow 0$$




Soft Singularity

Separate infinity, cancel with virtual graphs

The Plan

Collinear Divergences

$$|\mathcal{M}|^2 \xrightarrow{z \rightarrow 1} \frac{2}{(1-z)} \frac{(1+x^2)}{(1-x)}$$



Plan

- 1) Separate ∞ at $z=1$
- 2) "Absorb" into PDF

Looks like a PDF splitting function


Method

Need to regulate ∞

Choices

- 1) Dimensional Regularization
- 2) Quark Mass
- 3) θ Cut

Soft Singularities



Plan

- 1) Separate ∞ at $x=1$
- 2) Cancel between Real and Virtual graphs

Method

Need to regulate ∞


Choices

- 1) Dimensional Regularization
- 2) Gluon Mass
- 3) ...

Dimensional Regularization meets Freshman E&M

M. Hans, Am.J.Phys. 51 (8) August (1983). p.694
C. Kaufman, Am.J.Phys. 37 (5), May (1969) p.560
B. Delamotte, Am.J.Phys. 72 (2) February (2004) p.170

Regularization, Renormalization, and Dimensional Analysis:
Dimensional Regularization meets Freshman E&M.
Olness & Scalise, arXiv:0812.3578 [hep-ph]




$$dV = \frac{1}{4\pi\epsilon_0} \frac{dQ}{r} \quad \lambda = Q/y$$

$$V = \frac{\lambda}{4\pi\epsilon_0} \int_{-\infty}^{+\infty} dy \frac{1}{\sqrt{x^2 + y^2}} = \infty$$

*Note: ∞ can
be very useful*

Scale Invariance



$$\begin{aligned} V(kx) &= \\ &= \frac{\lambda}{4\pi\epsilon_0} \int_{-\infty}^{+\infty} dy \frac{1}{\sqrt{(kx)^2 + y^2}} \\ &= \frac{\lambda}{4\pi\epsilon_0} \int_{-\infty}^{+\infty} d\left(\frac{y}{k}\right) \frac{1}{\sqrt{x^2 + (y/k)^2}} \\ &= \frac{\lambda}{4\pi\epsilon_0} \int_{-\infty}^{+\infty} dz \frac{1}{\sqrt{x^2 + z^2}} \\ &= V(x) \end{aligned}$$

$$V(kx) = V(x)$$

Naively Implies:
 $V(kx) - V(x) = 0$

Note: $\infty + c = \infty$
 $\therefore \infty - \infty = c$

*How do we distinguish
this from*

$$\infty - \infty = c + 17$$

Cutoff Method

$$V = \frac{\lambda}{4\pi\epsilon_0} \int_{-L}^{+L} dy \frac{1}{\sqrt{x^2 + y^2}}$$

$$V = \frac{\lambda}{4\pi\epsilon_0} \log \left[\frac{+L + \sqrt{L^2 + x^2}}{-L + \sqrt{L^2 + x^2}} \right]$$

$V(x)$ depends on artificial regulator L

We cannot remove the regulator L


All physical quantities are independent of the regulator:

Electric Field $E(x) = \frac{-dV}{dx} = \frac{\lambda}{2\pi\epsilon_0} \frac{L}{x\sqrt{L^2 + x^2}} \rightarrow \frac{\lambda}{2\pi\epsilon_0} \frac{1}{x}$

Energy $\delta V = V(x_1) - V(x_2) \xrightarrow{L \rightarrow \infty} \frac{\lambda}{4\pi\epsilon_0} \log \left[\frac{x_2^2}{x_1^2} \right]$

**Problem solved at the expense of an extra scale L
AND we have a broken symmetry: translation invariance**

Broken Translational Symmetry



Shift: $y \rightarrow y' = y - c$

$$y = [+L+c, -L+c]$$

$$V = \frac{\lambda}{4\pi\epsilon_0} \int_{-L+c}^{+L+c} dy \frac{1}{\sqrt{x^2 + y^2}}$$

$$V = \frac{\lambda}{4\pi\epsilon_0} \log \left[\frac{+(L+c) + \sqrt{(L+c)^2 + x^2}}{-(L-c) + \sqrt{(L-c)^2 + x^2}} \right]$$

V(r) depends on "y" coordinate!!!

*In QFT,
gauge symmetries
are important.
E.g., Ward identities*

Dimensional Regularization

Compute in n-dimensions

$$dy \rightarrow d^n y = \frac{d\Omega_n}{2} y^{n-1} dy$$

$$\Omega_n = \int d\Omega_n = \frac{2\pi^{n/2}}{\Gamma(n/2)}$$

$$\Omega_{1,2,3,4} = \{2, 2\pi, 4\pi, 2\pi^2\}$$


$$V = \frac{\lambda}{4\pi\epsilon_0} \int_0^{+\infty} d\Omega_n \frac{y^{n-1}}{\mu^{n-1}} \frac{dy}{\sqrt{x^2 + y^2}}$$

Each term is individually dimensionless


New scale μ

$$n = 1 - 2\epsilon$$

$$V = \frac{\lambda}{4\pi\epsilon_0} \left(\frac{\mu^{2\epsilon}}{x^{2\epsilon}} \frac{\Gamma[\epsilon]}{\pi^\epsilon} \right)$$



Why do we need an extra scale μ ???



$$dV = \frac{1}{4\pi\epsilon_0} \frac{dQ}{r}$$

$$\lambda = Q/y$$

$$V = \frac{\lambda}{4\pi\epsilon_0} f(x)$$

Dimensional Regularization

All physical quantities are independent of the regulators:

$$\text{Electric Field} \quad E(x) = \frac{-dV}{dx} = \frac{\lambda}{4\pi\epsilon_0} \left[\frac{2\epsilon\mu^{2\epsilon}\Gamma[\epsilon]}{\pi^\epsilon x^{1+2\epsilon}} \right] \xrightarrow{\epsilon \rightarrow 0} \frac{\lambda}{2\pi\epsilon_0} \frac{1}{x}$$

$$\text{Energy} \quad \delta V = V(x_1) - V(x_2) \xrightarrow{\epsilon \rightarrow 0} \frac{\lambda}{4\pi\epsilon_0} \log \left[\frac{x_2^2}{x_1^2} \right]$$

Problem solved at the expense of an extra scale μ **AND** regulator ϵ

Translation invariance is preserved!!!

Dimensional Regularization respects symmetries

Renormalization

$$V \rightarrow \frac{\lambda}{4\pi\epsilon_0} \left[\frac{1}{\epsilon} + \ln \left[\frac{e^{-\gamma_E}}{\pi} \right] + \ln \left[\frac{\mu^2}{x^2} \right] \right] \quad \text{Original}$$

$$V \rightarrow \frac{\lambda}{4\pi\epsilon_0} \left[\frac{1}{\epsilon} + \ln \left[\frac{e^{-\gamma_E}}{\pi} \right] + \ln \left[\frac{\mu^2}{x^2} \right] \right] \quad \text{MS}$$

$$V \rightarrow \frac{\lambda}{4\pi\epsilon_0} \left[\frac{1}{\epsilon} + \ln \left[\frac{e^{-\gamma_E}}{\pi} \right] + \ln \left[\frac{\mu^2}{x^2} \right] \right] \quad \text{MS-Bar}$$

Physical quantities are independent of renormalization scheme!

$$V_{\overline{MS}}(x_1) - V_{\overline{MS}}(x_2) = \delta V = V_{MS}(x_1) - V_{MS}(x_2)$$

But only if performed consistently:

$$V_{\overline{MS}}(x_1) - V_{MS}(x_2) \neq \delta V \neq V_{MS}(x_1) - V_{\overline{MS}}(x_2)$$

Connection to QFT

This was the potential from our “Toy” calculation:

$$V \rightarrow \frac{\lambda}{4\pi\epsilon_0} \left[\frac{1}{\epsilon} + \ln \left[\frac{e^{-\gamma_E}}{1\pi} \right] + \ln \left[\frac{\mu^2}{x^2} \right] \right]$$



This is a partial result from
a real NLO Drell-Yan Calculation:
Cf., B. Potter

$$\frac{D(\epsilon)}{\epsilon} = \left(\frac{4\pi\mu^2}{Q^2} \right) \frac{\Gamma(1-\epsilon)}{\Gamma(1-2\epsilon)} \rightarrow \left[\frac{1}{\epsilon} + \ln \left[\frac{e^{-\gamma_E}}{4\pi} \right] + \ln \left[\frac{\mu^2}{Q^2} \right] \right]$$

Recap

Regulator provides unique definition of V, f, ω

Cutoff regulator L :
simple, but does NOT respect symmetries

Dimensional regulator ϵ :
respects symmetries: translation, Lorentz, Gauge invariance
introduces new scale μ


All physical quantities (E, dV, σ) are independent of the regulator
AND the new scale μ
Renormalization group equation: $d\sigma/d\mu=0$

We can define renormalized quantities (V, f, ω)
Renormalized (V, f, ω) are scheme dependent and arbitrary
Physical quantities (E, dV, σ) are unique and scheme independent
if we apply the scheme consistently

Apply
Dimensional
Regularization
to QFT

$$d\sigma = \frac{1}{2s} |\mathcal{M}|^2 d\Gamma$$

$$d\Gamma_i = \frac{d^D k_i}{(2\pi)^D} (2\pi) \delta(k_i^2) \quad \text{1-particle}$$



$$d\Gamma = d\Gamma_3 d\Gamma_4 (2\pi)^D \delta^D(k_1 + k_2 - k_3 - k_4) \quad \text{Final state}$$

$$d\Gamma = \frac{1}{16\pi} \left(\frac{s}{16\pi} \right)^{-\epsilon} \frac{(1 - z^2)^{-\epsilon}}{\Gamma[1 - \epsilon]} dz \quad \text{Final state}$$

$$g \rightarrow g \mu^\epsilon \quad \text{Enter, } \mu \text{ scale}$$

$$d\Gamma = \frac{1}{16\pi} \left(\frac{16\pi\mu^2}{Q^2} \right)^{+\epsilon} \frac{1}{\Gamma[1 - \epsilon]} \frac{x^\epsilon}{(1 - x)^\epsilon} (1 - z^2)^{-\epsilon} dz \quad \text{All the pieces}$$

Soft Singularities

#1) Show:

$$\frac{d^3 p}{(2\pi)^3 2E} = \frac{d^4 p}{(2\pi)^4} (2\pi) \delta^+(p^2 - m^2)$$


This relation is often useful as the RHS is manifestly Lorentz invariant

#2) Show that the 2-body phase space can be expressed as:

$$d\Gamma = \frac{d^3 p_3}{(2\pi)^3 2E_3} \frac{d^3 p_4}{(2\pi)^3 2E_4} (2\pi)^4 \delta^4(p_1 + p_2 - p_3 - p_4) = \frac{d\cos(\theta)}{16\pi}$$

Note, we are working with massless partons, and θ is in the partonic CMS frame

Soft Singularities




$$\underbrace{\frac{x^\epsilon}{(1-x)^\epsilon}}_{\text{From phase space}} \underbrace{\frac{1}{(1-x)}}_{\text{Soft Singularity}} = \underbrace{\frac{1}{(1-x)_+}}_{\text{Finite remainder}} - \underbrace{\frac{1}{\epsilon} \delta(1-x)}_{\text{To be canceled by virtual diagram}}$$

This only makes sense under the integral

$$\frac{f(x)}{(1-x)_+} = \frac{f(x) - f(1)}{(1-x)}$$

$$\int_0^1 dx f(x) \frac{x^\epsilon}{(1-x)^{1+\epsilon}} = \int_0^1 dx \frac{f(x) - f(1)}{(1-x)} - \frac{1}{\epsilon} \int_0^1 dx \delta(1-x) f(x)$$

Collinear Singularities



Collinear Singularity

$$\int_{-1}^1 (1 - z^2)^{-\epsilon} |\mathcal{M}|^2 \simeq -\frac{1}{\epsilon} \frac{(1 + x^2)}{(1 - x)} + \frac{1 - 4x + 4(1 + x^2) \ln 2}{2(1 - x)}$$

This should be “absorbed” in the PDF
This is finite for $z \in [-1, 1]$

... looks like a splitting kernel

Key Points

- 1) “Absorb $1/\epsilon$ into PDF
- 2) This defines how to regularize PDF
- 3) Need to match schemes of ω and PDF
... MS , $MS\text{-}Bar$, DIS , ...
- 4) Note we have regulator ϵ and extra scale μ

How do we know what to
“absorb” into PDFs ???

Compute NLO Subtractions
for a partonic target

Application of Factorization Formula at Leading Order (LO)

Basic Factorization Formula

$$\sigma = f \otimes \omega + \mathcal{O}(\Lambda^2/Q^2)$$

At Zeroth Order:

$$\sigma^0 = f^0 \otimes \omega^0 + \mathcal{O}(\Lambda^2/Q^2)$$


Use: $f^0 = \delta$ for a **parton** target.

Therefore:

$$\sigma^0 = f^0 \otimes \omega^0 = \delta \otimes \omega^0 = \omega^0$$

$$\sigma^0 = \omega^0$$

Higher Twist



Warning: This trivial result leads to many misconceptions at higher orders

Application of Factorization Formula at NLO

Basic Factorization Formula

$$\sigma = f \otimes \omega + \mathcal{O}(\Lambda^2/Q^2)$$

At First Order:

$$\sigma^1 = f^1 \otimes \omega^0 + f^0 \otimes \omega^1$$



$$\sigma^1 = f^1 \otimes \sigma^0 + \omega^1$$

We used: $f^0 = \delta$ for a **parton** target.

Therefore:

$$\omega^1 = \sigma^1 - f^1 \otimes \sigma^0$$

$$\omega^1 =$$



$$f^1 \sim \frac{\alpha_s}{2\pi} P^{(1)}$$

$P^{(1)}$ defined by
scheme choice

Application of Factorization Formula at NLO

Combined Result:

$$\underbrace{\omega^0 + \omega^1}_{\text{TOT}} = \underbrace{\sigma^0}_{\text{LO}} + \underbrace{\sigma^1}_{\text{NLO}} - \underbrace{f^1 \otimes \sigma^0}_{\text{SUB Subtraction}}$$


TOT

LO

NLO

SUB
Subtraction

$$\text{TOT} = \text{LO} + \text{NLO} - \text{SUB}$$



HOMEWORK PROBLEM: NNLO WILSON COEFFICIENTS

Use the Basic Factorization Formula

$$\sigma = f \otimes \omega \otimes d + \mathcal{O}(\Lambda^2/Q^2)$$

At Second Order (NNLO):

$$\sigma^2 = f^2 \otimes \omega^0 \otimes d^0 + \dots + f^1 \otimes \omega^1 \otimes d^0 + \dots$$

Therefore:


$$\omega^2 = ???$$

Compute ω^2 at second order.
Make a diagrammatic representation of each term.

Include Fragmentation
Functions d

Do we get different answers if we
“absorb” different terms
into PDFs ???

Pictorial Demonstration of Scheme Consistency




Parton Model


$$\sigma(Q^2) = f(\mu, \alpha_s) \otimes \hat{\omega}(Q^2, \mu^2, \alpha_s)$$

Evolution Equation

$$\frac{df}{\ln[\mu^2]} = P \otimes f$$



Pictorial Demonstration of Scheme Consistency



$$[\delta + f^1] \otimes [\sigma_0 + \underbrace{\sigma_1 - f^1 \otimes \sigma_0}_{\text{Complete NLO Term}}]$$

$$f^1 \sim \frac{\alpha_s}{2\pi} P^{(1)}$$

$P^{(1)}$ defined by
scheme choice

$$\sigma_0 + \sigma_1 + f^1 \otimes \sigma_0 - f^1 \otimes \sigma_0 + \mathcal{O}(\alpha_s^2)$$

Contains BOTH collinear and non-collinear region

From PDF Evolution

From NLO Subtraction

QCD is
Bullet-proof

Do we get different answers if we
“absorb” different terms
into PDFs ???

NO !!!

- NLO Theoretical Calculations:
 - Essential for accurate comparison with experiments
- We encounter singularities:
 - Soft singularities: cancel between real and virtual diagrams
 - Collinear singularities: “absorb” into PDF
- Regularization and Renormalization:
 - Regularize & Renormalize intermediate quantities
 - Physical results independent of regulators (e.g., L , or μ and ϵ)
 - Renormalization introduces scheme dependence (MS-bar, DIS)
- Factorization works:
 - Hard cross section $\hat{\sigma}$ or ω is not the same as σ
 - Scheme dependence cancels out (if performed consistently)

END OF LECTURE 3