

New results for algebraic NLO

# Tensor reduction of Feynman integrals

Tord Riemann

DESY, Zeuthen, Germany

Based on work done in collaboration with Jochem Fleischer and Valery Yundin  
Workshop on QCD at LHC, Aug 21 – Aug 25, 2011, St. Andrews, Scotland

## A simple example

### 1-loop self-energy:

$$\begin{aligned} I_2^\mu &= \int \frac{d^d k}{i\pi^{d/2}} \frac{k^\mu}{[k^2 - M_1^2][(k+p)^2 - M_2^2]} \\ &= p_\mu B_1 \end{aligned}$$

### Solve:

$$p_\mu I_2^\mu = p^2 B_1(p, M_1, M_2)$$

$$\begin{aligned} &= \int \frac{d^d k}{i\pi^{d/2}} \frac{pk}{[k^2 - M_1^2][(k+p)^2 - M_2^2]} = \int \frac{d^d k}{i\pi^{d/2}} \frac{pk}{D_1 D_2} \\ &= \int \frac{d^d k}{i\pi^{d/2}} \left[ \frac{D_2 - (p^2 - M_2^2 - M_1^2) - D_1}{D_1 D_2} \right], \end{aligned}$$

$$B_1(p, M_1, M_2) = \frac{1}{2p^2} \left[ A_0(M_1) - A_0(M_2) - (p^2 - M_2^2 - M_1^2) B_0(p, M_1, M_2) \right]$$

A **tensor** Feynman integral is expressed in terms of **scalar** Feynman integrals.

Systematic approach to tensor reductions:

1- to 4-point functions: Passarino, Veltman 1978 [1]

Need in addition a library of scalar functions:

'tHooft, Veltman 1979 [2]

State of the art + open source programs:

K. Ellis and G. Zanderighi, QCDloop/FF [25, 4] 2007, 1990

T. Hahn, LoopTools/FF [3, 4] 1998, 1990

To our knowledge, there is no open source program for 5-point reductions without certain restrictions (?)

→ C++ code PJFry by V. Yundin, released this Summer 2011

This talk: Efficient reduction formulae in the algebraic  
Fleischer-Davydychev-Tarasov approach

Recent developments in the Fleischer-Davydychev-Tarasov approach:

- Get tensor reduction with  $\dots$ :
- $\dots$  arbitrary masses
- $\dots$  killed pentagon Gram determinants
- $\dots$  treatment of full kinematics, also with small sub-diagram Gram determinants
  - presented by J. Fleischer at QCD@LHC@Trento2010
- → c++ code PJFry by V. Yundin
- $\dots$  multiple sums made efficient by contracting with external momenta

# Outline

- [5] 1991 Davydychev, . . . Reducing Feynman diagrams to scalar integrals
- [6] 1996 Tarasov, Connection [of] Feynman integrals [with] different . . . space-time dimensions
- [7] 1999 Fleischer et al., Algebraic reduction of one-loop Feynman graph amplitudes

- 1 Introduction
- 2 Recursions
- 3 Simplifying
- 4 Numbers:  $D_{1111}$
- 5 Small Grams
- 6 PJFry
- 7 External momenta
- 8 Summary

## References:

- [9] 2010 Diakonidis et al., PLB 683, . . . recursive reduction of tensor Feynman integrals
  - [15] 2011 Fleischer, T.R., PRD 83, Complete . . . reduction of . . . tensor Feynman integrals
  - [19] 2011 Fleischer, T.R., PLB 701, . . . contracted tensor Feynman integrals
- subm. Aug. 2011: V. Yundin, PhD thesis [with PJFry code]

## Notations: Gram and modified Cayley determinant, signed minors [Melrose:1965]

Gram determinant  $G_n$ :

$$G_n = |2q_i q_j|, i, j = 1, \dots, n \quad (1)$$

Modified Cayley determinant  $()_N$  of a diagram with  $N$  internal lines and chords  $q_j$ ; for a choice  $q_n = 0$ , both determinants are related:

$$()_N \equiv \begin{vmatrix} 0 & 1 & 1 & \dots & 1 \\ 1 & Y_{11} & Y_{12} & \dots & Y_{1N} \\ 1 & Y_{12} & Y_{22} & \dots & Y_{2N} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & Y_{1N} & Y_{2N} & \dots & Y_{NN} \end{vmatrix} = -G_{N-1}, \quad (2)$$

where  $D_i = (k - q_i)^2 - m_i^2$  [with  $q_i$  = chord], and the matrix elements

$$Y_{ij} = -(q_i - q_j)^2 + m_i^2 + m_j^2, \quad (i, j = 1 \dots N) \quad (3)$$

⇒ The Gram determinant  $()_N$  does not depend on the masses.

## Notations: signed minors [Melrose:1965]

signed minors of  $(\cdot)_N$  are constructed by deleting  $m$  rows and  $m$  columns from  $(\cdot)_N$ , and multiplying with a sign factor:



$$\begin{aligned} \left( \begin{array}{cccc} j_1 & j_2 & \cdots & j_m \\ k_1 & k_2 & \cdots & k_m \end{array} \right)_N &\equiv \\ &\equiv (-1)^{\sum_i (j_i + k_i)} \operatorname{sgn}_{\{j\}} \operatorname{sgn}_{\{k\}} \left| \begin{array}{c|c} \text{rows } j_1 \cdots j_m \text{ deleted} \\ \hline \text{columns } k_1 \cdots k_m \text{ deleted} \end{array} \right| \end{aligned} \quad (4)$$

where  $\operatorname{sgn}_{\{j\}}$  and  $\operatorname{sgn}_{\{k\}}$  are the signs of permutations that sort the deleted rows  $j_1 \cdots j_m$  and columns  $k_1 \cdots k_m$  into ascending order.

Example:

$$\left( \begin{array}{cc} 0 & 0 \end{array} \right)_N \equiv \left| \begin{array}{cccc} Y_{11} & Y_{12} & \cdots & Y_{1N} \\ Y_{12} & Y_{22} & \cdots & Y_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ Y_{1N} & Y_{2N} & \cdots & Y_{NN} \end{array} \right|, \quad (5)$$

## Example: Getting a 4-point function from a six-point function I



**Figure:** A six-point topology (a) leading to four-point functions (b) with realistically vanishing Gram determinants.

## Example: Getting a 4-point function from a six-point function II

The example is taken from [10].

The corresponding 4-point tensor integrals are, in LoopTools [3, 11] notation:

$$\text{D0i(id, 0, 0, } s_{\bar{\nu}u}, t_{ed}, t_{\bar{e}\mu}, s_{\mu\bar{\nu}u}, 0, M_Z^2, 0, 0). \quad (6)$$

The Gram determinant is:

$$()_4 = -2t_{\bar{e}\mu}[s_{\mu\bar{\nu}u}^2 + s_{\bar{\nu}u}t_{ed} - s_{\mu\bar{\nu}u}(s_{\bar{\nu}u} + t_{ed} - t_{\bar{e}\mu})], \quad (7)$$

It vanishes if:

$$t_{ed} \rightarrow t_{ed,\text{crit}} = \frac{s_{\mu\bar{\nu}u}(s_{\mu\bar{\nu}u} - s_{\bar{\nu}u} + t_{\bar{e}\mu})}{s_{\mu\bar{\nu}u} - s_{\bar{\nu}u}}. \quad (8)$$

In terms of a dimensionless scaling parameter  $x$ ,

$$t_{ed} = (1 + x)t_{ed,\text{crit}}, \quad (9)$$

## Example: Getting a 4-point function from a six-point function III

the Gram determinant becomes:

$$()_4 = 2 \times s_{\mu\bar{\nu}u} t_{\bar{e}\mu} (s_{\mu\bar{\nu}u} - s_{\bar{\nu}u} + t_{\bar{e}\mu}). \quad (10)$$

We will also need the modified Cayley determinant:

$$\begin{aligned} \begin{pmatrix} 0 \\ 0 \end{pmatrix}_4 &= \begin{pmatrix} 2M_Z^2 & M_Z^2 & M_Z^2 - s_{\mu\bar{\nu}u} & M_Z^2 \\ M_Z^2 & 0 & -s_{\bar{\nu}u} & M_Z^2 \\ M_Z^2 - s_{\mu\bar{\nu}u} & -s_{\bar{\nu}u} & 0 & -t_{ed} \\ M_Z^2 & -t_{\bar{e}\mu} & -t_{ed} & 0 \end{pmatrix} \\ &= s_{\mu\bar{\nu}u}^2 t_{\bar{e}\mu}^2 + 2 M_Z^2 t_{\bar{e}\mu} [-2s_{\bar{\nu}u} t_{ed} + s_{\mu\bar{\nu}u} (s_{\bar{\nu}u} + t_{ed} - t_{\bar{e}\mu})] \\ &\quad + M_Z^4 (s_{\bar{\nu}u}^2 + (t_{ed} - t_{\bar{e}\mu})^2 - 2s_{\bar{\nu}u} (t_{ed} + t_{\bar{e}\mu})). \end{aligned}$$

## Dimensional shifts and recurrence relations for pentagons (I)

Following [Davydychev:1991 [5]]

Replace tensors by scalar integrals in higher dimensions:

Example  $R = 3$ :

$$\begin{aligned} I_5^{\mu\nu\lambda} &= \int \frac{d^{4-2\epsilon} k}{i\pi^{d/2}} \prod_{r=1}^5 c_r^{-1} k^\mu k^\nu k^\lambda \\ &= - \sum_{i,j,k=1}^4 q_i^\mu q_j^\nu q_k^\lambda n_{ijk} I_{5,ijk}^{[d+]^3} + \frac{1}{2} \sum_{i=1}^{n-1} (g^{\mu\nu} q_i^\lambda + g^{\mu\lambda} q_i^\nu + g^{\nu\lambda} q_i^\mu) I_{5,i}^{[d+]^2}, \end{aligned} \tag{11}$$

and  $n_{ijk} = (1 + \delta_{ij})(1 + \delta_{ik} + \delta_{jk})$ .

$$[d+]^l = 4 - 2\epsilon + 2l$$

$I_{5,i}^{[d+]^2}$  – scratch the line  $i$  from  $I_5^{[d+]^2}$ .

## Dimensional shifts and recurrence relations for pentagons (II)

'Naive', direct approach – just perform dimensional recurrences

Following [Tarasov:1996,Fleischer:1999 [6, 7]]

apply **recurrence relations**, relating scalar integrals of different dimensions, in order to get rid of the dimensionalities  $[d+]^l = 4 - 2\epsilon + 2l$ :

$$\nu_j(\mathbf{j}^+ I_5^{[d+]}) = \frac{1}{\binom{d}{0}_5} \left[ -\binom{j}{0}_5 + \sum_{k=1}^5 \binom{j}{k}_5 \mathbf{k}^- \right] I_5 \quad (12)$$

$$(d - \sum_{i=1}^5 \nu_i + 1) I_5^{[d+]} = \frac{1}{\binom{d}{0}_5} \left[ \binom{0}{0}_5 - \sum_{k=1}^5 \binom{0}{k}_5 \mathbf{k}^- \right] I_5, \quad (13)$$

where the operators  $\mathbf{i}^\pm, \mathbf{j}^\pm, \mathbf{k}^\pm$  act by shifting the indices  $\nu_i, \nu_j, \nu_k$  by  $\pm 1$ .

## The result of simplifying manipulations

... and collecting all contributions, our final result for e.g. the tensor of rank  $R = 3$  can be written as follows:

$$I_5^{\mu \nu \lambda} = \sum_{i,j,k=1}^4 q_i^\mu q_j^\nu q_k^\lambda E_{ijk} + \sum_{k=1}^4 g^{[\mu\nu} q_k^{\lambda]} E_{00k}, \quad (14)$$

with:

$$E_{00j} = \sum_{s=1}^5 \frac{1}{\binom{0}{0}_5} \left[ \frac{1}{2} \binom{0s}{0j}_5 I_4^{[d+],s} - \frac{d-1}{3} \binom{s}{j}_5 I_4^{[d+]^2,s} \right], \quad (15)$$

$$E_{ijk} = - \sum_{s=1}^5 \frac{1}{\binom{0}{0}_5} \left\{ \left[ \binom{0j}{sk}_5 I_4^{[d+]^2,s}_{4,i} + (i \leftrightarrow j) \right] + \binom{0s}{0k}_5 \nu_{ij} I_4^{[d+]^2,s}_{4,ij} \right\}. \quad (16)$$

✓ no scalar 5-point integrals in higher dimensions

✓ no inverse Gram det.  $({})_5$

We have yet:

† scalar 4-point integrals in higher dimensions:  $I_4^{[d+]^2,s}_{4,ij}$  etc.

† inverse Gram det.  $\binom{0}{0}_5 \equiv ({})_4$

Reduce  $I_{4,ij\dots}^{[d+],s}$  to  $I_4^{[d+],s}$  plus simpler objects !

By nontrivial manipulations we get e.g.:

$$I_{4,i}^{[d+],s} = \frac{1}{\binom{0s}{0s}_5} \left[ -\binom{0s}{is}_5 (d-3) I_4^{[d+],s} + \sum_{t=1}^5 \binom{0st}{0si}_5 I_3^{st} \right] \quad (17)$$

$$\nu_{ij} I_{4,ij}^{[d+]^2} = \frac{\binom{0}{i}_4 \binom{0}{j}_4}{\binom{0}{0}_4 \binom{0}{0}_4} (d-2)(d-1) I_4^{[d+]^2} + \frac{\binom{0i}{0j}_4}{\binom{0}{0}_4} I_4^{[d+]} \\ - \frac{\binom{0}{j}_4}{\binom{0}{0}_4} \frac{d-2}{\binom{0}{0}_4} \sum_{t=1}^4 \binom{0t}{0i}_4 I_3^{[d+],t} + \frac{1}{\binom{0}{0}_4} \sum_{t=1}^4 \binom{0t}{0j}_4 I_3^{[d+],t} \quad (18)$$

These equations are free of inverse Gram determinants  $(\cdot)_4$ .  
 But they contain yet the generic 4-point and (partly indexed) 3-point functions in higher dimensions,  $I_4^{[d+],s}$ ,  $I_3^{[d+],t}$ , etc.

Last step: evaluate the  $I_4^{[d+],s}$ ,  $I_3^{[d+],t}$ , etc. I

Several strategies are now possible:

- Just evaluate them **analytically** in  $d + 2l - 2\epsilon$  dimensions – if you may do that
- Just evaluate them **numerically** in  $d + 2l - 2\epsilon$  dimensions
- **Reduce** them further by recurrences – buy the towers of  $1/()_4 \rightarrow$  apply (13)
- Make a **small Gram determinant expansion**  $\rightarrow$  apply (13) another way round

Last two items are done here.

Reduction of scalars  $I_4^D$  to the generic dimension  $\rightarrow I_4^d = D_0$ ,  $I_3^d = C_0$  !

Non-small 4-point Gram determinants:

Direct, iterative use of (13) yields e.g.:

$$I_4^{[d+]^l} = \left[ \frac{\binom{0}{0}_4}{\binom{l}{4}} I_4^{[d+]^{l-1}} - \sum_{t=1}^4 \frac{\binom{t}{0}_4}{\binom{l}{4}} I_3^{[d+]^{l-1}, t} \right] \frac{1}{d+2l-5} \quad (19)$$

$$I_3^{[d+]^l, t} = \left[ \frac{\binom{0t}{0t}_4}{\binom{t}{t}_4} I_3^{[d+]^{l-1}, t} - \sum_{u=1, u \neq t}^4 \frac{\binom{ut}{0t}_4}{\binom{t}{t}_4} I_2^{[d+]^{l-1}, tu} \right] \frac{1}{d+2l-4} \quad (20)$$

And we are done.

This works fine if  $\binom{0}{4}$  is not small [and also the  $\binom{t}{t}_4$ ].

# Make a small Gram expansion I

Again use (13):

$$(0)_4(d - \sum_{i=1}^4 \nu_i + 1) I_4^{[d+]} = \left[ \binom{0}{0}_4 I_4 - \sum_{k=1}^4 \binom{0}{k}_4 I_3^k \right]$$

If  $(0)_4 = 0$ , then it follows ( $n = 4$ ):

$$I_n^D = \sum_k^n \frac{\binom{0}{k}_n}{\binom{0}{0}_n} I_{n-1}^{D,k} \quad (21)$$

If  $(0)_4 \ll 1$ , re-write (13), as follows:

$$I_n^D = \sum_k^n \frac{\binom{0}{k}_n}{\binom{0}{0}_n} I_{n-1}^{D,k} - \frac{(0)_n}{\binom{0}{0}_n} [(D+1) - \sum_i^n \nu_i] I_n^{D+2}. \quad (22)$$

Effectively we may evaluate  $I_n^D$  in terms of simpler functions  $I_{n-1}^{D,k}$  with a small correction depending on  $I_n^{D+2}$ .

We may go a step further, and insert into (22) for  $I_n^{D+2}$  the rhs. of (21), taken now at  $D' = D + 2$ :

$$\begin{aligned} I_n^D &= \sum_k^n \frac{\binom{0}{k}_n}{\binom{0}{0}_n} I_{n-1}^{D,k} \\ &\quad - \frac{(\_)_n}{\binom{0}{0}_n} [(D+1) - \sum_i^n \nu_i] \\ &\quad \times \left[ \sum_k^n \frac{\binom{0}{k}_n}{\binom{0}{0}_n} I_{n-1}^{D+2,k} - \frac{(\_)_n}{\binom{0}{0}_n} [(D+3) - \sum_i^n \nu_i] I_n^{D+4} \right]. \end{aligned}$$

The terms proportional to  $[(\_)_n / \binom{0}{0}_n]^a$ ,  $a = 0, 1$  may be evaluated at the correct kinematics. They depend on three-point functions, and their reduction by normal recurrences will not introduce the unwanted powers of  $1/(\_)_4$ . The last term, suppressed by the factor  $[(\_)_n / \binom{0}{0}_n]^2$ , depends on  $I_n^{D+4}$ . It may either be taken approximately at  $(\_)_n = 0$ , where it can also be represented by 3-point functions (and their reductions), or it may be evaluated more correctly by another iteration based on (21).

And so on and so on ...

In the numerical example – next section – we worked out up to 10 stable iterations.

A quite similar attempt to perform such a series of approximations was undertaken in [14] (see equation (5) there), where a specific **example, forward light-by-light scattering through a massless fermion loop**, was studied. The approach was then not further followed.

W. Giele, E. W. N. Glover, and G. Zanderighi,  
in: Proceedings of Loops ans Legs 2004:  
*Numerical evaluation of one-loop diagrams near exceptional momentum configurations,*

Following Davydychev, [5], one gets

$$I_4^{\mu\nu\lambda} = \int^d \frac{k^\mu k^\nu k^\lambda}{\prod_{r=1}^n c_r} = - \sum_{i,j,k=1}^n q_i^\mu q_j^\nu q_k^\lambda \nu_{ijk} I_{n,ijk}^{[d+]^3} + \frac{1}{2} \sum_{i=1}^n g^{[\mu\nu} q_i^{\lambda]} I_{n,i}^{[d+]^2} \quad (23)$$

We identify the tensor coefficients  $D_{11\dots}$  a la LoopTools,  
e.g.:

$$D_{111} = I_{4,222}^{[d+]^3} \quad (24)$$

Similarly:

$$D_{1111} = I_{4,2222}^{[d+]^4} \quad (25)$$

## Rank $R = 4$ tensor $D_{1111}$ – Numerics with dimensional recurrences

From (22) we see that a “small Gram determinant” expansion will be useful when the following dimensionless parameter becomes small:



$$R = \frac{()_4}{\binom{0}{0}_4} \times s, \quad (26)$$

where  $s$  is a typical scale of the process, e.g. we will choose  $s = s_{\mu\bar{\nu}u}$ .  
Following [10], we further choose:

$$\begin{aligned} s_{\mu\bar{\nu}u} &= 2 \times 10^4 \text{ GeV}^2, \\ s_{\bar{\nu}u} &= 1 \times 10^4 \text{ GeV}^2, \\ t_{\bar{e}\mu} &= -4 \times 10^4 \text{ GeV}^2, \end{aligned}$$

and get  $t_{ed,\text{crit}} = -6 \times 10^4 \text{ GeV}^2$ . For  $x=1$ , the Gram determinant becomes  $()_4 = 4.8 \times 10^{13} \text{ GeV}^3$ .

The small expansion parameter  $R(x)$  and  $D_{1111}$  are shown in figure 2.



## New for QCD at LHC: small Gram expansion and Pade approximation I

[Fleischer,TR: PRD 2011 [15] ]

Tables have been taken from there.

They were shown already at [QCD@LHC@Trento2010](#)

The use of appropriate Pade approximations is explained there.  
Convergence in the small Gram determinant region is considerably improved.

$x$	$\Re e D_{1111}$	$\Im m D_{1111}$
0. [exp 0,0]	2.05969289730 E-10	1.55594910118 E-10
$10^{-8}$ [exp x,2] [exp 0,2]	2.05969289342 E-10	1.55594909187 E-10
	2.05969289349 E-10	1.55594909187 E-10
$10^{-4}$ [exp x,5] [exp 0,5]	2.05965609497 E-10	1.55585605343 E-10
	2.05965609495 E-10	1.55585605343 E-10
0.001 [exp 0,6] [exp x,6]	2.05932484380 E-10	1.55501912433 E-10
	2.05932484381 E-10	1.55501912433 E-10
$I_{4,2222}^{[d+4]}$ $D_{1111}$	2.02292295240 E-10	1.54974785467 E-10
	2.01707671668 E-10	1.62587142251 E-10
0.005 [exp 0,6] [pade 0,3]	2.05786054801 E-10	1.55131031024 E-10
	2.05785198947 E-10	1.55131031003 E-10
[exp x,6] [pade x,3]	2.05786364440 E-10	1.55131031024 E-10
	2.05785199805 E-10	1.55131030706 E-10
$I_{4,2222}^{[d+4]}$ $D_{1111}$	2.05778894114 E-10	1.55135794453 E-10
	2.05779811490 E-10	1.55136343923 E-10
0.01 [exp 0,6] [pade 0,3]	2.05703298143 E-10	1.54669910676 E-10
	2.05600940065 E-10	1.54669907784 E-10
[exp 0,10] [pade 0,5]	2.05600964693 E-10	1.54669910676 E-10
	2.05600955381 E-10	1.54669910676 E-10
[exp x,10] [pade x,5]	2.05600963675 E-10	1.54669910676 E-10
	2.05600955381 E-10	1.54669910676 E-10
$I_{4,2222}^{[d+4]}$ $D_{1111}$	2.05600013702 E-10	1.54670651917 E-10
	2.05600239280 E-10	1.54670771210 E-10

**Table:** Numerical values for the tensor coefficient  $D_{1111}$ . Values marked by  $D_{1111}$  are evaluated with LoopTools, the  $I_{4,2222}^{[d+4]}$  corresponds to (28) The labels [exp 0,2n] and [pade 0,n] denote iteration 2n and Pade approximant [n, n] when the small Gram determinant expansion starts at  $x = 0$ , and [exp x,2n] and [pade x,n] are the corresponding numbers for an expansion starting at  $x$ .

$x$	$\Re D_{1111}$	$\Im D_{1111}$
0.01 [exp 0,6] [pade 0,3] [exp 0,10] [pade 0,5] [exp x,10] [pade x,5]	2.05703298143 E-10 2.05600940065 E-10 2.05600964693 E-10 2.05600955381 E-10 2.05600963675 E-10 2.05600955381 E-10	1.54669910676 E-10 1.54669907784 E-10 1.54669910676 E-10 1.54669910676 E-10 1.54669910676 E-10 1.54669910676 E-10
$\text{I}_{4,2222}^{[d+]} D_{1111}$	2.05600013702 E-10 2.05600239280 E-10	1.54670651917 E-10 1.54670771210 E-10
0.05 [exp 0,6] [pade 0,3] [exp 0,20] [pade 0,10] [exp x,20] [pade x,10]	4.83822963052 E-09 2.01518061131 E-10 2.04218962072 E-10 2.04122727654 E-10 2.04190274030 E-10 2.04122727971 E-10	1.51077429118 E-10 1.50591643209 E-10 1.51077424143 E-10 1.51077424149 E-10 1.51077424143 E-10 1.51077423985 E-10
$\text{I}_{4,2222}^{[d+]} D_{1111}$	2.04122726387 E-10 2.04122726601 E-10	1.51077422901 E-10 1.51077423320 E-10
0.1 [exp 0,26] [pade 0,13] [exp x,26] [pade x,13]	2.20215264409 E-08 2.01749674352 E-10 2.08190721550 E-08 2.03995221326 E-10	1.46815247004 E-10 1.46681287362 E-10 1.46815247004 E-10 1.46785977364 E-10
$\text{I}_{4,2222}^{[d+]} D_{1111}$	2.02269485177 E-10 2.02269485217 E-10	1.46815247061 E-10 1.46815247051 E-10
1. $\text{I}_{4,2222}^{[d-]} D_{1111}$	1.72115440143 E-10 1.72115440148 E-10	9.74550747662 E-11 9.74550747662 E-11

**Table:** Numerical values for the tensor coefficient  $D_{1111}$ . Values marked by  $D_{1111}$  are evaluated with

LoopTools, the  $\text{I}_{4,2222}^{[d+]}$  corresponds to (28) The labels [exp 0,2n] and [pade 0,n] denote iteration 2n and Pade approximant [n, n] when the small Gram determinant expansion starts at  $x = 0$ , and [exp x,2n] and [pade x,n] are the corresponding numbers for an expansion starting at  $x$ .

# PJFry - an open source c++ program by V. Yundin I

## PJFry 1.0.0 - one loop tensor integral library

- More information and the latest source code:  
project page: <https://github.com/Vayu/PJFry/>
- → how to install
- → how to use
- → samples
- See also:  
V. Yundin's **talk** at LHCphenoNet meeting, Valencia, Feb 2011:  
“One loop tensor reduction program PJFRY”

# PJFry - an open source c++ program by V. Yundin II

- Yundin's PhD thesis, submitted Aug 2011 at Humboldt University

## PJFry — numerical package [from V.Y. Valencia 2011]


Numerical implementation of described algorithms:

C++ package **PJFry** by **V. Yundin** [see project webpage]

- Reduction of **5-point** 1-loop tensor integrals up to **rank 5**
- No limitations on internal/external masses combinations
- Small Gram determinants treatment by expansion
- Interfaces for C, C++, FORTRAN and MATHEMATICA

### Example:


Relative accuracy of  $E_{3333}$  coef. around small Gram4 region



## PJFry — small Gram region example [from V.Y. Valencia 2011] |

**Example:**  $E_{3333}$  coefficient in small Gram region ( $x = 0$ )

Comparison of Regular and Expansion formulae:



$$x=0: E_{3333}(0, 0, -6 \times 10^4, 0, 0, 10^4, -3.5 \times 10^4, 2 \times 10^4, -4 \times 10^4, 1.5 \times 10^4, 0, 6550, 0, 0, 8315)$$

## Contractions with external momenta [or with CHORDS] I

[Fleischer,TR: PLB 2011 [19]]

After having tensor reductions with basis functions  $I_n^D$ , which are independent of the indices  $i, j, k, \dots$ , one may use **contractions with external momenta** in order to perform all the sums over  $i, j, k, \dots$ .

This leads to a **significant simplification and shortening** of calculations.

**Reminder:**

One option was to avoid the appearance of inverse Gram determinants  $1/(\cdot)_5$ . For rank  $R = 5$ , e.g.,

$$\begin{aligned} I_5^{\mu\nu\lambda\rho\sigma} &= \sum_{s=1}^5 \left[ \sum_{i,j,k,l,m=1}^5 q_i^\mu q_j^\nu q_k^\lambda q_l^\rho q_m^\sigma E_{ijklm}^s + \sum_{i,j,k=1}^5 g^{[\mu\nu} q_i^\lambda q_j^\rho q_k^\sigma] E_{00ijk}^s \right. \\ &\quad \left. + \sum_{i=1}^5 g^{[\mu\nu} g^{\lambda\rho} q_i^\sigma] E_{0000i}^s \right] \end{aligned} \tag{27}$$

## Contractions with external momenta [or with CHORDS] I

The tensor coefficients are expressed in terms of integrals  $I_{4,i\dots}^{[d+]^l,s}$ , e.g.:

$$\begin{aligned} E_{ijklm}^s &= -\frac{1}{\binom{0}{0}_5} \left\{ \left[ \binom{0l}{sm}_5 n_{ijk} I_{4,ijk}^{[d+]^4,s} + (i \leftrightarrow l) + (j \leftrightarrow l) + (k \leftrightarrow l) \right] \right. \\ &\quad \left. + \binom{0s}{0m}_5 n_{jkl} I_{4,ijkl}^{[d+]^4,s} \right\}. \end{aligned}$$

Now, in a next step, one may avoid the appearance of inverse sub-Gram determinants  $(\cdot)_4$ .

The complete dependence on the indices  $i$  of the tensor coefficients is contained now in the pre-factors with signed minors. One can say that the indices *decouple* from the integrals.

As an example, we reproduce the 4-point part of

$$\begin{aligned} n_{jkl} I_{4,ijkl}^{[d+]^4} &= \frac{\binom{0}{i}}{\binom{0}{0}_3} \frac{\binom{0}{j}}{\binom{0}{0}_3} \frac{\binom{0}{k}}{\binom{0}{0}_3} \frac{\binom{0}{l}}{\binom{0}{0}_3} d(d+1)(d+2)(d+3) I_4^{[d+]^4} \\ &+ \frac{\binom{0i}{0j} \binom{0}{k} \binom{0}{l} + \binom{0i}{0k} \binom{0}{j} \binom{0}{l} + \binom{0j}{0k} \binom{0}{i} \binom{0}{l} + \binom{0i}{0l} \binom{0}{j} \binom{0}{k} + \binom{0j}{0l} \binom{0}{i} \binom{0}{k} + \binom{0k}{0l} \binom{0}{i} \binom{0}{j}}{\binom{0}{0}_3^3} d(d+1) I_4^{[d+]^3} \\ &+ \frac{\binom{0i}{0l} \binom{0j}{0k} + \binom{0j}{0l} \binom{0i}{0k} + \binom{0k}{0l} \binom{0i}{0j}}{\binom{0}{0}_2^2} I_4^{[d+]^2} + \dots \end{aligned} \tag{28}$$

## Contractions with external momenta [or with CHORDS]

In (28), one has to understand the 4-point integrals to carry the corresponding index  $s$  and the signed minors are  $\binom{0}{k} \rightarrow \binom{0s}{ks}_5$  etc.

## Contractions with external momenta [or with CHORDS] I

A chord is the momentum shift of an internal line due to external momenta,  $D_i = (k - q_i)^2 - m_i^2 + i\epsilon$ , and  $q_i = (p_1 + p_2 + \dots + p_i)$ , with  $q_n = 0$ .

The tensor 5-point integral of rank  $R = 1$  yields, when contracted with a chord,

$$q_{a\mu} I_5^\mu = -\frac{1}{\binom{0}{0}_5} \sum_{s=1}^5 \left[ \sum_{i=1}^4 (q_a \cdot q_i) \binom{0i}{0s}_5 \right] I_4^s. \quad (29)$$

In fact, the sum over  $i$  may be performed explicitly:

$$\Sigma_a^{1,s} \equiv \sum_{i=1}^4 (q_a \cdot q_i) \binom{0s}{0i}_5 = +\frac{1}{2} \left\{ \binom{s}{0}_5 (Y_{a5} - Y_{55}) + \binom{0}{0}_5 (\delta_{as} - \delta_{5s}) \right\},$$

## Contractions with external momenta I

We get immediately

$$q_{a\mu} I_5^\mu = - \frac{1}{\binom{0}{0}_5} \sum_{s=1}^5 \Sigma_a^{1,s} I_4^s. \quad (30)$$

## Contractions with external momenta I

The tensor 5-point integral of rank  $R = 2$

$$I_5^{\mu\nu} = \sum_{i,j=1}^4 q_i^\mu q_j^\nu E_{ij} + g^{\mu\nu} E_{00}, \quad (31)$$

has the following tensor coefficients free of  $1/()_5$ :

$$E_{00} = -\sum_{s=1}^5 \frac{1}{2} \frac{1}{\binom{0}{0}_5} \binom{s}{0}_5 I_4^{[d+],s}, \quad (32)$$

$$E_{ij} = \sum_{s=1}^5 \frac{1}{\binom{0}{0}_5} \left[ \binom{0i}{sj}_5 I_4^{[d+],s} + \binom{0s}{0j}_5 I_{4,i}^{[d+],s} \right]. \quad (33)$$

## Contractions with external momenta I

Equation (31) yields for the contractions with chords:

$$q_{a\mu} q_{b\nu} I_5^{\mu\nu} = \sum_{i,j=1}^4 (q_a \cdot q_i)(q_b \cdot q_j) E_{ij} + (q_a \cdot q_b) E_{00}. \quad (34)$$

and finally (34) simply reads

$$\begin{aligned} q_{a\mu} q_{b\nu} I_5^{\mu\nu} &= \frac{1}{4} \sum_{s=1}^5 \left\{ \frac{\binom{s}{0}_5}{\binom{0s}{0s}_5} (\delta_{ab} \delta_{as} + \delta_{5s}) + \frac{\binom{s}{0s}_5}{\binom{0s}{0s}_5} [(\delta_{as} - \delta_{5s})(Y_{b5} - Y_{55}) \right. \\ &\quad \left. + (\delta_{bs} - \delta_{5s})(Y_{a5} - Y_{55}) + \frac{\binom{s}{0}_5}{\binom{0}{0}_5} (Y_{a5} - Y_{55})(Y_{b5} - Y_{55})] \right\} I_4^{[d+],s} \\ &\quad + \frac{1}{\binom{0}{0}_5} \sum_{s=1}^5 \frac{\Sigma_b^{1,s}}{\binom{0s}{0s}_5} \sum_{t=1}^5 \Sigma_a^{2,st} I_3^{st}, \end{aligned}$$

## Contractions with external momenta I

with

$$\begin{aligned}\Sigma_a^{2,st} &\equiv \sum_{i=1}^4 (q_a \cdot q_i) \binom{0st}{0si}_5 \\ &= \frac{1}{2} (1 - \delta_{st}) \left\{ \binom{ts}{0s}_5 (Y_{a5} - Y_{55}) + \binom{0s}{0s}_5 (\delta_{at} - \delta_{5t}) - \binom{0s}{0t}_5 (\delta_{as} - \delta_{5s}) \right\}.\end{aligned}$$

This has been extended also to higher ranks.

We need at most double sums, e.g.:

$$\begin{aligned}\Sigma_{ab}^{2,s} &\equiv \sum_{i,j=1}^4 (q_a \cdot q_i)(q_b \cdot q_j) \binom{si}{sj}_5 \frac{1}{2} (q_a \cdot q_b) \binom{s}{s}_5 \\ &= -\frac{1}{4} ()_5 (\delta_{ab}\delta_{as} + \delta_{5s}),\end{aligned}\tag{35}$$

## Contractions with external momenta I

Many of the sums over signed minors, weighted with scalar products of chords are given in PLB 2011 [[19]], and an almost complete list may be obtained on request from J. Fleischer, T.R.

## Summary

- Recursive treatment of hexagon and pentagon tensor integrals of rank  $R$  in terms of pentagons and boxes of rank  $R - 1$
- Systematic derivation of expressions which are explicitly free of inverse Gram determinants ()<sub>5</sub> until pentagons of rank  $R = 5$
- Proper isolation of inverse Gram determinants of subdiagrams of the type  $\binom{s}{s}_n 4$ , which cannot be completely avoided
- Numerical C++ package PJFry (V. Yundin, open source) for C, c++, Mathematica, Fortran
- Perform multiple sums with signed minors and scalar products after contractions with chords or external momenta

## References |

-  G. Passarino and M. Veltman, *One loop corrections for  $e^+ e^-$  annihilation into  $\mu^+ \mu^-$  in the Weinberg model*, *Nucl. Phys.* **B160** (1979) 151.
-  G. 't Hooft and M. Veltman, *Scalar one loop integrals*, *Nucl. Phys.* **B153** (1979) 365–401.
-  T. Hahn and M. Perez-Victoria, *Automatized one loop calculations in four-dimensions and D-dimensions*, *Comput.Phys.Commun.* **118** (1999) 153–165, [[hep-ph/9807565](#)].
-  G. J. van Oldenborgh, *FF: A Package to evaluate one loop Feynman diagrams*, *Comput. Phys. Commun.* **66** (1991) 1–15.
-  A. I. Davydychev, *A simple formula for reducing Feynman diagrams to scalar integrals*, *Phys. Lett.* **B263** (1991) 107–111.
-  O. Tarasov, *Connection between Feynman integrals having different values of the space-time dimension*, *Phys. Rev.* **D54** (1996) 6479–6490, [[hep-th/9606018](#)].
-  J. Fleischer, F. Jegerlehner, and O. Tarasov, *Algebraic reduction of one-loop Feynman graph amplitudes*, *Nucl. Phys.* **B566** (2000) 423–440, [[hep-ph/9907327](#)].
-  T. Diakonidis, J. Fleischer, J. Gluza, K. Kajda, T. Riemann, and J. Tausk, *A complete reduction of one-loop tensor 5- and 6-point integrals*, *Phys. Rev.* **D80** (2009) 036003, [[0812.2134](#)].
-  T. Diakonidis, J. Fleischer, T. Riemann, and J. B. Tausk, *A recursive reduction of tensor Feynman integrals*, *Phys. Lett.* **B683** (2010) 69–74, [[0907.2115](#)].

## References II

-  A. Denner, *Techniques and concepts for higher order calculations*, Introductory [Lecture](#) at DESY Theory Workshop on Collider Phenomenology, Hamburg, 29 Sep - 2 Oct 2009.
-  T. Hahn, [LoopTools 2.5 User's Guide](#), [LT25Guide.pdf](#).
-  T. Binoth, J. Guillet, G. Heinrich, E. Pilon, and C. Schubert, *An algebraic / numerical formalism for one-loop multi-leg amplitudes*, *JHEP* **10** (2005) 015, [[hep-ph/0504267](#)].
-  A. Denner and S. Dittmaier, *Reduction schemes for one-loop tensor integrals*, *Nucl. Phys.* **B734** (2006) 62–115, [[hep-ph/0509141](#)].
-  W. Giele, E. W. N. Glover, and G. Zanderighi, *Numerical evaluation of one-loop diagrams near exceptional momentum configurations*, *Nucl. Phys. Proc. Suppl.* **135** (2004) 275–279, [[hep-ph/0407016](#)].
-  J. Fleischer and T. Riemann, *Complete algebraic reduction of one-loop tensor Feynman integrals*, *Phys. Rev.* **D83** (2011) 073004, [[1009.4436](#)].
-  D. Shanks, *Non-linear transformations of divergent and slowly convergent sequences*, *J. Math. Phys.* **34** (1955) 1–42.
-  P. Wynn, *On a device for calculating the  $e_m(s_n)$  transformation*, *Math Tables and A.C.* **10** (1956) 91–96.
-  J. G. A. Baker and P. Graves-Morris, *Pade Approximants. Part I: Basic Theory*. Addison-Wesley, Reading, Massachusetts, 1981.

## References III

-  J. Fleischer and T. Riemann, *Calculating contracted tensor Feynman integrals*, *Phys.Lett.* **B701** (2011) 646–653, [[1104.4067](#)].
-  T. Binoth, J. P. Guillet, G. Heinrich, E. Pilon, and T. Reiter, *Golem95: a numerical program to calculate one-loop tensor integrals with up to six external legs*, *Comput. Phys. Commun.* **180** (2009) 2317–2330, [[0810.0992](#)].
-  T. Riemann, G. Mann, and D. Ebert.  
*Nonconservation of lepton number in Z decay*, in: F. Kaschluhn (ed.), Proc. XVth Int. Symp. Ahrenshoop on Special Topics In Gauge Field Theories, Nov 5-12, 1981, Ahrenshoop, GDR, AdW, Zeuthen (1981) PHE 81-07, pp. 88-91.
-  G. Mann and T. Riemann, *Effective flavor changing weak neutral current in the standard theory and Z boson decay*, *Annalen Phys.* **40** (1984) 334.
-  J. Fleischer and F. Jegerlehner, *Radiative Corrections to Higgs Decays in the Extended Weinberg-Salam Model*, *Phys. Rev.* **D23** (1981) 2001–2026.
-  T. Diakonidis, J. Fleischer, J. Gluza, K. Kajda, T. Riemann, and J. Tausk, *On the tensor reduction of one-loop pentagons and hexagons*, *Nucl. Phys. Proc. Suppl.* **183** (2008) 109–115, [[0807.2984](#)].
-  R. K. Ellis and G. Zanderighi, *Scalar one-loop integrals for QCD*, *JHEP* **02** (2008) 002, [[0712.1851](#)].
-  J. Campbell, E. W. N. Glover, and D. Miller, *One-loop tensor integrals in dimensional regularisation*, *Nucl. Phys.* **B498** (1997) 397–442, [[hep-ph/9612413](#)].

## References IV

-  A. Denner and S. Dittmaier, *Reduction of one-loop tensor 5-point integrals*, *Nucl. Phys.* **B658** (2003) 175–202, [[hep-ph/0212259](#)].
-  T. Binoth, J. P. Guillet, and G. Heinrich, *Reduction formalism for dimensionally regulated one-loop N-point integrals*, *Nucl. Phys.* **B572** (2000) 361–386, [[hep-ph/9911342](#)].
-  Z. Bern, L. J. Dixon, and D. A. Kosower, *Dimensionally regulated pentagon integrals*, *Nucl. Phys.* **B412** (1994) 751–816, [[hep-ph/9306240](#)].
-  G. Ossola, C. Papadopoulos, and R. Pittau, *Reducing full one-loop amplitudes to scalar integrals at the integrand level*, *Nucl. Phys.* **B763** (2007) 147–169, [[hep-ph/0609007](#)].