

# NLO QCD corrections to vector-boson scattering at the LHC

Lucia Hošeková in collaboration with A. Denner

IFIC Valencia





20.3.2012

LHCphenOnet



- 1 Motivation
  - Search for a light Higgs boson
  - Study of Electroweak Symmetry Breaking
- 2 What kinds of diagrams are involved?
- 3 Previous studies
  - Tree-level
  - NLO QCD corrections
- 4 Our project
  - Block Structure
  - Elements of calculation
  - Monte Carlo and cuts
- 5 Summary

# SM Higgs Search







- vector boson fusion ( $qq \rightarrow qqH$ )
  - possible discovery mode for a light SM Higgs boson
  - second largest cross-section for the light Higgs
  - statistically significant thanks to forward jet tagging and suppressed QCD in the central region
  - relatively low luminosity needed for discovery
- vector boson scattering ( $qq \rightarrow qqVV$ ) - background to  $H \rightarrow VV$  decay mode via VBF
- VBF can be used to determine Higgs properties and constraint its couplings to gauge bosons

# No light-Higgs scenario

- the scattering of longitudinal W's grows with energy and violates unitarity





$$\propto g_W^2 \frac{E^2}{M_W^2}$$






$$\propto -g_W^2 \frac{E^2}{M_W^2}$$

- without Higgs, new mechanism of EW symmetry breaking must be considered
- new physics (composite Higgs, extra dimensions,...) predicts new resonances and modify VBF
- $qq \rightarrow qqWW$  very sensitive channel to new interaction
- minimizes the background from transversely polarized WW

# Diagrams - Leading Order



- following types of diagrams have to be included (to preserve gauge invariance)





# Tree-level studies

- first partial results [Cahn, Dawson \(1984\)](#)
- $pp \rightarrow qqWW$  in effective gauge boson approximation, only for longitudinal polarization [Duncan, Kane, Repko \(1986\)](#)
- exact calculation of  $pp \rightarrow qqWW$ , all polarizations [Dicus, Vega \(1986\)](#)
- $pp \rightarrow qqZZ$ , effective gauge boson approximation [Abbasabadi, Repko \(1988\)](#)
- $pp \rightarrow qqZZ \rightarrow qqllll$ , narrow width approximation [Baur, Glover \(1990\)](#)
- $pp \rightarrow (qqZW \rightarrow qqZW) + X$ , effective gauge boson approximation, longitudinal polarization [Dobado, Herrero, Terron \(1991\)](#)
- $pp \rightarrow qqZW$ , full tree-level, leptonic decay correlations [Barger, Cheung, Han, Stange, Zeppenfeld \(1992\)](#)
- $pp \rightarrow qqWW$  - electroweak chiral lagrangian formalism, semileptonic decay [Butterworth, Cox, Forshaw \(2002\)](#)
- $pp \rightarrow qqllll$  - complete parton level analysis, SM and SILH [Ballestrero, Accomando, Bevilacqua, Franzosi, Maina \(2006-2010\)](#)
- multiple BSM studies for the LHC [Han, Krohn, Wang, Zhu \(2009\)](#), [Cheung, Chiang, Yuan \(2008\)](#)...


# Diagrams - NLO QCD Contributions

## Virtual corrections






## Counterterms





## Real corrections





## Initial state gluon

- spacelike gauge bosons





- timelike gauge bosons




# NLO QCD calculation

- full tree-level calculation and NLO QCD corrections (real and virtual contributions)
  - 2006 - Jäger, Oleari, Zeppenfeld:  $qq \rightarrow jjW^+W^- \rightarrow jjllll$
  - 2006 - Jäger, Oleari, Zeppenfeld:  $qq \rightarrow jjZZ \rightarrow jjllll$
  - 2007 - Bozzi, Jäger, Oleari, Zeppenfeld:  $qq \rightarrow jjWZ \rightarrow jjllll$
  - 2009 - Jäger, Oleari, Zeppenfeld:  $qq \rightarrow jjW^\pm W^\pm \rightarrow jjllll$




# Block structure

- EW and QCD parts are completely independent and can be evaluated separately and reused
- introducing so called "leptonic tensors"
- separating QCD and EW blocks
  - simplifies calculation
  - speeds up Monte Carlo simulations






# Block structure

- EW and QCD parts are completely independent and can be evaluated separately and reused
- introducing so called "leptonic tensors"
- separating QCD and EW blocks
  - simplifies calculation
  - speeds up Monte Carlo simulations



# Block structure

- EW and QCD parts are completely independent and can be evaluated separately and reused
- introducing so called "leptonic tensors"
- separating QCD and EW blocks
  - simplifies calculation
  - speeds up Monte Carlo simulations




# How it works - polarization sums

- relatively small number of building blocks required to construct large number of diagrams

$$\mathcal{M} = \mathcal{M}_{QCD\mu} A^\mu = \mathcal{M}_{QCD\mu} g^{\mu\nu} A_\nu \quad \text{and} \quad g_{\mu\nu} = - \sum_i \varepsilon(k)_{i\mu} \varepsilon(k)_{i\nu} + \frac{k_\mu k_\nu}{k^2}$$

$$\begin{aligned} \mathcal{M} &= -(\mathcal{M}_{QCD} \cdot \varepsilon_+)(\mathcal{A} \cdot \varepsilon_+) - (\mathcal{M}_{QCD} \cdot \varepsilon_-)(\mathcal{A} \cdot \varepsilon_-) \\ \Rightarrow & -(\mathcal{M}_{QCD} \cdot \varepsilon_0)(\mathcal{A} \cdot \varepsilon_0) + \frac{1}{k^2}(\mathcal{M}_{QCD} \cdot k)(\mathcal{A} \cdot k) \end{aligned}$$




# How it works - polarization sums

- relatively small number of building blocks required to construct large number of diagrams

$$\mathcal{M} = \mathcal{M}_{QCD\mu} \mathcal{A}^\mu = \mathcal{M}_{QCD\mu} g^{\mu\nu} \mathcal{A}_\nu \quad \text{and} \quad g_{\mu\nu} = - \sum_i \varepsilon(k)_{i\mu} \varepsilon(k)_{i\nu} + \frac{k_\mu k_\nu}{k^2}$$

$$\begin{aligned} \mathcal{M} &= -(\mathcal{M}_{QCD} \cdot \varepsilon_+) (\mathcal{A} \cdot \varepsilon_+) - (\mathcal{M}_{QCD} \cdot \varepsilon_-) (\mathcal{A} \cdot \varepsilon_-) \\ \Rightarrow & -(\mathcal{M}_{QCD} \cdot \varepsilon_0) (\mathcal{A} \cdot \varepsilon_0) + \frac{1}{k^2} (\mathcal{M}_{QCD} \cdot k) (\mathcal{A} \cdot k) \end{aligned}$$





# Elements of Calculation

- diagrams generated with FeynArts
  - EW and QCD blocks are generated independently
- analytical expressions generated with FormCalc and modified in Mathematica and exported to Fortran
- Weyl-van-der-Waerden formalism - translates all kinematic objects into two-component WvdW spinors in chiral representation

$$\Psi = \begin{pmatrix} \phi_A \\ \psi^{\dot{A}} \end{pmatrix} \quad \psi_A \phi^A = (\psi \phi) \quad \psi_{\dot{A}} \phi^{\dot{A}} = \langle \psi \phi \rangle \quad 2k_\mu p^\mu = (k p) \langle k p \rangle$$

- s- and u-channel obtained via crossing which amounts to sign reversal of certain spinors

# Dipole subtraction method

$$\sigma = \int_m d\sigma^B + \int_{m+1} d\sigma^R + \int_m d\sigma^V$$

- $d\sigma^B$  - no singularities,  $d\sigma^R$  - collinear and soft singularities,  $d\sigma^V$  - IR singularities
- analytical integration problematic  $\rightarrow$  dipole subtraction

$$\sigma^{NLO} = \int_{m+1} (d\sigma^R - d\sigma^A) + \int_m (d\sigma^V + \int_1 d\sigma^A)$$

- $d\sigma^A$ : subtraction term containing all IR singularities and analytically integrable over 1-particle phase-space causing collinear and soft singularities
- $d\sigma^B$ ,  $d\sigma^R$ ,  $d\sigma^V$  and colour correlated Born matrix elements  $d\sigma^{B'}$
- IR, soft and collinear singularities are regularized in mass regularization (LoopTools, Coli) or dimensional regularization scheme (Coli)

# Monte Carlo and cuts

- custom-made multi-channel Monte Carlo is being developed
- choosing proper cuts is essential for distinguishing VBF from the background
- 'typical' VBF cuts include
  - tagging jets - two hard reconstructed jets with  $p_T \geq 20$  GeV and large rapidity separation  $\Delta y_{jj} > 4$  and invariant mass  $M_{jj} > 600$  GeV
  - parallel to the beam (within  $1^\circ$ ) -  $|\eta_j| < 4.9$
  - separation of jets and leptons  $\Delta R_{ll} > 0.2$ ,  $\Delta R_{jl} > 0.4$
  - jets in opposite hemispheres  $y_{j1} \times y_{j2} < 0$
  - cuts on invariant masses of the leptons

# Outlook

- all helicity MEs ( $qq \rightarrow jj4l$ ) incl. t-, s- and u-channel and their interferences  
- completed
- LO and real corrections ME's compared with independent results (MadGraph, FormCalc,...)
- total LO cross section compared with existing results (Zeppenfeld, Ballestrero) with different sets of cuts
- implementation of the custom-made MC at NLO in progress
- possibility of incorporating EW corrections and BSM physics in the future

# Summary

- vector boson scattering might turn out to be principal for
  - search for a light Higgs at the LHC
  - probing for new strong effects in the no-Higgs scenario
- full NLO corrections are fairly complicated (many legs, large number of diagrams) and require modular approach
- despite long history of studies NLO calculations started to emerge only recently (and only QCD corrections)
- our project - first independent verification of Zeppenfeld's calculation