

# Diphoton production at LHC (NNLO)

Leandro Cieri  
Universidad de Buenos Aires - Argentina  
& INFN Sezione di Firenze  
LHCphenonet

LHCphenonet Annual Meeting 2012  
March 22, 2012

# *Outline*

- Introduction
- Available theoretical tools
- Diphoton production with  $2\gamma$ NNLO
- Summary

In collaboration with S. Catani, D. de Florian, G. Ferrera and M. Grazzini

# *Outline*

## 📌 Introduction

- 📌 Why is diphoton production important?
- 📌 Photon production mechanisms and isolation

## 📌 Theoretical tools available

## 📌 Diphoton production with $2\gamma$ NNLO

## 📌 Summary

In collaboration with S. Catani, D. de Florian, G. Ferrera and M. Grazzini

# *Outline*

- Introduction
- Available theoretical tools
- Diphoton production with  $2\gamma$ NNLO
  - Features of the code
  - Results
- Summary

In collaboration with S. Catani, D. de Florian, G. Ferrera and M. Grazzini

# **Why is diphoton production important?**


- ➊ It is a channel that we can use to check the validity of perturbative Quantum Chromodynamics (pQCD)
  - ➌ Collinear factorization approach
  - ➌  $K_T$  factorization approach
  - ➌ Soft gluon logarithmic resummation techniques
  
- ➋ It constitutes an irreducible background for new physics searches
  - ➌ Universal Extra Dimensions
  - ➌ Randall-Sundrum ED
  - ➌ Supersymmetry
  - ➌ New heavy resonances
  
- ➌ **Irreducible background**
  - ➌ **In searches for a low mass Higgs boson decaying into photon pairs**

# **Why is diphoton production important?**

- ➊ It is a channel that we can use to check the validity of perturbative Quantum Chromodynamics (pQCD)
  - ➌ Collinear factorization approach
  - ➌  $K_T$  factorization approach
  - ➌ Soft gluon logarithmic resummation techniques
  
- ➋ It constitutes an irreducible background for new physics searches
  - ➌ Universal Extra Dimensions
  - ➌ Randall-Sundrum ED
  - ➌ Supersymmetry
  - ➌ New heavy resonances


## **Irreducible background**

**➌ In searches for a low mass Higgs boson decaying into photon pairs**




# The search for the SM Higgs boson

- Direct searches at LEP2 experiments  
Phys. Lett. B 565 (2003) 61



One of the most promising channels at the LHC is the rare decay of the Higgs boson into a pair of photons

$$H \rightarrow \gamma\gamma$$



- Combined results ATLAS - CMS  
Phys. Lett. B 705 (2011) 452–470  
ATLAS-CONF-2011-149

CMS-PAS-HIG-11-021  
arXiv:1201.3084 [hep-ph]



# The search for the SM Higgs boson

- Direct searches at LEP2 experiments  
Phys. Lett. B 565 (2003) 61




One of the most promising channels at the LHC is the rare decay of the Higgs boson into a pair of photons

$$H \rightarrow \gamma\gamma$$

In order to understand the signal we have to control the background to this process in the best way that we can.

- Combined results ATLAS - CMS  
Phys. Lett. B 705 (2011) 452–470  
ATLAS-CONF-2011-149

CMS-PAS-HIG-11-021  
arXiv:1201.3084 [hep-ph]




# The search for the SM Higgs boson

- Direct searches at LEP2 experiments  
Phys. Lett. B 565 (2003) 61

$M_{\text{Higgs}} < 114 \text{ GeV}/c^2$   
(95% C.L.)

## After Moriond (CMS)

$114.4 \text{ GeV}/c^2 < M_{\text{Higgs}} < 127.5 \text{ GeV}/c^2$   
(95% C.L.)



## After Moriond (ATLAS)

$117.5 \text{ GeV}/c^2 < M_{\text{Higgs}} < 118.5 \text{ GeV}/c^2$   
 $122.5 \text{ GeV}/c^2 < M_{\text{Higgs}} < 129 \text{ GeV}/c^2$   
(95% C.L.)


One of the most promising channels at the LHC is the rare decay of the Higgs boson into a pair of photons

$$H \rightarrow \gamma\gamma$$

In order to understand the signal we have to control the background to this process in the best way that we can.

- Combined results ATLAS - CMS  
Phys. Lett. B 705 (2011) 452–470  
ATLAS-CONF-2011-149


CMS-PAS-HIG-11-021  
arXiv:1201.3084 [hep-ph]




$141 \text{ GeV}/c^2 < M_{\text{Higgs}} < 476 \text{ GeV}/c^2$   
(95% C.L.)

# Photon production

When dealing with the production of photons we have to consider two production mechanisms:



**Direct component:** photon directly produced through the hard interaction



Fragmentation function:  
to be fitted from data

**Fragmentation component:** photon produced from non-perturbative fragmentation of a hard parton (analogously to a hadron)  
Single and double resolved (collinear fragmentation)  
Calculations of cross sections with photons have additional singularities in the presence of QCD radiation.  
(i.e. When we go beyond LO)


When quark and photon are collinear  $\rightarrow$  singular propagator

See R. Hernandez-Pinto's and  
P. Artoisenet's presentation!

# Photon production

- Experimentally photons must be isolated
- Isolation reduces fragmentation component
- Experimentalist may choose:

$$\sum_{\delta < R_0} E_T^{had} \leq \epsilon_\gamma p_T^\gamma$$




$$\sum_{\delta < R_0} E_T^{had} \leq E_T^{max}$$

Using conventional isolation, only the sum of the direct and fragmentation contributions is meaningful.

# Photon production

- Experimentally photons must be isolated
- Isolation reduces fragmentation component
- Experimentalist may choose:

$$\sum_{\delta < R_0} E_T^{had} \leq \epsilon_\gamma p_T^\gamma$$



$$\sum_{\delta < R_0} E_T^{had} \leq E_T^{max}$$

Using conventional isolation, only the sum of the direct and fragmentation contributions is meaningful.

But there is a way to isolate and make the direct cross section physical  
(Infrared safe)

## Smooth cone Isolation

S. Frixione, Phys.Lett. B429 (1998) 369-374,

Soft emission allowed arbitrarily close to the photon

$$\chi(\delta) = \epsilon_\gamma E_T^\gamma \left( \frac{1 - \cos(\delta)}{1 - \cos(R_0)} \right)^n$$


- no quark-photon collinear divergences
- no fragmentation component (only direct)
- direct well defined by itself

$E_T^{had}(\delta) \leq \chi(\delta)$  such that  $\lim_{\delta \rightarrow 0} \chi(\delta) = 0$

# *Available theoretical tools*


**DIPHOX** Full NLO for direct and fragmentation  
+ Box contribution (one piece of NNLO)

T. Binoth, J.Ph. Guillet, E. Pilon and M. Werlen



**gamma2MC** Full NLO (direct only) + Box  
+ correction to Box contribution partial N<sup>3</sup>LO term

Zvi Bern, Lance Dixon, and Carl Schmidt



**MCFM** Full NLO for direct, but only LO for fragmentation  
+ correction to Box contribution partial N<sup>3</sup>LO term

John M. Campbell, R.Keith Ellis, Ciaran Williams

**Resbos** NLL q<sub>T</sub> resummation for direct (with regulator  
for collinear singularities)  
C. Balázs, E. L. Berger, P. Nadolsky, and C.-P. Yuan  
Correction to Box contribution partial N<sup>3</sup>LO term

# **Available theoretical tools**

**DIPHOX** Full NLO for direct and fragmentation  
+ Box contribution (one piece of NNLO)

T. Binoth, J.Ph. Guillet, E. Pilon and M. Werlen

**gamma2MC** Full NLO (direct only) + Box  
+ correction to Box contribution partial N<sup>3</sup>LO term

Zvi Bern, Lance Dixon, and Carl Schmidt

**MCFM** Full NLO for direct, but only LO for fragmentation  
+ correction to Box contribution partial N<sup>3</sup>LO term

John M. Campbell, R.Keith Ellis, Ciaran Williams

**Resbos** NLL q<sub>T</sub> resummation for direct (with regulator  
C. Balázs, E. L. Berger, P. Nadolsky, and C.-P. Yuan for collinear singularities)  
Correction to Box contribution partial N<sup>3</sup>LO term

Results typically in good agreement with data, but some differences observed:

- **Azimuth separation for diphoton production**
- **Low mass region of the invariant mass distribution**

**It is desireable to count on a NNLO description of the phenomenology of diphoton production**

# Diphoton production with 2 $\gamma$ NNLO

- Based on the  $q_T$  subtraction formalism
- Fully exclusive NNLO description (direct contribution) for  $pp(\bar{p}) \rightarrow \gamma\gamma$
- No fragmentation contribution
- Also corrections to Box contribution, partial N<sup>3</sup>LO terms available

Zvi Bern, Lance Dixon, and Carl Schmidt








S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

S. Catani, M. Grazzini

Frixione Isolation

(Available, but not present in the following analysis)

Full NNLO means full control of the  $\mathcal{O}(\alpha_s^2)$  diagrams:



+ ...

# Diphoton production with 2 $\gamma$ NNLO

- Based on the  $q_T$  subtraction formalism
- Fully exclusive NNLO description (direct contribution) for  $pp(\bar{p}) \rightarrow \gamma\gamma$
- No fragmentation contribution
- Also corrections to Box contribution, partial N<sup>3</sup>LO terms available

Zvi Bern, Lance Dixon, and Carl Schmidt


S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

S. Catani, M. Grazzini

Frixione Isolation

(Available, but not present in the following analysis)

Full NNLO means full control of the  $\mathcal{O}(\alpha_s^2)$  diagrams:




# $q_T$ subtraction method

S. Catani, M. Grazzini (2007)

Let us consider a specific, though important class of processes: the production of colourless high-mass systems  $F$  in hadron collisions  
(  $F$  may consist of lepton pairs, vector bosons, Higgs bosons.....)

At LO it starts with  $c\bar{c} \rightarrow F$



**Strategy:** start from NLO calculation of  $F+\text{jet(s)}$  and observe that as soon as the transverse momentum of the  $F$ ,  $q_T \neq 0$ , one can write:

$$d\sigma_{(N)NLO}^F|_{q_T \neq 0} = d\sigma_{(N)LO}^{F+\text{jets}}$$

Define a counterterm to deal with singular behaviour at  $q_T \rightarrow 0$

But.....

the singular behaviour of  $d\sigma_{(N)LO}^{F+\text{jets}}$  is well known from the resummation program of large logarithmic contributions at small transverse momenta

G. Parisi, R. Petronzio (1979)

J. Collins, D.E. Soper, G. Sterman (1985)

S. Catani, D. de Florian, M.Grazzini (2000)

# $q_T$ subtraction method

S. Catani, M. Grazzini (2007)

choose

$$d\sigma^{CT} \sim d\sigma^{(LO)} \otimes \Sigma^F(q_T/Q)$$

where

$$\Sigma^F(q_T/Q) \sim \sum_{n=1}^{\infty} \left(\frac{\alpha_S}{\pi}\right)^n \sum_{k=1}^{2n} \Sigma^{F(n;k)} \frac{Q^2}{q_T^2} \ln^{k-1} \frac{Q^2}{q_T^2}$$

Then the calculation can be extended to include the  $q_T = 0$  contribution:

$$d\sigma_{(N)NLO}^F = \mathcal{H}_{(N)NLO}^F \otimes d\sigma_{LO}^F + \left[ d\sigma_{(N)LO}^{F+\text{jets}} - d\sigma_{(N)LO}^{CT} \right]$$

where I have subtracted the truncation of the counterterm at (N)LO and added a contribution at  $q_T = 0$  to restore the correct normalization

The function  $\mathcal{H}^F$  can be computed in QCD perturbation theory

$$\mathcal{H}^F = 1 + \left(\frac{\alpha_S}{\pi}\right) \mathcal{H}^{F(1)} + \left(\frac{\alpha_S}{\pi}\right)^2 \mathcal{H}^{F(2)} + \dots$$

# $q_T$ subtraction method

S. Catani, M. Grazzini (2007)

For a generic  $pp \rightarrow F + X$  process:

- At NLO we need a LO calculation of  $d\sigma^{F+\text{jet(s)}}$  plus the knowledge of  $d\sigma_{LO}^{CT}$  and  $\mathcal{H}^{F(1)}$ 
  - the counterterm  $d\sigma_{LO}^{CT}$  requires the resummation coefficients  $A^{(1)}, B^{(1)}$  and the one loop anomalous dimensions
  - the general form of  $\mathcal{H}^{F(1)}$  is known D. de Florian, M. Grazzini (2000)  
G. Bozzi, S. Catani, D. de Florian, M. Grazzini (2005)
- At NNLO we need a NLO calculation of  $d\sigma^{F+\text{jet(s)}}$  plus the knowledge of  $d\sigma_{NLO}^{CT}$  and  $\mathcal{H}^{F(2)}$ 
  - the counterterm  $d\sigma_{NLO}^{CT}$  depends also on the resummation coefficients  $A^{(2)}, B^{(2)}$  and on the two loop anomalous dimensions
  - we have computed  $\mathcal{H}^{F(2)}$  for Higgs and vector boson production!
  - generalized to any process with final state colorless system **F**

S. Catani, M. Grazzini (2007)

S. Catani, L. C, G.Ferrera, D. de Florian, M. Grazzini (2009)

S. Catani, L. C, G.Ferrera, D. de Florian, M. Grazzini (2011)

# $q_T$ subtraction method

S. Catani, M. Grazzini (2007)

For a generic  $pp \rightarrow F + X$  process:

This is enough to compute NNLO corrections for any process in this class provided that  $F + \text{jet}$  is known up to NLO and the two loop amplitude for  $\bar{c}c \rightarrow F$  is known

- At NNLO we need a NLO calculation of  $d\sigma^{F+\text{jet}(s)}$  plus the knowledge of  $d\sigma_{NLO}^{CT}$  and  $\mathcal{H}^{F(2)}$ 
  - the counterterm  $d\sigma_{NLO}^{CT}$  depends also on the resummation coefficients  $A^{(2)}, B^{(2)}$  and on the two loop anomalous dimensions
  - we have computed  $\mathcal{H}^{F(2)}$  for Higgs and vector boson production!
  - generalized to any process with final state colorless system  $F$

S. Catani, M. Grazzini (2007)

S. Catani, L. Cieri, G.Ferrera, D. de Florian, M. Grazzini (2009)

S. Catani, L. Cieri, G.Ferrera, D. de Florian, M. Grazzini (2011)

# $q_T$ subtraction method

S. Catani, M. Grazzini (2007)

**In our case**

## **DiPhoton production at NNLO**

Two-loop amplitudes available C.Anastasiou, E.W.N.Glover, M.E.Tejeda-Yeomans

Di-photon + jet at NLO computed V.Del Duca, F.Maltoni, Z.Nagy, Z.Trocsanyi

implemented in NLOJet++

# $q_T$ subtraction method

S. Catani, M. Grazzini (2007)

In our case

**DiPhoton production at NNLO**

$\mathcal{H}^{F(2)}$

Two-loop amplitudes available • C.Anastasiou, E.W.N.Glover, M.E.Tejeda-Yeomans

Di-photon + jet at NLO computed V.Del Duca, F.Maltoni, Z.Nagy, Z.Trocsanyi

implemented in NLOJet++

# $q_T$ subtraction method

S. Catani, M. Grazzini (2007)

In our case

**DiPhoton production at NNLO**

Two-loop amplitudes available

• C.Anastasiou, E.W.N.Glover, M.E.Tejeda-Yeomans

Di-photon + jet at NLO computed

• V.Del Duca, F.Maltoni, Z.Nagy, Z.Trocsanyi

implemented in NLOJet++

# $q_T$ subtraction method

S. Catani, M. Grazzini (2007)

In our case

**DiPhoton production at NNLO**


Two-loop amplitudes available

• C.Anastasiou, E.W.N.Glover, M.E.Tejeda-Yeomans

Di-photon + jet at NLO computed

• V.Del Duca, F.Maltoni, Z.Nagy, Z.Trocsanyi

implemented in NLOJet++



Fully exclusive NNLO code for  $pp \rightarrow F$

**2 $\gamma$ NNLO**


First exclusive NNLO in pp collisions with two final state particles  
S.Catani, L.Cieri, D.de Florian, G.Ferrera, M.Grazzini (2011)

# Diphoton production at NNLO

S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

**First** results using  $2\gamma$ NNLO



- $\sqrt{S} = 14 \text{ TeV}$
- $p_T^{\gamma \text{ hard}} \geq 40 \text{ GeV}$
- $p_T^{\gamma \text{ soft}} \geq 25 \text{ GeV}$
- $|\eta^\gamma| \leq 2.5$
- $20 \text{ GeV} \leq M_{\gamma\gamma} \leq 250 \text{ GeV}$
- $\mu_R = \mu_F = M_{\gamma\gamma}$

NNLO effect about +50 % in the peak region


Box only  $\sim 22\%$  of NNLO correction

# Diphoton production at NNLO

S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

**First** results using  $2\gamma$ NNLO



$$\sqrt{S} = 14 \text{ TeV}$$

$$p_T^{\gamma hard} \geq 40 \text{ GeV}$$

$$p_T^{\gamma soft} \geq 25 \text{ GeV}$$

$$|\eta^\gamma| \leq 2.5$$

$$20 \text{ GeV} \leq M_{\gamma\gamma} \leq 250 \text{ GeV}$$

$$\mu_R = \mu_F = M_{\gamma\gamma}$$

$$\frac{\sigma^{NNLO}}{\sigma^{NLO+Box}} \sim 1.35$$


$$\frac{\sigma^{NNLO}}{\sigma^{NLO}} \sim 1.55$$

# Diphoton production at NNLO

S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

$p_T$  of harder and softer photon




The requirement  
 $p_{T1}^\gamma \geq 40 \text{ GeV}$   
implies that at LO  
also the  
softer photon  
must have  
 $p_T^\gamma \geq 40 \text{ GeV}$

# Diphoton production at NNLO

S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

$p_T$  of harder and softer photon




- Substantial contribution from radiation in the region  $25 \text{ GeV} < pT < 40 \text{ GeV}$
- Unphysical peak in  $p_T^\gamma$  at  $p_T^\gamma = 40 \text{ GeV}$

# Diphoton production at NNLO


S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

Discrepancy between NLO and experimental data



PRD 85, 012003 (2012)




Same discrepancies found by CDF: Phys.Rev.Lett.107:102003,2011.

# Diphoton production at NNLO


S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

Discrepancy between NLO and experimental data



PRD 85, 012003 (2012)



JHEP 01(2012)133


Same discrepancies found by CDF: Phys.Rev.Lett.107:102003,2011.

# Diphoton production at NNLO

S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

Discrepancy between NLO and experimental data




Same discrepancies found by CDF: Phys.Rev.Lett.107:102003,2011.

# Diphoton production at NNLO


S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

Discrepancy between NLO and experimental data



PRD 85, 012003 (2012)



JHEP 01(2012)133


Same discrepancies found by CDF: Phys.Rev.Lett.107:102003,2011.

# Diphoton production at NNLO

S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

Discrepancy between NLO and experimental data




Same discrepancies found by CDF: Phys.Rev.Lett.107:102003,2011.

# Diphoton production at NNLO

S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC

First exclusive NNLO with two final state particles

Discrepancy between NLO and experimental data at low  $\Delta\phi_{\gamma\gamma}$




# Diphoton production at NNLO

Preliminary results

S.Catani, D. de Florian, G.Ferrera, M.Grazzini, LC


NNLO Corrections much larger in some kinematical regions  
NLO effectively lowest order

“away from back-to-back configuration”




NNLO corrections essential to understand the background

invariant mass below the LO threshold



"No back-to-back"



This discrepancy can be related to the discrepancy observed in the  $\Delta\phi_{\gamma\gamma}$  distribution.


# Summary


- Sizeable NNLO corrections to the  $\gamma\gamma$  mass distribution in kinematical regions related to Higgs boson searches  
40-55% effect over NLO
- NNLO very large away from back-to-back configuration (effectively NLO)  
needed to understand LHC data
- At NNLO starts to reliably predict values of cross sections in all kinematical regions (with very few exceptions; e.g  $p_{T\gamma\gamma} \rightarrow 0$ )
- Cross section with “smooth” isolation, is a lower bound for cross section with standard isolation.
- Work in progress: release a public version of **2 $\gamma$ NNLO**  
+ approximation of standard isolation

# *Backup Slides*


# Why do we need NNLO corrections?

NNLO QCD corrections in diphoton production

$\gamma\gamma$  production  some NNLO terms known to be as large as Born!



$O(\alpha_s^2)$  but  $gg$  Luminosity




$O(\alpha_s^0)$  but  $q\bar{q}$  Luminosity


- Box contribution already included in NLO calculation

DIPHOX: T.Binoth, J.P.Gillet, E.Pilon,  
M.Werlen


# Why do we need NNLO corrections?

NNLO QCD corrections in diphoton production

$\gamma\gamma$  production  some NNLO terms known to be as large as Born!



$O(\alpha_s^2)$  but  $gg$  Luminosity




$O(\alpha_s^0)$  but  $q\bar{q}$  Luminosity

- Box contribution already included in NLO calculation      DIPHOX: T.Binoth, J.P.Gillet, E.Pilon, M.Werlen
- Full NNLO control of Di-photon production is desired (main light Higgs bkg)

# Diphoton production at NNLO

S.Catani, L.Cieri, D. de Florian, G.Ferrera, M.Grazzini

First exclusive NNLO with two final state particles



**Tevatron**


$p_T^\gamma \geq 17 \text{ GeV}$   
 $p_T^\gamma \geq 15 \text{ GeV}$   
 $|\eta^\gamma| < 1$

- Impact of NNLO corrections a bit smaller than at the LHC but still important
- NNLO effect about +30%

# Diphoton production at NNLO

S.Catani, L.Cieri, D. de Florian, G.Ferrera, M.Grazzini

First exclusive NNLO with two final state particles



$$\sqrt{s} = 7 \text{ TeV}$$

$$p_T^{\gamma \text{ hard}} \geq 40 \text{ GeV}$$

$$p_T^{\gamma \text{ soft}} \geq 30 \text{ GeV}$$

$$100 \text{ GeV} \leq M_{\gamma\gamma} \leq 160 \text{ GeV}$$

$$|\eta^\gamma| \leq 2.5$$

$$\text{excluding } 1.4442 \leq |\eta^\gamma| \leq 1.566$$


$$\epsilon = 0.05$$

- Scale does not represent TH uncertainties at LO and NLO → new channels
- All channels open at NNLO → estimate of TH uncertainties

# Diphoton production at NNLO

S.Catani, L.Cieri, D. de Florian, G.Ferrera, M.Grazzini

First exclusive NNLO with two final state particles



- Scale does not represent TH uncertainties at LO and NLO → **new channels**
- All channels open at NNLO → **estimate of TH uncertainties**

$$\sqrt{s} = 7 \text{ TeV}$$

$$p_T^{\gamma \text{ hard}} \geq 40 \text{ GeV}$$

$$p_T^{\gamma \text{ soft}} \geq 30 \text{ GeV}$$


$$100 \text{ GeV} \leq M_{\gamma\gamma} \leq 160 \text{ GeV}$$

$$|\eta^\gamma| \leq 2.5$$

$$\text{excluding } 1.4442 \leq |\eta^\gamma| \leq 1.566$$

$$\epsilon = 0.05$$

$$\alpha_s^3 \quad \text{Bern, Dixon, Schmidt (2002)}$$




Some  $\text{N}^3\text{LO}$  terms known to contribute  $\sim 5\%$

# Diphoton production at NNLO

S.Catani, L.Cieri, D. de Florian, G.Ferrera, M.Grazzini

First exclusive NNLO with two final state particles



- Scale does not represent TH uncertainties at LO and NLO → **new channels**
- All channels open at NNLO → **estimate of TH uncertainties**

$$\sqrt{s} = 7 \text{ TeV}$$

$$p_T^{\gamma \text{ hard}} \geq 40 \text{ GeV}$$

$$p_T^{\gamma \text{ soft}} \geq 30 \text{ GeV}$$



$$100 \text{ GeV} \leq M_{\gamma\gamma} \leq 160 \text{ GeV}$$

$$|\eta^\gamma| \leq 2.5$$

$$\text{excluding } 1.4442 \leq |\eta^\gamma| \leq 1.566$$



$$\epsilon = 0.05$$

$$\alpha_s^3 \quad \text{Bern, Dixon, Schmidt (2002)}$$



Some **N<sup>3</sup>LO** terms known to contribute ~5%

# With Higgs search cuts at 7 TeV



$$p_T^{\gamma \text{ hard}} \geq 40 \text{ GeV}$$

$$p_T^{\gamma \text{ soft}} \geq 25 \text{ GeV}$$

$$100 \text{ GeV} \leq M_{\gamma\gamma} \leq 160 \text{ GeV}$$

$$|\eta^\gamma| \leq 2.37$$

excluding  $1.37 \leq |\eta^\gamma| \leq 1.52$

$$\epsilon = 0.05$$

$$p_T^{\gamma \text{ hard}} \geq 40 \text{ GeV}$$

$$p_T^{\gamma \text{ soft}} \geq 30 \text{ GeV}$$


$$100 \text{ GeV} \leq M_{\gamma\gamma} \leq 160 \text{ GeV}$$

$$|\eta^\gamma| \leq 2.5$$

excluding  $1.4442 \leq |\eta^\gamma| \leq 1.566$

$$\epsilon = 0.05$$

# Channels



# Higgs search at 7 TeV

