



Contribution ID: 199

Type: Poster

## **O(4) scaling analysis in two-flavor QCD at finite temperature and density with improved Wilson quarks**

*Tuesday, July 26, 2016 8:00 PM (1 hour)*

We study scaling behavior of a chiral order parameter in the low density region, performing a simulation of two-flavor lattice QCD with improved Wilson quarks. It has been confirmed that the scaling behavior of the chiral order parameter defined by a Ward-Takahashi identity agrees with the scaling function of the three-dimensional O(4) spin model at zero chemical potential. We discuss the scaling properties of the chiral phase transition at finite density, applying the reweighting method and calculating derivatives of the chiral order parameter with respect to the chemical potential. In the comparison between the scaling functions of the O(4) spin model and QCD at low density, there is a fit parameter which can be interpreted as the curvature of the chiral phase transition curve in the QCD phase diagram with respect to temperature and chemical potential. We determine the curvature of the phase boundary by the fitting. The physical scale is set by the gradient flow.

**Author:** Dr UMEDA, Takashi (Hiroshima Univ.)

**Co-authors:** Mr UJI, Atsushi (Niigata University); Dr OHNO, Hiroshi (University of Tsukuba); Prof. KANAYA, Kazuyuki (CiRfSE, Univ. Tsukuba); Mr WAKABAYASHI, Naoki (Niigata University); Mr IWAMI, Ryo (Niigata University); Dr EJIRI, Shinji (Niigata University); Dr YOSHIDA, Shinsuke (Central China Normal Univ.)

**Presenter:** Dr UMEDA, Takashi (Hiroshima Univ.)

**Session Classification:** Poster

**Track Classification:** Nonzero Temperature and Density