

Charm physics: f_D and f_{D_s} status and outlook

Justus Tobias Tsang

University of Southampton

29 July 2016



RBC-UKQCD Collaborations

BNL and RBRC

Mattia Bruno
Tomomi Ishikawa
Taku Izubuchi
Chulwoo Jung
Christoph Lehner
Meifeng Lin
Taichi Kawanai
Hiroshi Ohki
Shigemi Ohta (KEK)
Amarjit Soni
Sergey Syritsyn

CERN

Marina Marinkovic
Columbia University

Ziyuan Bai
Norman Christ
Luchang Jin

Christopher Kelly
Bob Mawhinney
Greg McGlynn
David Murphy
Daiqian Zhang
Jiqun Tu

University of Connecticut

Tom Blum

Edinburgh University

Peter Boyle
Guido Cossu
Luigi Del Debbio
Richard Kenway
Julia Kettle

Ava Khamseh
Brian Pendleton
Antonin Portelli
Oliver Witzel
Azusa Yamaguchi

K.E.K.
Julien Frison
Peking University
Xu Feng
Plymouth University
Nicolas Garron
University of Southampton
Jonathan Flynn
Vera Gülpers
James Harrison
Andreas Jüttner
Andrew Lawson
Edwin Lizarazo
Chris Sachrajda
Francesco Sanfilippo
Matthew Spraggs
Tobias Tsang
York University (Toronto)
Renwick Hudspith




Outline

1 Motivation

2 Decay Constant Analysis

3 Current Projects

Motivation - Flavour Physics



CKMfitter Group (J. Charles et al.), Eur. Phys. J. C41, 1-131 (2005) [hep-ph/0406184], updated results and plots available at: <http://ckmfitter.in2p3.fr>

Experiment


- Belle, BaBar, CLEO-c
- LHCb, Belle II

Theory

- K , D and B physics to test unitarity of the CKM matrix.

⇒ Place tight bounds on SM predictions

D and B physics



Goal: $D_{(s)}$ and $B_{(s)}$ phenomenology

- masses
- decay constants
- semi-leptonics

Outcome:

- Test of SM
- Reduce systematic errors by direct computation

Charm physics in the RBC/UKQCD program


- Charm phenomenology (f_D , f_{D_s} , Bag, ξ , semi-leptonics)
- 2+1+1f ensemble generation
- GIM mechanism
(see Rare Kaon talk by **A. Lawson Thurs 17⁵⁰**)
- Charm contribution to the HVP
(see talk by **M. Spraggs Tue 17¹⁰**)
- Semi-leptonic B decays in RHQ e.g. $B \rightarrow D$
(see talks by **E. Lizarazo Fri 17¹⁰** & **O. Witzel Wed 10²⁰**)
- Ratio Method to push from c to b (*arXiv:0909.3187*)

Our Action: Domain Wall Fermions

- Chiral fermions
- Automatically $\mathcal{O}(a)$ -improved
- **Physical Pion Mass ensembles:** Moebius Domain Wall Fermions (*arXiv:0409118*)
- Tiny Chiral extrapolation is done with Shamir DWFs (*arXiv:9303005*)
- Tested with quenched **PILOT STUDY.**
⇒ **Quenched study as proof of concept:**
JHEP 05 (2015) 072 (*arXiv:1504.01630 [hep-lat]*)
JHEP 04 (2016) 037 (*arXiv:1602.04118 [hep-lat]*)

Dynamical Ensembles

2 ensembles with physical pion masses



$L^3 \times T/a^4$	a^{-1}/GeV	m_π/MeV
$48^3 \times 96$	1.73	139
$24^3 \times 64$	1.78	340
$24^3 \times 64$	1.78	430
$64^3 \times 128$	2.36	139
$32^3 \times 64$	2.38	300
$32^3 \times 64$	2.38	360
$48^3 \times 96$	2.77	230

(C0 + M0: arXiv:1411.7017)

Set up of Simulation

Valence Quarks:

- light and strange: unitary ($M_5 = 1.8$)
 - heavy from quenched study: (*arXiv:1602.04118*)
 $am_h \leq 0.4$, $M_5 = 1.6$, $L_s = 12$
 - Expect same qualitative features in $N_f = 2 + 1$
- ⇒ **Mixed action** between the (light+strange) and the heavy quark sector.



Computational Details:

- Volume averaging by using \mathbb{Z}_2 -Wall sources.
- HDCG (*arXiv:1402.2585*) for light and strange. CG for heavy
- Monitored time-slice residual (*arXiv:0508023*):
$$r_t = \max_t \frac{|D\psi - \eta|_t}{|\psi|_t}$$
- Many source position per configuration

Decay Constant Data

Masses and MEs from combined fit to $\langle AP \rangle$ and $\langle PP \rangle$ including 1st excited states: $\mathcal{O}(a, m_l, m_s, m_h)$.¹


$$\Phi_P = f_P \sqrt{m_P}$$



¹so far: Z_A from conserved light-light current

Ratio of Bag parameters ξ

$$B_P = \frac{\langle P^0 | O_{VV+AA} | \bar{P}^0 \rangle}{\frac{8}{3} f_P^2 m_P^2}, \quad \xi = \frac{f_{hs} \sqrt{B_{hs}}}{f_{hl} \sqrt{B_{hl}}}$$



PLAN:

- Renormalisation of mixed action for Bag parameters.
(A. Khamseh Wed 10²⁰)
- Extrapolate to B via ratio method. arXiv:0909.3187

Analysis

$$\mathcal{O}(a, m_I, m_s, m_h) \rightarrow \mathcal{O}(a = 0, m_I^{\text{phys}}, m_s^{\text{phys}}, m_c^{\text{phys}})$$

- Extrapolate to m_s^{phys} (from global fit arXiv:1411.7017).
- Global fit: CL+heavy quark mass + light quark mass

$$\begin{aligned} \mathcal{O}(a, m_I, m_h) &= \mathcal{O}(0, m_I^{\text{phys}}, m_h^{\text{phys}}) \\ &\quad + C_{CL}^0 a^2 \\ &\quad + C_\chi^0 \left(m_\pi^2 - m_\pi^{2\text{phys}} \right) \\ &\quad + C_{P_h}^0 \Delta m_{P_h}^{-1} \end{aligned} \tag{1}$$

where $\Delta m_{P_h}^{-1} = m_{P_h}^{-1} - m_{P_h}^{\text{phys}}{}^{-1}$ with $P_h = D, D_s, \eta_c$

Analysis

$$\mathcal{O}(a, m_l, m_s, m_h) \rightarrow \mathcal{O}(a = 0, m_l^{\text{phys}}, m_s^{\text{phys}}, m_c^{\text{phys}})$$

- Extrapolate to m_s^{phys} (from global fit arXiv:1411.7017).
- Global fit: CL+heavy quark mass + light quark mass

$$\begin{aligned} \mathcal{O}(a, m_l, m_h) &= \mathcal{O}(0, m_l^{\text{phys}}, m_h^{\text{phys}}) \\ &\quad + [C_{CL}^0 + C_{CL}^1 \Delta m_{P_h}^{-1}] a^2 \\ &\quad + [C_\chi^0 + C_\chi^1 \Delta m_{P_h}^{-1}] (m_\pi^2 - m_\pi^{2\text{phys}}) \\ &\quad + [C_{P_h}^0] \Delta m_{P_h}^{-1} \end{aligned} \tag{1}$$

where $\Delta m_{P_h}^{-1} = m_{P_h}^{-1} - m_{P_h}^{\text{phys}-1}$ with $P_h = D, D_s, \eta_c$

Strange Quark Mass Correction

- Slight mistuning between unitary and physical strange quark mass.

ensemble	am_s^{unitary}	am_s^{physical}	mismatch
coarse	0.03620	0.03580(16)	-1.1%
medium	0.02661	0.02539(17)	-4.8%
fine	0.02144	0.02132(17)	-0.6%

- Parameterise mistuning in terms of dimensionless α :


$$\mathcal{O}^{\text{phys}} = \mathcal{O}^{\text{uni}} \left(1 + \alpha_{\mathcal{O}} \frac{m_s^{\text{phys}} - m_s^{\text{uni}}}{m_s^{\text{phys}}} \right)$$

- Find $\alpha_{\mathcal{O}}$ from subset of ensembles and apply to others.

Strange Quark Mass Correction

$$\mathcal{O}^{\text{phys}} = \mathcal{O}^{\text{uni}} \left(1 + \alpha_{\mathcal{O}} \frac{m_s^{\text{phys}} - m_s^{\text{uni}}}{m_s^{\text{phys}}} \right)$$


- Calculate 2 values of m_s on C1 and M1



Strange Quark Mass Correction

$$\mathcal{O}^{\text{phys}} = \mathcal{O}^{\text{uni}} \left(1 + \alpha_{\mathcal{O}} \frac{m_s^{\text{phys}} - m_s^{\text{uni}}}{m_s^{\text{phys}}} \right)$$


- Calculate 2 values of m_s on C1 and M1
- Extrapolate to F1 masses



Strange Quark Mass Correction


$$\mathcal{O}^{\text{phys}} = \mathcal{O}^{\text{uni}} \left(1 + \alpha_{\mathcal{O}} \frac{m_s^{\text{phys}} - m_s^{\text{uni}}}{m_s^{\text{phys}}} \right)$$

- Calculate 2 values of m_s on C1 and M1
- Extrapolate to F1 masses
- Extrapolate to F1 lattice spacing



f_{D_s}/f_D Global Fit


$$\mathcal{O}(a, m_I, m_h) = \mathcal{O}(0, m_I^{\text{phys}}, m_h^{\text{phys}}) + C_{CL}^0 a^2 + C_\chi^0 \Delta m_\pi^2 + C_{P_h}^0 \Delta m_{P_h}^{-1}$$



$$f_{D_s}/f_D = 1.167(8)_{\text{stat}}$$

f_{D_s}/f_D Systematics

m_π and m_h , dependence




$$f_{D_s}/f_D = 1.167(8)_{\text{stat}} \left({}^{+4}_{-2} \right)_{\text{fit}}$$

To Do: FV, Isospin, fully correlated fits, m_{η_c} -disconnected and sea-charm

Φ_D


$$\begin{aligned} \mathcal{O}(a, m_I, m_h) &= \mathcal{O}(0, m_I^{\text{phys}}, m_h^{\text{phys}}) + C_{CL}^0 a^2 \\ &\quad + \left(C_\chi^0 + C_\chi^1 \Delta m_{P_h}^{-1} \right) \Delta m_\pi^2 + C_{P_h}^0 \Delta m_{P_h}^{-1} \end{aligned} \quad (2)$$



$$\Phi_D = 0.2857(44)_{\text{stat}} \text{GeV}^{3/2}$$

Φ_{D_s}


$$\begin{aligned}\mathcal{O}(a, m_I, m_h) &= \mathcal{O}(0, m_I^{\text{phys}}, m_h^{\text{phys}}) + \left(C_{CL}^0 + C_{CL}^1 \Delta m_{P_h}^{-1} \right) a^2 \\ &\quad + C_\chi^0 \Delta m_\pi^2 + C_{P_h}^0 \Delta m_{P_h}^{-1}\end{aligned}\quad (3)$$



$$\Phi_{D_s} = 0.3495(26)_{\text{stat}} \text{GeV}^{3/2}$$

comparison with FLAG (decay constants)


adapted from FLAG 2016 (arXiv:1607.00299)



PRELIMINARY and STATISTICAL ERROR ONLY

V_{cd} and V_{cs} comparison with literature

adapted from FLAG 2016
 (arXiv:1607.00299)
 using data from arXiv:1509.02220



PRELIM, STAT ERROR ONLY

Fit systematics:

- m_π^{cuts}
- ways to set m_c
- mass-dependent chiral and CL coeffs.

Not finalised yet

- Take all correlation into account
- FV
- Isospin
- sea-charm effects
- renormalisation

New set-up


Problems

- $am_h^{\max} < am_h^{\text{phys}}$ on Coarse
- Poor Signal-to-noise for $m_\pi = 139\text{MeV}$ and $m_h \gtrsim m_c^{\text{phys}}$

Solutions?

- Stout Smeared charm
- Gaussian Smearing for source + sink


Noise growth



“Smeared Runs”

WORK IN PROGRESS:

First Data (Limited statistic)



am_h reach

- Sweetspot: $M_5 = 1.0$
 3 hits of stout smearing
 $(\rho = 0.1)$
 $\Rightarrow am_h \lesssim 0.7$
- **Can reach charm** on all ensembles
- **Physical program:**
 f_D , f_{D_s} , semi-leptonics,
 bag parameters, ξ , HVP
- Ratio Method to reach b .

Summary

What we have done:

- D and D_s decay constants
- at **Physical Pion Masses** ($2 + 1f$ simulation) in an automatically $\mathcal{O}(a)$ -improved setting.
- Continuum Limit with 3 lattice spacings.
- Global fit with full control of systematics

Current work:

- Finalising systematic error budget
- Produce smeared data on other ensembles
- semi-leptonics, ratio method, combined analysis of all data

ADDITIONAL SLIDES

Definition of the Residual Mass

From the axial Ward identity we define the **residual mass** am_{res} as


$$am_{\text{res}} = \frac{\sum_x \langle J_{5q}(x) P(0) \rangle}{\sum_x \langle P(x) P(0) \rangle}$$

where J_{5q} is the pseudoscalar density in the centre of the 5th dimension.


⇒ measure of the residual chiral symmetry breaking

Quenched Study: M_5 and bound on am_h - quenched

m_{res} , am_h and M_5




Hints of flat continuum limit




JHEP 04 (2016) 037 (arXiv:1602.04118 [hep-lat])

$am_h \lesssim 0.4$ bound in 2+1f tested on C0



$am_h \lesssim 0.4$ bound still applies

Quenched Study: Flat continuum limit scaling



JHEP 04 (2016) 037 (arXiv:1602.04118 [hep-lat])


Definitions

- Decay constants: $\langle 0 | A_{cq}^\mu | D_q(p) \rangle = f_{D_q} p_{D_q}^\mu$
 \Rightarrow Extraction of $|V_{cd}|$ and $|V_{cs}|$
- Bag Parameters: $B_P = \frac{\langle P^0 | O_{VV+AA} | \bar{P}^0 \rangle}{8/3 f_P^2 m_P^2}$ where
 $O_{VV+AA} = (\bar{h}\gamma_\mu q)(\bar{h}\gamma_\mu q) + (\bar{h}\gamma_5\gamma_\mu q)(\bar{h}\gamma_5\gamma_\mu q)$
- ξ :
$$\xi = \frac{f_{hs}\sqrt{B_{hs}}}{f_{hl}\sqrt{B_{hl}}}$$


 \Rightarrow Extraction of $|V_{td}/V_{ts}|$

Limitations of current formalism

Residual mass




reach in m_h



Restriction $am_h \lesssim 0.4$ in agreement with quenched
 \Rightarrow **Cannot reach physical charm on the coarsest ensembles**


f_{D_s} and f_D Systematics

Φ_D



Can resolve C_χ^1

Φ_{D_s}



Can resolve C_{CL}^1

To Do:

- Renormalisation
- FV
- Isospin
- Fully correlated fits
- m_{η_c} -disconnected
- sea-charm


Smeared Kernel: Set-up

Smearing

Found sweetspot for


- $M_5 = 1.0$
 - 3 hits of stout smearing
 - Standard Stout parameter $\rho = 0.1$
 - Gaussian smearing of light and strange
 - Kernel smearing of heavy quarks
- ⇒ Now test the reach of am_h

hit comparison




Smeared Kernel

Residual Mass



am_h reach



- Can reach charm
- Limited statistic
- $am_h \lesssim 0.7$