



Contribution ID: 133

Type: **Talk**

## Landau Levels in Lattice QCD

*Friday, July 29, 2016 5:50 PM (20 minutes)*

The spectrum of the two-dimensional continuum Dirac operator in the presence of a uniform background magnetic field consists of Landau levels, which are degenerate and separated by gaps. On the lattice the Landau levels are spread out by discretization artefacts, but a remnant of their structure is clearly visible (Hofstadter butterfly). If one switches on a non-Abelian interaction, the butterfly structure will be smeared out, but the lowest Landau level will still be separated by a gap from the rest of the spectrum. In this talk we discuss how the eigenmodes of the four-dimensional QCD Dirac operator are built out of the two-dimensional eigenmodes of the Dirac operator diagonalized on each slice at fixed  $(z,t)$ . In particular, starting from this decomposition, we consider how well certain physical quantities are approximated by using only the two-dimensional eigenmodes belonging to the lowest Landau level.

**Primary authors:** Dr BRUCKMANN, Falk (University of Regensburg); Dr PITTLER, Ferenc (Eötvös University); Dr ENDRODI, Gergely (University of Frankfurt); Mr WELLNHOFER, Jacob (Universität Regensburg); Dr GIORDANO, Matteo (Eotvos University, Budapest); Prof. KATZ, Sándor (Eötvös University, Budapest); Prof. KOVÁCS, Tamás (MTA Atomki, Debrecen)

**Presenter:** Dr PITTLER, Ferenc (Eötvös University)

**Session Classification:** Nonzero Temperature and Density

**Track Classification:** Nonzero Temperature and Density