



Contribution ID: 138

Type: **Talk**

## Quark masses and strong coupling constant with Highly-Improved Staggered Quarks

*Tuesday, July 26, 2016 2:00 PM (20 minutes)*

We present new results on the charm and bottom quark masses as well as the strong coupling constant in 2+1-flavor QCD using Highly Improved Staggered Quark action and gauge configurations generated by HotQCD Collaboration.

Our approach is based on calculating the moments of meson correlators and we use a wide range of lattice spacing up to  $a^{-1} \sim 4.9$  GeV in our study.

The ratios of quark masses  $m_c/m_s$  and  $m_b/m_c$  are obtained from the combinations of the pseudo-scalar and vector meson masses and found to be  $m_c/m_s = 11.871(96)$  and  $m_b/m_c = 4.528(57)$  in the continuum limit.

The lattice results of the hyper-fine splitting of the charmonium can reproduce the experimental value in the continuum.

We also perform the determination of the strong coupling constant in the  $\overline{\text{MS}}$  scheme from the moments of pseudo-scalar charmonium correlators and find  $\alpha_s(\mu = m_c) = 0.3697(56)$ , which is the determination of  $\alpha_s$

at lowest energy scale so far.

For the charm quark mass we obtain  $m_c(\mu = m_c) = 1.2668(10)$  GeV.

**Primary author:** Dr MAEZAWA, Yu (YITP, Kyoto University)

**Co-author:** Dr PETRECZKY, Peter (Brookhaven National Laboratory)

**Presenter:** Dr MAEZAWA, Yu (YITP, Kyoto University)

**Session Classification:** Standard Model Parameters and Renormalization

**Track Classification:** Standard Model Parameters and Renormalization