

The Dark Side of the Propagators

Analytical approach to QCD
in the infrared of Minkowski space

Fabio Siringo

Department of Physics and Astronomy
University of Catania, Italy



LATTICE 2016 - Southampton, 25-30 July 2016

Overview: the whole talk in a slide



Overview: the whole talk in a slide

- The Dream:

Overview: the whole talk in a slide

- The Dream:
Do all the work analytically and forget about Lattice People!

Overview: the whole talk in a slide

- The Dream:
Do all the work analytically and forget about Lattice People!

Change the expansion point \rightarrow Massive Expansion
also known as Optimized Perturbation Theory (OPT) and
evaluate everything from first principles



Overview: the whole talk in a slide

- The Dream:
Do all the work analytically and forget about Lattice People!

Change the expansion point \rightarrow Massive Expansion
also known as Optimized Perturbation Theory (OPT) and
evaluate everything from first principles
- Reality:
Not self-consistent yet \rightarrow Optimization by Lattice



Overview: the whole talk in a slide

- The Dream:
Do all the work analytically and forget about Lattice People!

Change the expansion point \rightarrow Massive Expansion
also known as Optimized Perturbation Theory (OPT) and
evaluate everything from first principles

- Reality:
Not self-consistent yet \rightarrow Optimization by Lattice

but

we can analytically continue to Minkowski space!



Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR
(Tissier+Wschebor,2010,2011 - Landau Gauge)

Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR
(Tissier+Wschebor,2010,2011 - Landau Gauge)
- OPT is an old story: $(\mathcal{L}_0 + \delta\mathcal{L}_m) + (\mathcal{L}_{int} - \delta\mathcal{L}_m)$

Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR
(Tissier+Wschebor,2010,2011 - Landau Gauge)
- OPT is an old story: $(\mathcal{L}_0 + \delta\mathcal{L}_m) + (\mathcal{L}_{int} - \delta\mathcal{L}_m)$
- [Massive propagator] + [mass counterterm $\delta\Gamma = m^2$]



Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR
(Tissier+Wschebor,2010,2011 - Landau Gauge)
- OPT is an old story: $(\mathcal{L}_0 + \delta\mathcal{L}_m) + (\mathcal{L}_{int} - \delta\mathcal{L}_m)$
- [Massive propagator] + [mass counterterm $\delta\Gamma = m^2$]

Nice features:



Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR
(Tissier+Wschebor,2010,2011 - Landau Gauge)
- OPT is an old story: $(\mathcal{L}_0 + \delta\mathcal{L}_m) + (\mathcal{L}_{int} - \delta\mathcal{L}_m)$
- [Massive propagator] + [mass counterterm $\delta\Gamma = m^2$]

Nice features:

- The mass kills itself at tree-level:

$$\Sigma_{tree} = \text{---}\times\text{---} = \delta\Gamma = m^2 \quad \Longrightarrow \quad (-p^2 + m^2) - \Sigma = -p^2$$



Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR (Tissier+Wschebor,2010,2011 - Landau Gauge)
- OPT is an old story: $(\mathcal{L}_0 + \delta\mathcal{L}_m) + (\mathcal{L}_{int} - \delta\mathcal{L}_m)$
- [Massive propagator] + [mass counterterm $\delta\Gamma = m^2$]

Nice features:

- The mass kills itself at tree-level:

$$\Sigma_{tree} = \text{---}\times\text{---} = \delta\Gamma = m^2 \quad \implies \quad (-p^2 + m^2) - \Sigma = -p^2$$

- Mass divergences cancel in loops:

$$\text{---}\bigcirc\text{---} + \text{---}\bigcirc\times\text{---} = \text{IR finite}$$



Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR (Tissier+Wschebor,2010,2011 - Landau Gauge)
- OPT is an old story: $(\mathcal{L}_0 + \delta\mathcal{L}_m) + (\mathcal{L}_{int} - \delta\mathcal{L}_m)$
- [Massive propagator] + [mass counterterm $\delta\Gamma = m^2$]

Nice features:

- The mass kills itself at tree-level:

$$\Sigma_{tree} = \text{---}\times\text{---} = \delta\Gamma = m^2 \quad \implies \quad (-p^2 + m^2) - \Sigma = -p^2$$

- Mass divergences cancel in loops:

$$\text{---}\bigcirc\text{---} + \text{---}\bigcirc\times\text{---} = \text{IR finite}$$

- The original Lagrangian is not modified



Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR (Tissier+Wschebor,2010,2011 - Landau Gauge)
- OPT is an old story: $(\mathcal{L}_0 + \delta\mathcal{L}_m) + (\mathcal{L}_{int} - \delta\mathcal{L}_m)$
- [Massive propagator] + [mass counterterm $\delta\Gamma = m^2$]

Nice features:

- The mass kills itself at tree-level:

$$\Sigma_{tree} = \text{---}\times\text{---} = \delta\Gamma = m^2 \implies (-p^2 + m^2) - \Sigma = -p^2$$

- Mass divergences cancel in loops:

$$\text{---}\bigcirc\text{---} + \text{---}\bigcirc\times\text{---} = \text{IR finite}$$

- The original Lagrangian is not modified
- Standard UV behavior



Breaking the Dogma:

- Perturbation Theory (PT) works well in the IR (Tissier+Wschebor,2010,2011 - Landau Gauge)
- OPT is an old story: $(\mathcal{L}_0 + \delta\mathcal{L}_m) + (\mathcal{L}_{int} - \delta\mathcal{L}_m)$
- [Massive propagator] + [mass counterterm $\delta\Gamma = m^2$]

Nice features:

- The mass kills itself at tree-level:

$$\Sigma_{tree} = \text{---}\times\text{---} = \delta\Gamma = m^2 \implies (-p^2 + m^2) - \Sigma = -p^2$$

- Mass divergences cancel in loops:

$$\text{---}\bigcirc\text{---} + \text{---}\bigcirc\times\text{---} = \text{IR finite}$$

- The original Lagrangian is not modified
- Standard UV behavior
- From first principles



Do mass divergences cancel at any order?

A simple argument:

- Mass divergences arise from the massive propagator
- No mass divergences in the exact (scaleless) theory
- The Lagrangian is not modified



Do mass divergences cancel at any order?

A simple argument:

- Mass divergences arise from the massive propagator
- No mass divergences in the exact (scaleless) theory
- The Lagrangian is not modified

The counterterms $\delta\Gamma = m^2$ must cancel the divergences.
How many do we need?




Do mass divergences cancel at any order?

A simple argument:

- Mass divergences arise from the massive propagator
- No mass divergences in the exact (scaleless) theory
- The Lagrangian is not modified

The counterterms $\delta\Gamma = m^2$ must cancel the divergences.
How many do we need?



$$\frac{1}{-p^2 + m^2} \implies \frac{1}{-p^2 + m^2} m^2 \frac{1}{-p^2 + m^2}$$

The integral is less divergent at each insertion.
A finite number of insertions makes any loop integral convergent:




Do mass divergences cancel at any order?

A simple argument:

- Mass divergences arise from the massive propagator
- No mass divergences in the exact (scaleless) theory
- The Lagrangian is not modified

The counterterms $\delta\Gamma = m^2$ must cancel the divergences.
How many do we need?



A Feynman diagram consisting of a horizontal line on the left that enters a circle. From the right side of the circle, a horizontal line exits. A small 'x' is drawn on the right side of the circle, indicating a counterterm insertion.

$$\frac{1}{-p^2 + m^2} \implies \frac{1}{-p^2 + m^2} m^2 \frac{1}{-p^2 + m^2}$$

The integral is less divergent at each insertion.
A finite number of insertions makes any loop integral
convergent: divergences must cancel at a finite order



One-Loop third-order double expansion

(Landau Gauge)

Yang-Mills \rightarrow F.S., Nucl. Phys. B **907** 572 (2016)

QCD \rightarrow F.S., arXiv:1607.02040

$$\Sigma_{gh} = - \text{diagram} + \text{diagram}$$

The first diagram is a dashed line with a gluon loop on top. The second diagram is the same as the first, but with a large 'X' over the loop.

$$\Pi = \text{diagram} + \text{diagram} + \text{diagram} + \text{diagram} + \text{diagram} + \text{diagram} + \text{diagram} + \text{diagram} + \text{diagram}$$

The diagrams are: 1. A gluon loop with a ghost loop on top, marked with an 'X'. 2. A gluon loop with a ghost loop on top. 3. A gluon loop with a ghost loop on top, marked with an 'X'. 4. A gluon loop with a ghost loop on top, marked with two 'X's. 5. A ghost loop with a gluon loop on top. 6. A gluon loop with a ghost loop on top, marked with an 'X'. 7. A gluon loop with a ghost loop on top. 8. A gluon loop with a ghost loop on top, marked with an 'X'. 9. A ghost loop with a gluon loop on top.

$$\Sigma_q = \text{diagram} + \text{diagram} + \text{diagram} + \text{diagram}$$

The diagrams are: 1. A solid line with a ghost loop on top, marked with an 'X'. 2. A solid line with a gluon loop on top. 3. A solid line with a gluon loop on top, marked with an 'X'. 4. A solid line with a gluon loop on top, marked with an 'X'.



UNIVERSAL SCALING

Ignoring RG effects, $\alpha \sim N\alpha_s$

$$\Sigma(p) = \alpha \Sigma^{(1)}(p) + \alpha^2 \Sigma^{(2)}(p, N) + \dots \quad (1)$$

$$\frac{\Sigma(p)}{\alpha p^2} = -F(p^2/m^2) + \mathcal{O}(\alpha); \quad F(p^2/m^2) = -\frac{\Sigma^{(1)}}{p^2} \quad (2)$$

$$\Delta(p) = \frac{Z}{p^2 - \Sigma(p)} = \frac{J(p)}{p^2} \quad (3)$$

Setting $Z = z(1 + \alpha\delta Z)$ (one-loop):

$$z J(p)^{-1} = 1 + \alpha [F(p^2/m^2) - \delta Z] + \mathcal{O}(\alpha^2) \quad (4)$$


$$z J(p)^{-1} = 1 + \alpha [F(p^2/m^2) - F(\mu^2/m^2)] + \mathcal{O}(\alpha^2) \quad (5)$$

Must exist x, y, z :

$$\boxed{z J(p/x)^{-1} + y = F(p^2/m^2) + F_0 + \mathcal{O}(\alpha)} \quad (6)$$


UNIVERSAL SCALING

GLUON INVERSE DRESSING FUNCTION



UNIVERSAL SCALING


GLUON INVERSE DRESSING FUNCTION



UNIVERSAL SCALING

GHOST INVERSE DRESSING FUNCTION

Denoting by $G(s)$ the ghost universal function ($F(s) \rightarrow G(s)$)




UNIVERSAL SCALING

GHOST INVERSE DRESSING FUNCTION

The ghost universal function is just

$$G(s) = \frac{1}{12} \left[2 + \frac{1}{s} - 2s \log s + \frac{1}{s^2} (1+s)^2 (2s-1) \log(1+s) \right]$$



UNIVERSAL SCALING

GHOST INVERSE DRESSING FUNCTION

The ghost universal function is just

$$G(s) = \frac{1}{12} \left[2 + \frac{1}{s} - 2s \log s + \frac{1}{s^2} (1+s)^2 (2s-1) \log(1+s) \right]$$




TABLE of OPTIMIZED RENORMALIZATION

CONSTANTS:

$$z J(p/x)^{-1} + y = F(p^2/m^2) + F_0$$

arXiv:1607.02040

Data set	N	N_f	x	y	z	y'	z'
Bogolubsky et al.	3	0	1	0	3.33	0	1.57
Duarte et al.	3	0	1.1	-0.146	2.65	0.097	1.08
Cucchieri-Mendes	2	0	0.858	-0.254	1.69	0.196	1.09
Ayala et al.	3	0	0.933	-	-	0.045	1.17
Ayala et al.	3	2	1.04	-	-	0.045	1.28
Ayala et al.	3	4	1.04	-	-	0.045	1.28

Table: Scaling constants x, y, z (gluon) and y', z' (ghost). The constant shifts $F_0 = -1.05$, $G_0 = 0.24$ and the mass $m = 0.73$ GeV are optimized by requiring that $x = 1$ and $y = y' = 0$ for the lattice data of Bogolubsky et al. (2009)




Running Coupling

Pure Yang-Mills SU(3)

RG invariant product (Landau Gauge – MOM-Taylor scheme):

$$\alpha_s(\mu) = \alpha_s(\mu_0) \frac{J(\mu)\chi(\mu)^2}{J(\mu_0)\chi(\mu_0)^2}$$

What if $\delta F_0 = \delta G_0 = \pm 25\%$?



$\mu_0 = 2$ GeV, $\alpha_s = 0.37$, data of Bogolubsky et al.(2009).




Running Coupling

Pure Yang-Mills SU(3)

RG invariant product (Landau Gauge – MOM-Taylor scheme):

$$\alpha_s(\mu) = \alpha_s(\mu_0) \frac{J(\mu)\chi(\mu)^2}{J(\mu_0)\chi(\mu_0)^2}$$

What if $\delta F_0 = \delta G_0 = \pm 25\%$?



$\mu_0 = 0.15$ GeV, $\alpha_s = 0.2$, data of Bogolubsky et al.(2009).




Running Coupling

Pure Yang-Mills SU(3)

RG invariant product (Landau Gauge – MOM-Taylor scheme):

$$\alpha_s(\mu) = \alpha_s(\mu_0) \frac{J(\mu)\chi(\mu)^2}{J(\mu_0)\chi(\mu_0)^2}$$

What if $\delta F_0 = \delta G_0 = \pm 25\%$?



$\mu_0 = 0.67$ GeV, $\alpha_s = 1.21$, data of Bogolubsky et al.(2009).



FAQ I

SILLY POINTS MADE BY THE REFEREES

(more serious list at the end)



FAQ I

SILLY POINTS MADE BY THE REFEREES

(more serious list at the end)

- Does the photon acquire a mass by the same method?

FAQ I

SILLY POINTS MADE BY THE REFEREES

(more serious list at the end)

- Does the photon acquire a mass by the same method?
- How can you get anything new by adding "zero" to the Lagrangian ? (without inserting any physical ansatz or any model for NP physics)

FAQ I

SILLY POINTS MADE BY THE REFEREES

(more serious list at the end)

- Does the photon acquire a mass by the same method?
- How can you get anything new by adding "zero" to the Lagrangian ? (without inserting any physical ansatz or any model for NP physics)
- The method must be wrong otherwise it would also explain Chiral Symmetry Breaking

FAQ I

SILLY POINTS MADE BY THE REFEREES

(more serious list at the end)


- Does the photon acquire a mass by the same method?
- How can you get anything new by adding "zero" to the Lagrangian ? (without inserting any physical ansatz or any model for NP physics)
- The method must be wrong otherwise it would also explain Chiral Symmetry Breaking
- The method must be wrong otherwise the propagator could be analytically continued to Minkowski space where the gluon would get a physical dynamical pole



ANALYTIC CONTINUATION

arXiv:1605.07357

GLUON PROPAGATOR - SU(3)




Lattice data are from Bogolubsky et al. (2009)



ANALYTIC CONTINUATION

arXiv:1605.07357

GLUON PROPAGATOR - SU(3)




Lattice data are from Bogolubsky et al. (2009)



ANALYTIC CONTINUATION

arXiv:1605.07357



GENERALIZED SPECTRAL FUNCTION

HOW TO DEFINE A SPECTRAL FUNCTION WITH COMPLEX POLES ?

If $G(p)$ has complex poles then

$$G(p^2) = G^R(p^2) + \delta G(p^2)$$

where the *rational* function G^R just contains the poles

$$G^R(z) = \frac{R}{z - \alpha - i\beta} + \frac{R^*}{z - \alpha + i\beta}$$

and the finite part δG satisfies usual dispersion relations


$$\text{Re } \delta G(p^2) = PV \int_0^{+\infty} \frac{\rho(\omega)}{p^2 - \omega} d\omega$$

$$\rho(\omega) = -\frac{1}{\pi} \text{Im } \delta G(\omega + i\epsilon) = -\frac{1}{\pi} \text{Im } G(\omega + i\epsilon)$$

$G^R(p^2)$ cannot be reconstructed from $\text{Im } G$

ANALYTIC CONTINUATION

Dispersion relations with complex poles \rightarrow arXiv:1606.03769




Lattice data are from Bogolubsky et al. (2009)

BACK TO EUCLIDEAN SPACE

$$G^R(z) = \frac{R}{z - \alpha - i\beta} + \frac{R^*}{z - \alpha + i\beta} \Rightarrow \frac{p_E^2 + (\alpha + t\beta)}{p_E^4 + 2\alpha p_E^2 + (\alpha^2 + \beta^2)}$$

where $t = (\text{Im } R)/(\text{Re } R) = \tan[\arg(R)]$


RGZ model!



ANALYTIC CONTINUATION


Ghost dressing function: $\mathcal{G}(p^2) = \frac{\chi(p^2)}{p^2}$

$$\rho(p^2) = -\frac{1}{\pi} \text{Im} \mathcal{G}(p^2 + i\varepsilon) = \chi(0) \delta(p^2) - \frac{1}{\pi} \frac{\text{Im} \chi(p^2)}{p^2}$$



ANALYTIC CONTINUATION



Ghost dressing function: $\mathcal{G}(p^2) = \frac{\chi(p^2)}{p^2}$



Lattice data are from Bogolubsky et al. (2009)



Optimized by the Lattice $N_f = 2, m = 0.8 \text{ GeV} \quad M = ?$



Lattice data are for two light quarks, from Ayala et al. (2012)

What about poles ?

2 pairs of complex conjugated poles




CHIRAL QCD

Gluon sector

Optimized by the Lattice:

$m = 0.8 \text{ GeV}, M = 0.65 \text{ GeV}$

$m_1^2 = (0.54 \pm 0.52i) \text{ GeV}^2, \quad m_2^2 = (1.69 \pm 0.1i) \text{ GeV}^2$





$$\Sigma_q = \text{---} \mathbf{X} \text{---} + \text{---} \text{---} + \text{---} \mathbf{X} \text{---} + \text{---} \mathbf{X} \text{---}$$

- The counterterm $\delta\Gamma = -M$ cancels the mass at tree-level
- A massive propagator from *loops* $\rightarrow S(p) = \frac{Z(p)}{\not{p} - M(p)}$
- A new parameter $x = M/m$

but





$$\Sigma_q = \text{---} \times \text{---} + \text{---} \text{---} + \text{---} \times \text{---} + \text{---} \times \text{---}$$

- The counterterm $\delta\Gamma = -M$ cancels the mass at tree-level
- A massive propagator from *loops* $\rightarrow S(p) = \frac{Z(p)}{p-M(p)}$
- A new parameter $x = M/m$


but

- Agreement not as good as for pure YM theory ($Z(p)$ is decreasing)
- $M(p)$ depends on α_s
- Optimization is not easy without RG corrections!




CHIRAL QCD

Quark sector – $N_f = 2$



CHIRAL QCD

Quark sector – $N_f = 2$




Lattice data are: *unquenched*, $N_f = 2$, in the CHIRAL limit
Bowman et al. (2005)



CHIRAL QCD

Quark sector – $N_f = 2$




Lattice data are: *unquenched*, $N_f = 2$, in the CHIRAL limit
Bowman et al. (2005)



CHIRAL QCD

Quark sector – $N_f = 2$



Lattice data are: *unquenched*, $N_f = 2$, in the CHIRAL limit
Bowman et al. (2005)



Quark propagator:

$$S(p) = S_p(p^2)\not{p} + S_M(p^2)$$

NO COMPLEX POLES \implies Standard Dispersion Relations



Quark propagator:

$$S(p) = S_p(p^2)\not{p} + S_M(p^2)$$

NO COMPLEX POLES \implies Standard Dispersion Relations

$$\rho_M(p^2) = -\frac{1}{\pi} \text{Im } S_M(p^2)$$

$$\rho_p(p^2) = -\frac{1}{\pi} \text{Im } S_p(p^2)$$

$$S(p) = \int_0^\infty dq^2 \frac{\rho_p(q^2)\not{p} + \rho_M(q^2)}{p^2 - q^2 + i\varepsilon}.$$



Quark propagator:

$$S(p) = S_p(p^2)\not{p} + S_M(p^2)$$

NO COMPLEX POLES \implies Standard Dispersion Relations

$$\rho_M(p^2) = -\frac{1}{\pi} \text{Im } S_M(p^2)$$

$$\rho_p(p^2) = -\frac{1}{\pi} \text{Im } S_p(p^2)$$

$$S(p) = \int_0^\infty dq^2 \frac{\rho_p(q^2)\not{p} + \rho_M(q^2)}{p^2 - q^2 + i\varepsilon}.$$

Positivity Conditions:


$$\rho_p(p^2) \geq 0,$$

$$p \rho_p(p^2) - \rho_M(p^2) \geq 0$$




CHIRAL QCD

Quark sector – $N_f = 2$




CHIRAL QCD

Quark sector – $N_f = 2$




CHIRAL QCD

Quark sector – $N_f = 2$



CHIRAL QCD

Quark sector: $N_f = 2$, $M = 0.65$ GeV, $m = 0.7$ GeV




Positivity Conditions:

$$\rho_p(p^2) \geq 0, \quad p \rho_p(p^2) - \rho_M(p^2) \geq 0$$



CHIRAL QCD

Quark sector: $N_f = 2$, $M = 0.65$ GeV, $m = 0.7$ GeV



Positivity Conditions:

$$\rho_p(p^2) \geq 0, \quad p \rho_p(p^2) - \rho_M(p^2) \geq 0$$



FAQ II.

MORE SERIOUS LIST (that the Referees did not raise)



- What about the *Longitudinal Polarization* ?
The Landau gauge is very special!

FAQ II.

MORE SERIOUS LIST (that the Referees did not raise)

- What about the *Longitudinal Polarization* ?
The Landau gauge is very special!
- Can we *Optimize* the expansion (without Lattice Data) ?
May be by real observables, like glueball masses

FAQ II.

MORE SERIOUS LIST (that the Referees did not raise)

- What about the *Longitudinal Polarization* ?
The Landau gauge is very special!
- Can we *Optimize* the expansion (without Lattice Data) ?
May be by real observables, like glueball masses
- Is there any proof of renormalizability ?



FAQ II.

MORE SERIOUS LIST (that the Referees did not raise)

- What about the *Longitudinal Polarization* ?
The Landau gauge is very special!
- Can we *Optimize* the expansion (without Lattice Data) ?
May be by real observables, like glueball masses
- Is there any proof of renormalizability ?
- What about improving the expansion by RG ?



FAQ II.

MORE SERIOUS LIST (that the Referees did not raise)

- What about the *Longitudinal Polarization* ?
The Landau gauge is very special!
- Can we *Optimize* the expansion (without Lattice Data) ?
May be by real observables, like glueball masses
- Is there any proof of renormalizability ?
- What about improving the expansion by RG ?
- More questions and remarks are welcome!

THANK YOU

