



Contribution ID: 379

Type: **Talk**

Hadron Matrix Elements and the Feynman-Hellman Theorem

Monday, July 25, 2016 4:45 PM (20 minutes)

Motivated by the Feynman-Hellman Theorem, we develop an improved method for computing matrix elements of external currents utilizing only two-point correlation functions.

The contamination from excited states is shown to be Euclidean-time dependent allowing for a significantly improved ability to reliably determine and control the systematics.

We demonstrate the utility of our method with a calculation of the nucleon axial-charge, performed at a single lattice spacing and a moderate unphysical pion mass.

The Feynman-Hellman Theorem can be derived from the long Euclidean-time limit of correlation functions determined with functional derivatives of the partition function.

This elucidates the generic applicability of our new method: one can determine matrix elements of any external current by computing only two-point correlation functions, including non-zero momentum transfer and flavor-changing matrix elements.

Primary author: WALKER-LOUD, Andre (LBNL)

Co-authors: Dr BOUCHARD, Chris (William & Mary); Prof. ORGINOS, Kostas (College of William and Mary/JLAB); Dr KURTH, Thorsten (LBNL)

Presenter: WALKER-LOUD, Andre (LBNL)

Session Classification: Hadron Structure

Track Classification: Hadron Structure