

Contribution ID: 85 Type: Talk

Chiral Perturbation Theory at finite volume or with twisted boundary conditions

Thursday, 28 July 2016 14:00 (20 minutes)

Recent progress in mesonic Chiral Perturbation Theory at finite volume and/or with twisted boundary conditions will be discussed. Topics include the finite volume at two-loop order calculations of masses and decay constants for the pseudoscalar of the normal case for two and three flavours [1], partially quenched three flavour [2] and the case for different patterns of spontaneous symmetry breaking [3].

Twisted boundary conditions lead to extra terms in Ward identities and form-factors which affects finite volume corrections and relations often used. This will be discussed for a number of cases, the older work [4] and the application to $K_{\ell 3}$ for staggered partially quenched [5]. Vector two-point functions relevant for the g-2 HVP calculation will be treated at two-loop order [6] allowing a better estimate of twisting and finite volume corrections.

- [1] Finite Volume at two-loops in Chiral Perturbation Theory,
- J. Bijnens and T. Rössler,[arXiv:1411.6384] JHEP 1501 (2015) 034
- [2] Finite Volume for Three-Flavour Partially Quenched Chiral Perturbation Theory through NNLO in the Meson Sector,
- J. Bijnens and T. Rössler, [arXiv:1508.07238] JHEP 1511 (2015) 097
- [3] Finite Volume and Partially Quenched QCD-like Effective Field Theories,
- J. Bijnens and T. Rössler, [arXiv:1509.04082] JHEP 1511 (2015) 017 $\,$
- [4] Masses, Decay Constants and Electromagnetic Form-factors with Twisted Boundary Conditions, J. Bijnens and J. Relefors, [arXiv:1402.1385] JHEP 05 (2014) 015
- [5] C. Bernard, J. Bijnens, E. Gamiz and J. Relefors, work in progress
- [6] J. Bijnens and J. Relefors, work in progress

Primary author: Prof. BIJNENS, Johan (Lund University)

Presenter: Prof. BIJNENS, Johan (Lund University)

Session Classification: Weak Decays and Matrix Elements

Track Classification: Weak Decays and Matrix Elements