



Lattice simulations of vector mesons in strong magnetic field

Olga Solovjeva
in collaboration with E.V. Luschevskaya , O.V. Teryev

The 34th annual “International Symposium on Lattice Field Theory”
Highfield Campus, University of Southampton
24-30 July 2016

Details of calculations

For the generation of $SU(3)$ gauge configurations the tadpole improved Lüscher-Symanzik action was used

$$S = \beta_{imp} \sum_{plq} S_{plq} - \frac{\beta_{imp}}{20u_0^2} \sum_{rt} S_{rt},$$

where $S_{plq,rt} = (1/3)(1 - U_{plq,rt})$
 $u_0 = (W_{1 \times 1})^{1/4} = \langle (1/3)U_{plq} \rangle^{1/4}$ is the tadpole factor, calculated
at zero temperature (V.G.Bornyakov, E.-M.Ilgenfritz, and
M.Müller-Preussker, Phys. Rev. D 72, 054511 (2005)) .

Details of calculations

Parameters of the ensembles

| Ensemble | $N_t \times N_s^3$ | β_{imp} | $a,$ | N_{conf} | m_u, MeV |
|----------|--------------------|---------------|-------|------------|----------------------|
| A_{16} | 16^4 | 8.20 | 0.115 | 245 | 34.26 |
| A_{18} | 18^4 | 8.10 | 0.125 | 250-285 | 34.26, 17.13 , 11.99 |
| B_{18} | 18^4 | 8.20 | 0.115 | 200 | 34.26 |
| C_{18} | 18^4 | 8.30 | 0.105 | 235 | 34.26 |
| D_{18} | 18^4 | 8.45 | 0.095 | 195 | 34.26 |
| E_{18} | 18^4 | 8.60 | 0.084 | 180 | 34.26 |
| A_{20} | 20^4 | 8.20 | 0.115 | 275 | 34.26 |
| B_{20} | 20^4 | 8.45 | 0.095 | 195 | 34.26 |

Details of calculations

2-pt correlation function : $C = \langle O_1 O_2 \rangle_A$

Interpolating operators: $O_{1,2} = \psi^\dagger(x) \gamma_i \psi(x)$,

where $i = 1, 2, 3 \rightarrow x, y, z$ components respectively.

$$C^{VV}(s_z = \pm 1) = C_{xx}^{VV} + C_{yy}^{VV} \pm i(C_{xy}^{VV} - C_{yx}^{VV})$$

$$\langle \bar{\psi} \gamma_\mu \psi \bar{\psi} \gamma_\nu \psi \rangle_A = -[\gamma_\mu D^{-1}(x, y) \gamma_\nu D^{-1}(y, x)]$$

consider zero momentum $\langle p \rangle = 0$ for the ground energy state

$$C_{ij}^{VV}(n_t) = \langle \bar{\psi}(\mathbf{0}, n_t) \gamma_i \psi(\mathbf{0}, n_t) \bar{\psi}(\mathbf{0}, 0) \gamma_j \psi(\mathbf{0}, 0) \rangle_A =$$

$$\sum_k \langle 0 | \gamma_i | k \rangle \langle k | \gamma_j | 0 \rangle e^{-n_t a E_k}$$

Details of calculations

Solve Dirac equation numerically

$$D\psi_k = i\lambda_k \psi_k, \quad D = \gamma^\mu (\partial_\mu - iA_\mu)$$

We use the Neuberger overlap operator $D_{overlap}$


$$A_{\mu ij} \rightarrow A_{\mu ij} + A_\mu^B \delta_{ij}$$

$A_\mu^B(x) = \frac{B}{2}(x_1 \delta_{\mu,2} - x_2 \delta_{\mu,1})$ - Abelian magnetic field

Calculate the propagators

$$D^{-1}(x, y) = \sum_{k < M} \frac{\psi_k(x)\psi_k^\dagger(y)}{i\lambda_k + m}$$


Fitting the correlators



$$\tilde{C}_{fit}(n_t) = 2A_0 e^{-N_T a E_0 / 2} \cosh\left(\frac{N_T}{2} - n_t\right) a E_0$$

E_0 - ?

Lattice Stern–Gerlach experiment for the ρ mesons




Pointlike particle (lines): $E^2 = (1 + gs_z)qB + m^2$

Non-pointlike particle (dots):

$$E^2 = (1 + gs_z)qB + m^2 - 4\pi m\beta(qB)^2$$

Lack of the charged vector meson tachyonic mode



pointlike particle : $eB_c \approx 1 \text{ GeV}^2$

$$E^2 = |qB| - g s_z qB + m^2 - 4\pi m \beta (qB)^2 - 4\pi m \beta^h (qB)^4$$

The g-factor of ρ meson

| V_{latt} | $m_q(Mev)$ | $a(fm)$ | g -factor | $\chi^2/\text{d.o.f.}$ | N_{conf} |
|------------|---------------------|---------|-----------------|------------------------|------------|
| 18^4 | 11.99 | 0.115 | 2.02 ± 0.11 | 0.524 | 250 |
| 18^4 | 17.13 | 0.115 | 2.20 ± 0.12 | 0.766 | 250 |
| 18^4 | 34.26 | 0.115 | 2.10 ± 0.01 | 0.610 | 285 |
| 18^4 | 51.39 | 0.115 | 2.06 ± 0.13 | 2.363 | 250 |
| 18^4 | $m_q \rightarrow 0$ | 0.115 | 2.11 ± 0.10 | 0.805 | |
| 18^4 | 34.26 | 0.095 | 2.25 ± 0.08 | 1.102 | 200 |
| 20^4 | 34.26 | 0.115 | 2.04 ± 0.14 | 3.101 | 275 |

Experiment: $g_{exp} = 2.1 \pm 0.5$ D. G. Gudino and G. T. Sanchez (2015), Int.J. of Mod.Phys.A, 30:1550114 arXiv:1305.6345


Previous results:

Relativistic quark model : $g \approx 2.37$ A. M. Badalian, Yu. A. Simonov(ITEP), Phys. Rev. D 87, 074012 (2013)

QCD sum rules: $g = 2.4 \pm 0.4$ T. M. Aliev et al., Phys.Lett.B678

Lattice: $g \approx 3.25$ F.X.Lee et al., Phys.Rev. D,78,094502(2008)


Quark mass dependence of the g-factor of ρ meson



$$m_\rho = c_0 + c_1 \cdot m_u$$


$18^4, a = 0.115 \text{ fm}: m_q \rightarrow 0 \quad g = 2.11 \pm 0.10$

Lattice Shtern-Gerlach experiment for the charged K* meson




$$E^2 = (1 + gs_z)qB + m^2 - 4\pi m\beta(qB)^2 - 4\pi m(qB)^4$$

Lattice Shtern-Gerlach experiment for the charged K* meson



$$E^2 = (1 + g s_z) q B + m^2 - 4\pi m \beta (q B)^2 - 4\pi m (q B)^4$$

g-factor of charged K* meson



| m_u (Mev) | m_s/m_u | g-factor | $\chi^2/\text{d.o.f.}$ | N_{conf} |
|-------------|-----------|-----------------|------------------------|------------|
| 11.99 | 20 | 2.27 ± 0.18 | 1.845 | 250 |
| 11.99 | 25 | 2.23 ± 0.23 | 1.986 | 250 |
| 11.99 | 30 | 2.29 ± 0.19 | 1.366 | 250 |
| 17.13 | 20 | 2.43 ± 0.24 | 3.282 | 250 |

Conclusions:

- Splitting of ground state energy of the ρ and K^* mesons depending on its spin projection on the axis of the external magnetic field
- Lack of the tachyonic mode of charged ρ mesons
- g-factor of ρ meson has been estimated in the chiral limit
- g-factor of K^* meson has been established (extrapolations in future)


Thank you for your attention!
olga.solovjeva@itep.ru




Experiment

Data from BaBar Collaboration for $e^+e^- \rightarrow \pi^+\pi^-2\pi^0$
Obtained value: $g_{exp} = 2.1 \pm 0.5$

Extrapolation $V_{phys} \rightarrow \infty$ for ρ meson




Extrapolation $m_q \rightarrow 0$ for ρ meson




$m_q \rightarrow 0$: $\rho(770)$ -meson $m_\rho = 839 \pm 11 \text{ Mev}$

Unpolarized neutral ρ meson



Lattice spacing dependence $\beta_m^{|s_z|=1}(\rho^0)$



$$\beta_m^{|s_z|=1}(\rho^0) = (-0.0235 \pm 0.0023) \text{ Gev}^{-3}$$


$$\beta_m^{|s_z|=1}(\rho^0) = (-1.86 \pm 0.18) \text{ } 10^{-4} \text{ fm}^3$$

Magnetic polarizability of the ρ^0 meson with nonzero spin

| V_{latt} | $a(fm)$ | $\beta_m^{m_q=34MeV} (Gev^{-3})$ | Error (Gev^{-3}) | $\chi^2/d.o.f.$ |
|------------|---------------|----------------------------------|----------------------|-----------------|
| 18^4 | 0.084 | -0.0169 | 0.0012 | 1.235 |
| 18^4 | 0.095 | -0.0158 | 0.0009 | 0.730 |
| 18^4 | 0.115 | -0.0133 | 0.0008 | 0.754 |
| 18^4 | 0.125 | -0.0135 | 0.0006 | 0.832 |
| 18^4 | $a = 0$ extr. | -0.0235 | 0.0023 | 0.561 |
| | | $\beta_m^{ch. extr} (Gev^{-3})$ | | |
| 18^4 | 0.115 | -0.0138 | 0.0005 | 2.648 |
| 18^4 | 0.125 | -0.0161 | 0.0025 | 23.862 |

Table : The values of magnetic polarizability of the vector ρ^0 meson with nonzero spin for the bare quark mass $m_q = 34.26 Mev$, lattice volume 18^4 and various lattice spacings.

Extrapolation $m_q \rightarrow 0$ for K^* meson



$m_q \rightarrow 0$: $K^*(895)$ -meson $m_{K^*} = 894 \pm 12 MeV$