



Contribution ID: 12

Type: **Talk**

## **Critical endline of the finite temperature phase transition for 2+1 flavor QCD around the SU(3)-flavor symmetric point**

*Monday, 25 July 2016 15:35 (20 minutes)*

We investigate the critical endline of the finite temperature phase transition of QCD around the SU(3)-flavor symmetric point at zero chemical potential.

We employ the renormalization-group improved Iwasaki gauge action and non-perturbatively  $O(a)$ -improved Wilson-clover fermion action.

The critical endline is determined by using the intersection point of kurtosis, employing the multi-parameter, multi-ensemble reweighting method to calculate observables off the SU(3)-symmetric point, at the temporal size  $N_T=6$  and lattice spacing as low as  $a \approx 0.19$  fm.

We confirm that the slope of the critical endline takes the value of  $-2$ , and find that the second derivative is positive, at the SU(3)-flavor symmetric point on the Columbia plot parametrized with the strange quark mass  $m_s$  and degenerated up-down quark mass  $m_l$ .

**Primary author:** Dr NAKAMURA, Yoshifumi (RIKEN)

**Co-authors:** Dr UKAWA, Akira (RIKEN); Dr TAKEDA, Shinji (Kanazawa University); Prof. KURAMASHI, Yoshinobu (University of Tsukuba)

**Presenter:** Dr NAKAMURA, Yoshifumi (RIKEN)

**Session Classification:** Nonzero Temperature and Density

**Track Classification:** Nonzero Temperature and Density