



Contribution ID: 207

Type: Talk

Temperature dependence of topological susceptibility using gradient flow

Friday, July 29, 2016 5:10 PM (20 minutes)

Topological susceptibility brings important information of the QCD vacuum and contributes to a production of the axion. The temperature dependence of the topological susceptibility plays a crucial role in its production rate in the early universe.

It is a test whether the axion can be an appropriate candidate of the dark matter.

In this talk we report our recent study on the topological susceptibility in the finite temperature QCD with $N_f=2+1$ flavors.

The topological susceptibility is given by applying the gradient flow to the finite temperature system. We cover a wide range of temperature region $174 \leq T \leq 697$ MeV.

The u and d quark mass is rather heavy $m_\pi/m_\rho = 0.63$ and s quark mass is set to almost the physical value.

Primary author: Dr TANIGUCHI, Yusuke (University of Tsukuba)

Co-authors: Prof. SUZUKI, Hiroshi (Kyushu University); Prof. KANAYA, Kazuyuki (CiRFSE, Univ. Tsukuba); Dr KITAZAWA, Masakiyo (Osaka University); Mr WAKABAYASHI, Naoki (Niigata University); Mr IWAMI, Ryo (Niigata University); Dr EJIRI, Shinji (Niigata University); Dr UMEDA, Takashi (Hiroshima Univ.)

Presenter: Dr TANIGUCHI, Yusuke (University of Tsukuba)

Session Classification: Nonzero Temperature and Density

Track Classification: Nonzero Temperature and Density