



Contribution ID: 69

Type: Poster

Isovector Axial Charge with Current Improvement

Tuesday, July 26, 2016 8:00 PM (1 hour)

We employ dimension 4 operators to improve the local vector and axial vector currents, as the leading order approximation of the lattice conserved current, and then calculate the nucleon iso-vector axial coupling g_A^3 using overlap valence on Domain Wall Fermion sea. Using the equality of g_A^3 from A_i and A_4 components of the axial-vector current as a normalization condition in addition to axial Ward identity, we find two to three percent increase of g_A^3 towards the experimental value. The excited state contamination has been taken into account with three time separations between the source and sink. The improved axial charges $g_A^{IM}(24I) = 1.188(7)$, $g_A^{IM}(32I) = 1.177(9)$ are obtained on $24^3 \times 64$ and $32^3 \times 64$ lattices at the unitary point where the pion masses are 330 MeV and 300 MeV respectively.

Primary author: Dr LIANG, Jian (University of Kentucky)

Co-authors: Prof. ALEXANDRU, Andrei (The George Washington University); Prof. LIU, Keh-Fei (University of Kentucky); SUFIAN, Raza (University of Kentucky, United States); Prof. DRAPER, Terrence (University of Kentucky); Dr YANG, YIBO (university of Kentucky)

Presenter: Dr YANG, YIBO (university of Kentucky)

Session Classification: Poster

Track Classification: Hadron Structure