



Contribution ID : 160

Type : Talk

## The QCD deconfinement critical point as a function of $N_t$ with $N_f=2$ flavours of unimproved Wilson fermions

*Monday, 25 July 2016 16:45 (20)*

QCD at zero baryon density in the limit of infinite quark mass undergoes a first order deconfinement phase transition at a critical temperature  $T_c$  corresponding to the breaking of the global centre symmetry.

In the presence of dynamical quarks the global centre symmetry is explicitly broken. Lowering the quark mass the first order phase transition weakens and terminates in a second order  $Z(2)$  point. Beyond this line confined and deconfined regions are analytically connected by a crossover transition.

As the continuum limit is approached (i.e. the lattice spacing is decreased) the region of first order transitions expands towards lower masses. We study the deconfinement critical point with standard Wilson fermions and  $N_f=2$  flavours. Therefore we simulate several  $\kappa$  values on  $N_t=8$  and various aspect ratios in order to extrapolate to the thermodynamic limit, applying finite size scaling. We estimate if and when a continuum extrapolation is possible.

**Primary author(s) :** Mr CZABAN, Christopher (Goethe University Frankfurt am Main)

**Co-author(s) :** Prof. PHILIPSEN, Owe (Goethe University Frankfurt am Main)

**Presenter(s) :** Mr CZABAN, Christopher (Goethe University Frankfurt am Main)

**Session Classification :** Nonzero Temperature and Density

**Track Classification :** Nonzero Temperature and Density