



Contribution ID: 357

Type: **Talk**

Spectrum and mass anomalous dimension of SU(2) gauge theories with fermions in the adjoint representation: from $N_f = 1/2$ to $N_f = 2$

Monday, July 25, 2016 3:35 PM (20 minutes)

In this work I will summarize our results concerning the spectrum and mass anomalous dimension of SU(2) gauge theories with a different number of fermions in the adjoint representation, where each Majorana fermion corresponds effectively to half a Dirac flavour N_f . The most relevant examples for the extensions of the standard model are supersymmetric Yang-Mills theory ($N_f = 1/2$) and minimal walking technicolour ($N_f = 2$). In addition to these theories I will also consider the cases of $N_f = 1$ and $N_f = 3/2$.

The results will contain the particle spectrum of glueballs, triplet and singlet mesons, and possible fractionally charged spin half particles. In addition I will discuss our recent results for the mass anomalous dimension.

Author: Dr BERGNER, Georg (AEC ITP University of Bern)

Co-authors: Prof. MÜNSTER, Gernot (University of Münster); Prof. MONTVAY, Istvan (DESY Hamburg); Dr PIEMONTE, Stefano (University of Regensburg); Dr GIUDICE, Pietro (University of Münster)

Presenter: Dr BERGNER, Georg (AEC ITP University of Bern)

Session Classification: Physics Beyond the Standard Model

Track Classification: Physics Beyond the Standard Model