



Contribution ID: 212

Type: Poster

Tuning of hopping parameters in Oktay-Kronfeld action for heavy quarks on the $N_f = 2 + 1 + 1$ MILC HISQ ensemble.

Tuesday, July 26, 2016 7:00 PM (1 hour)

We determine hopping parameters of the Oktay-Kronfeld (OK) action for charm and bottom quarks. We use $N_f = 2 + 1 + 1$ MILC HISQ ensembles (with $a \approx 0.12\text{fm}$, $am_l = 0.0102$, $am_s = 0.0509$ and $am_c = 0.635$). As a key ingredient, we compute the masses of pseudoscalar and vector mesons B_s^* , D_s^* and their hyperfine splittings; the valence light quark is simulated with HISQ action. We also monitor the inconsistency parameters to confirm the improvement.

Primary authors: Dr GUPTA, Rajan (Los Alamos National Lab); Mr PARK, Sungwoo (Seoul National University); Prof. LEE, Weonjong (Seoul National University); Dr JANG, Yong-Chull (Los Alamos National Laboratory)

Co-authors: Dr TANMOY, Bhattacharya (Los Alamos National Laboratory); Dr YOON, Boram (Los Alamos National Laboratory); Mr JEONG, Hwancheol (Seoul National University); Mr LEEM, Jaehoon (Seoul National University); Dr BAILEY, Jon (Seoul National University)

Presenter: Mr PARK, Sungwoo (Seoul National University)

Session Classification: Poster

Track Classification: Hadron Spectroscopy and Interactions