



Contribution ID: 280

Type: **Talk**

## **Nucleon form factors and couplings with $N_f = 2 + 1$ Wilson fermions**

*Thursday, July 28, 2016 4:50 PM (20 minutes)*

We present updated results on the isovector electromagnetic form factors and axial coupling of the nucleon calculated using the CLS ensembles with  $N_f = 2 + 1$  flavours of Wilson fermions. Systematic effects are investigated by covering a range of lattice spacings and pseudoscalar masses. Efficient variance reduction is achieved with the truncated solver method. The strategy to renormalize nucleon matrix elements with the Rome-Southampton method is discussed.

**Author:** Dr HARRIS, Tim (Helmholtz-Institut Mainz)

**Co-authors:** Dr DJUKANOVIC, Dalibor (Helmholtz-Institut Mainz); Dr VON HIPPEL, Georg (University of Mainz); Prof. WITTIG, Hartmut (University of Mainz); Prof. MEYER, Harvey (Johannes-Gutenberg University Mainz); Dr JUNNARKAR, Parikshit (Helmholtz-Institut Mainz)

**Presenter:** Dr HARRIS, Tim (Helmholtz-Institut Mainz)

**Session Classification:** Hadron Structure

**Track Classification:** Hadron Structure