



Contribution ID: 260

Type: **Talk**

Form factors in the $B_s \rightarrow K l \nu$ decays using HQET and the lattice

Friday, July 29, 2016 5:50 PM (20 minutes)

We report on a recent computation of the form factors in semi-leptonic decays of the B_s using Heavy Quark Effective Theory (HQET) formalism applied on the lattice. The connection of the form factors with the 2-point and 3-point correlators on the lattice is explained, and the subsequent non-perturbative renormalization of HQET and its matching to $N_f = 2$ QCD is outlined. The results of the (static) leading-order calculation in the continuum limit is presented. Preliminary results on extending the leading-order computation by including the $1/m$ effects (where m is the heavy-quark mass) are mentioned.

Primary author: Dr BANERJEE, Debasish (NIC, DESY)

Co-authors: Dr JOSEPH, Anosh (DAMTP, University of Cambridge); Dr BERNADONI, Fabio (TU Dresden); Dr BAHN, Felix Tobias (NIC DESY); Dr SIMMA, Hubert (NIC DESY); Dr KOREN, Mateusz (NIC, DESY); Prof. SOMMER, Rainer (NIC, DESY)

Presenter: Dr BANERJEE, Debasish (NIC, DESY)

Session Classification: Weak Decays and Matrix Elements

Track Classification: Weak Decays and Matrix Elements