



Contribution ID: 56

Type: **Talk**

## **V<sub>us</sub> from inclusive determinations based on hadronic tau data**

*Thursday, July 28, 2016 3:20 PM (20 minutes)*

We discuss the current status of the determination of  $V_{us}$  based on flavor-breaking finite energy sum rule analyses of non-strange and strange hadronic tau decay data, highlighting the role of lattice input in providing a means of investigating the reliability of the OPE representation for the relevant flavor-breaking polarization combination employed in this analysis, and in quantifying the associated theoretical errors on  $V_{us}$ . We present results for  $V_{us}$  based on a new implementation of this approach motivated by continuum and lattice studies and show that this resolves the long-standing problem of  $V_{us}$  values more than 3 sigma low compared to 3-family unitarity expectations obtained in earlier implementations of this approach. We also discuss the practical limitations of this new implementation given what near-term improvements are likely in the experimental strange decay distributions, and, motivated by these limitations, propose a new method for obtaining  $V_{us}$  which involves dispersive analyses of the strange hadronic tau decay data alone in which lattice results for the relevant combination of the flavor  $us$   $J=0$  and 1 vector and axial vector polarizations can be used in place of the OPE. Details of the practical implementation of this approach, as well as preliminary results for  $V_{us}$ , will be presented in the accompanying talk by Hiroshi Ohki.

**Author:** MALTMAN, Kim (York University)

**Co-authors:** OHKI, Hiroshi (BNL); ZANOTTI, James (University of Adelaide); LEWIS, Randy (York University); Dr HUDSPITH, Renwick (York University); IZUBUCHI, Taku (BNL)

**Presenter:** MALTMAN, Kim (York University)

**Session Classification:** Weak Decays and Matrix Elements

**Track Classification:** Weak Decays and Matrix Elements